

*Helena Haapsamo*

A FOLLOW-UP STUDY  
OF CHILDREN'S  
COMMUNICATIVE  
DEVELOPMENT

ASSOCIATIONS TO SOCIAL-EMOTIONAL AND  
BEHAVIOURAL PROBLEMS AND COMPETENCES  
AND EXPERIENCED MATERNAL STRESS

UNIVERSITY OF OULU GRADUATE SCHOOL;  
UNIVERSITY OF OULU,  
FACULTY OF MEDICINE,  
INSTITUTE OF CLINICAL MEDICINE, DEPARTMENT OF PAEDIATRICS;  
OULU UNIVERSITY HOSPITAL;  
FACULTY OF EDUCATION,  
DEPARTMENT OF EDUCATIONAL SCIENCES AND TEACHER EDUCATION;  
UNIVERSITY OF MASSACHUSETTS BOSTON,  
DEPARTMENT OF PSYCHIATRY

D  
MEDICA





ACTA UNIVERSITATIS OULUENSIS  
D Medica 1179

*HELENA HAAPSAMO*

**A FOLLOW-UP STUDY OF  
CHILDREN'S COMMUNICATIVE  
DEVELOPMENT**

Associations to social-emotional and behavioural  
problems and competences and experienced maternal  
stress

Academic dissertation to be presented with the assent  
of the Doctoral Training Committee of Health and  
Biosciences of the University of Oulu for public defence  
in Auditorium F101 of the Department of Physiology  
(Aapistie 7), on 19 November 2012, at 12 noon

UNIVERSITY OF OULU, OULU 2012

Copyright © 2012  
Acta Univ. Oul. D 1179, 2012

Supervised by  
Professor Irma Moilanen  
Professor Hannu Soini  
Professor Hanna Ebeling

Reviewed by  
Docent Jari Sinkkonen  
Professor Eeva Aronen

ISBN 978-951-42-9961-2 (Paperback)  
ISBN 978-951-42-9962-9 (PDF)

ISSN 0355-3221 (Printed)  
ISSN 1796-2234 (Online)

Cover Design  
Raimo Ahonen

JUVENES PRINT  
TAMPERE 2012

**Haapsamo, Helena, A Follow-up study of children's communicative development. Associations to social-emotional and behavioural problems and competences and experienced maternal stress**

University of Oulu Graduate School; University of Oulu, Faculty of Medicine, Institute of Clinical Medicine, Department of Paediatrics, Clinic of Child Psychiatry, P.O. Box 5000, FI-90014 University of Oulu, Finland; Oulu University Hospital, P.O. Box 10, FI-90029 OYS; Faculty of Education, Department of Educational Sciences and Teacher Education, P.O. Box 2000, FI-90014 University of Oulu, Finland; University of Massachusetts Boston, Department of Psychiatry, 100 Morrissey Blvd., Boston, MA 02125-3393, USA

*Acta Univ. Oul. D 1179, 2012*

Oulu, Finland

***Abstract***

The aim of this study was to follow 8 to 36- month old children's communicative development and its' associations with social-emotional skills (the Brief Infant Toddler Social-Emotional Assessment, BITSEA) and behavioural problems. This study is the first study using the Finnish version of the BITSEA.

A total of 50 children participated in the Oulu region (first phases at year 2006 and 2007).

At the age of 8 months (at year 2006, n = 31), child participants were grouped into two conditions: 1) children possibly needing support for deficiencies in communicative and interaction skills and 2) children without noted areas of need of support in communicative and interaction skills. Through random sampling, three groups were formed and included children from both the above mentioned conditions. The groups met fortnightly for five months for a directed song-play session (enrichment intervention).

The sample size was increased at year 2007 (n = 19) and follow-up questionnaires were sent to all of the participating families (N = 50) at the child's age of 18, 24 and 36 months. The sample during the year 2007 did not receive any intervention.

Results suggest, that the children's communicative and social-emotional development may be linked to each other. Children who scored higher in the assessments in communicative skills were also more successful on assessments measuring social interaction and social-emotional competence. Children with better communication skills demonstrated higher scores after enrichment-intervention.

Scores on the BITSEA demonstrated an association with other indicators of children's development employed in the study suggesting the utility of the BITSEA as a follow-up assessment in Finnish sample.

Mothers, who rated high maternal stress (measured at the child's age of 8 months) also rated higher levels of social-emotional and behavioural problems in their children, especially when a child was 18 months. This effect appeared to decrease over time.

Results indicate the clinical importance of directly measuring not only a child's linguistic and social-emotional development, but also including assessment of a child's immediate environment, such as parents and siblings.

***Keywords:*** behavioural, children, communication, maternal stress, parent-child interaction, social-emotional development



## **Haapsamo, Helena, Lapsen kielellinen kehitys ja sen yhteys tunne-elämään, käyttäytymiseen ja sosiaalisuuteen sekä äidin kokemaan stressiin. Seurantatutkimus**

Oulun yliopiston tutkijakoulu; Oulun yliopisto, Lääketieteellinen tiedekunta, Kliinisen lääketieteen laitos, Lastentaudit, Lastenpsykiatrian klinikka, PL 5000, 90014 Oulun yliopisto; Oulun yliopistollinen sairaala, PL 10, 90029 OYS; Kasvatustieteiden tiedekunta, Kasvatustieteiden ja opettajankoulutuksen yksikkö, PL 2000, 90014 Oulun yliopisto; Department of Psychiatry, University of Massachusetts Boston, 100 Morrissey Blvd., Boston, MA 02125-3393, USA

*Acta Univ. Oul. D 1179, 2012*

Oulu

### ***Tiivistelmä***

Tutkimuksen tarkoituksena oli seurata 8-36 kuukauden ikäisten lasten kielellistä kehitystä ja sen yhteyttä sosioemotionaaliseen kehitykseen (Brief Infant Toddler Social-Emotional Assessment, BITSEA-lomake) sekä käyttäytymisen ongelmiin. Tämä oli ensimmäinen tutkimus Suomessa, jossa käytettiin BITSEAn suomenkielistä versiota.

Tutkimukseen osallistui 50 perhettä (aloitus vuosina 2006 ja 2007) Oulun alueelta.

Kahdeksan kuukauden iässä täytettyjen lomakkeiden perusteella (vuonna 2006,  $n = 31$ ) lapset luokiteltiin vuorovaikutus- ja kommunikaatiotaitojen mukaan kahteen joukkoon: 1) Vuorovaikutus- ja kommunikaatiotaitojen tukea mahdollisesti tarvitsevat lapset ja 2) Lapset, joilla ei todennäköisesti ollut tuen tarvetta vuorovaikutus- ja kommunikaatiotaidoissa. Satunnaisotannalla muodostettiin kolme pienryhmää (rikastuttamisryhmät), joihin kuului sekä mahdollista tukea tarvitsevia että tukea tarvitsemattomia lapsia. Ryhmät kokoontuivat joka toinen viikko viiden kuukauden ajan ohjattuun laulu-leikkituokioon.

Tutkimusta laajennettiin syksyllä 2007 ( $n = 19$ ) ja kaikille tutkimukseen osallistuneille perheille ( $N = 50$ ) lähetettiin seurantalomakkeita, jotka vanhempien tuli täyttää lapsen ollessa 18, 24 ja 36 kuukauden ikäinen. Vuoden 2007 otoksen perheille ei tarjottu perheinterventiota.

Tutkimuksessa kävi ilmi, että lapset, jotka saivat korkeat pisteet sosioemotionaalista kompetenssia kuvaavissa kartoituksissa, menestyivät paremmin myös vuorovaikutus- ja kommunikaatiotaitoja kuvaavissa arvioinneissa. Lapset, joilla oli jo ennestään korkeammat pisteet vuorovaikutus- ja kommunikaatiotaitojen kuvaavissa lomakkeissa, näyttivät saavan korkeammat pisteet myös intervention jälkeen.

Tutkimuksessa tuli esiin, että BITSEA -lomake korreloi hyvin muiden lasten kehityksen seurannassa käytettyjen mittareiden kanssa ja soveltuu siten hyvin tutkimuslomakkeeksi suomalaisessakin aineistossa. Lisäksi äidit, jotka arvioivat stressitasonsa korkealle (lapsen ollessa kahdeksan kuukauden ikäinen), arvioivat myöhemmin myös korkeampia pisteitä lasten sosioemotionaalisten ja käyttäytymisen ongelmien kyselylomakkeissa lapsen ollessa 18 kuukauden ikäinen. Tämä vaikutus kuitenkin väheni lapsen kasvaessa.

Tutkimustulokset osoittavat, että perhe- ja neuvolatyössä on lapsen kehityksen kannalta tärkeää huomioida myös hänen kasvuympäristönsä; vanhempien hyvinvointi ja sisarusten määrä vaikuttavat lapsen kielelliseen ja sosioemotionaaliseen kehitykseen.

*Asiasanat:* kommunikaatio, käyttäytyminen, lapset, sosioemotionaalinen kehitys, vanhempi-lapsisuhteen vuorovaikutus, äidin stressi





***To my family: Risto, Veeti and Henrietta***

*“Paka, paka, pullaa”...*



## Acknowledgements

This follow-up study was conducted at the Clinic of Child Psychiatry at Oulu University, the University Hospital of Oulu, and at five Health Care Centres in Oulu with the collaboration of Department of Educational Sciences and Teacher Education. I would like to express my sincerest and warmest gratitude to each and everyone, who has participated or has assisted me in so many different ways along this incredible journey of learning.

First, I would like to express my special thanks to my supervisors emerita, Professor Irma Moilanen, M.D., Ph.D., Clinic of Child Psychiatry, University and University Hospital of Oulu; Professor Hannu Soini Ph.D., Department of Teacher Education at the University of Oulu; Professor Hanna Ebeling, M.D, Ph.D., Clinic of Child Psychiatry, University and University Hospital of Oulu and co-authors Professor, Alice S. Carter, Ph.D., Department of Psychology, University of Massachusetts Boston, USA and Päivi Pihlaja, Ph.D., Department of Special Education at the University of Turku. Without my supervisors and co-authors experience, knowledge and enthusiasm, I would not have found the courage to start this project, and so I warmly thank them for their motivation, support, encouragement and guidance.

A truly exceptional and matchless expression of gratitude is due to the children and families who participated in this study. Without the infants' joy and their parent's faithful commitment this study would not have been possible.

I would like to also offer my deepest gratitude to Clinical Psychologist and Instructor of Psychology, Rachel Pollock-Wurman, Ph.D., University of Massachusetts Boston, USA, and Clinical Psychologist Sanna Kuusikko-Gauffin, Ph.D., Clinic of Child Psychiatry, University and University Hospital of Oulu for their time, patience and enthusiastic and thoughtful assistance with the statistical analyses and language review for this thesis. With their help and support this dissertation was actualized. I owe my sincerest thanks to Docent, Tero Timonen, Ph.D., from the Department of Psychology, University of Joensuu, for his invaluable comments at the beginning of the study and to Sari Ahlqvist –Björkroth, Psc.M., Tanja Lipasti, Psc.M., and Speech and Language Therapist, Päivi Savela, M.A. for offering their time and wisdom for the completion of this study. I would like to acknowledge and offer many thanks to Leena Joskitt, M.A., Risto Bloigu, Ph.D. and Anne Karjalainen, for their support and encouragement throughout this process, along with their guidance with statistical analyses at the beginning of the study.

Next I would like to thank Marja-Leena Laakso, Ph.D. and Paula Lyytinen, Ph.D., from the Department of Psychology, University of Jyväskylä and Raija Vanhala, M.D., Ph.D., from the University and University Hospital of Helsinki for their assistance with the developmental questionnaires for this study. And, thank you, infinitely, to Katja Larinen M.Psych.Ed., who has assisted this research project throughout all of its phases. Thanks are also due to co-authors Varpu Penninkilampi-Kerola Ph.D. and Päivi Pihlaja Ph.D.

I owe my special thanks to The Graduate School of Circumpolar Wellbeing, Health and Adaptation for their financial and educational support. Thank you, especially to Project Secretary Hannele Savela, Ph.D. for her warm and positive guidance throughout our work together. To all of my graduate school doctoral-student colleagues, I am indebted to you for your wisdom throughout our time together.

I also want to express my warmest thanks to my friends Taija and Hannu, who have brought so many memorable and joyful moments to my life. To my friends Taru and Martti, the discussions we have shared together have been nothing short of inspirational, thank you.

Innumerable thanks to my family: my mother Kaarina, you raised me to be curious and enthusiastic, my sister Mervi, you have supported, guided and helped me in so many ways throughout my life, and my sister Kirsti, you have always supported me in my career, and you have offered so many valuable comment for this thesis; I thank you with all of my heart. Warmest thanks are also due to my brothers Jari and Jarmo, for their concrete help in practical matters.

Finally, I offer my deepest and unconditional love and gratitude to my husband Risto, who has loved, supported and helped me endlessly throughout all phases of this thesis and throughout all phases of our lives together. Without his immeasurable love and endless faith in me and my work, this study could not have been initiated or completed. My love and gratitude is also boundless for my remarkable children Veeti and Henrietta; I would have been lost many times without their existence; the contagious presence of their smiles and joy, and the warmth of their love was the perfect recipe for the necessary encouragement and help to persevere in so many ways in life and throughout the writing of this thesis. This study was financially supported by the Alma and K.A. Snellman Foundation, Oulu, Finland, the Emil Aaltonen Foundation, from grants to Docent Hanna Ebeling, the Juselius Foundation, from grants to Professor, Emeritus Irma Moilanen, the Child Psychiatric Research Foundation, Finland, the Graduate School of

Circumpolar Wellbeing, Health and Adaptation, and the Child Psychiatry Research Foundation, Oulu, Finland.

Oulu, Finland, September 2012

Helena Haapsamo



## Abbreviations

ANOVA	Analysis of Variance
ANCOVA	Analysis of Covariance
ASD	Autism Spectrum Disorder
ASEBA	Achenbach System of Empirically Based Assessment
BITSEA	Brief Infant Toddler Social-Emotional Assessment
CBCL	Child Behaviour Checklist
MCDI	MacArthur Communicative Developmental Inventories
PC-ERA	Parent-Child Early Relational Assessment
PDD	Pervasive Developmental Disorder
PSI	Parenting Stress Index
RMANOVA	Repeated Measures Multivariate Analysis of Variance





## List of original publications

This thesis is based on the following publications which are referred to in the text by their Roman numerals. In addition, this thesis includes some unpublished data.

- I Haapsamo H, Kuusikko-Gauffin S, Ebeling H, Larinen K, Penninkilampi-Kerola V, Soini H & Moilanen I (2012) Communication development and characteristics of influencing factors; A follow-up study from 8 to 36 months. *Early Child Dev Care*, iFirst Article 23.4.2012.<http://www.tandfonline.com/doi/abs/10.1080/03004430.2012.674523>
- II Haapsamo H, Ebeling H, Soini H, Joskitt L, Larinen K, Penninkilampi-Kerola V, Carter AS & Moilanen I (2009) Screening infants with social and emotional problems; A pilot study of the Brief Infant Toddler Social and Emotional Assessment (BITSEA) in the Northern Finland. *Int J of Circumpolar Health* 68(4): 386–393
- III Haapsamo H, Kuusikko-Gauffin S, Carter A, Pollock-Wurman R, Ebeling H, Joskitt L, Larinen K, Soini H, Pihlaja P & Moilanen I (2011). A Pilot longitudinal follow-up study of the Brief Infant Toddler Social Emotional Assessment (BITSEA): Examining toddler's social-emotional, behavioural and communicative development. *Early Child Dev Care* 182(11): 1487–1502. <http://www.tandfonline.com/doi/abs/10.1080/03004430.2011.622756>
- IV Haapsamo H, Pollock-Wurman, R, Kuusikko-Gauffin S, Ebeling H, Larinen K, Soini H & Moilanen I Maternal stress and young children's behavioural development; A prospective pilot study from 8 to 36 months in a Finnish sample. Manuscript.



# Content

<b>Abstract</b>	
<b>Tiivistelmä</b>	
<b>Acknowledgements</b>	<b>9</b>
<b>Abbreviations</b>	<b>13</b>
<b>List of original publications</b>	<b>15</b>
<b>Content</b>	<b>17</b>
<b>1 Introduction</b>	<b>19</b>
<b>2 Review of literature</b>	<b>21</b>
2.1 Early interaction and communicative development from 8 to 36 months.....	21
2.1.1 Evaluative assessments and interventions.....	23
2.1.2 Communicative development with or without interventions in follow-up studies.....	25
2.2 Social-emotional and behavioural problems and competences from 18 to 36 months.....	27
2.3 The mother-infant interaction and maternal mood.....	29
<b>3 Aims of the study</b>	<b>33</b>
<b>4 Subjects and methods</b>	<b>35</b>
4.1 Procedure, participants and study design.....	35
4.2 Measures and assessments.....	38
4.2.1 MacArthur Communicative Development Inventories (MCDI).....	38
4.2.2 Parent-Child Early-Relational Assessment (PC-ERA).....	40
4.2.3 Brief Infant/Toddler Social Emotional Assessment (BITSEA).....	40
4.2.4 Child Behaviour Checklist (CBCL).....	42
4.2.5 Parenting Stress Index Short-Form (PSI).....	42
4.2.6 Family based enrichment-intervention.....	43
4.2.7 Statistical methods.....	44
<b>5 Results</b>	<b>45</b>
5.1 Sample.....	45
5.2 Reliability of measures.....	45
5.3 Communicative development (MCDI questionnaire) in children aged 8 months to 36 months (Study I).....	47
5.3.1 Children's communicative development and PC-ERA.....	48

5.3.2	Communicative development and family-based intervention.....	49
5.3.3	Social-emotional skills (the BITSEA) and behavioural problems (the CBCL) by MCDI groups.....	51
5.4	Social-emotional, behavioural problems and competencies in children aged 18 and 36 months (Studies II and III).....	53
5.4.1	Associations between age groups (18 months and 36 months) and the BITSEA, CBCL and MCDI (studies II and III).....	54
5.5	Maternal stress at infant age of 8 months and children's behavioural problems at the ages of 18 months and 36 months (Study IV).....	58
5.5.1	Maternal stress level.....	59
5.5.2	Associations between the PSI and the CBCL.....	59
5.5.3	Longitudinal associations between PSI groups (High and Low) and CBCL Total problem scores, Internalizing and Externalizing subscales.....	61
5.5.4	The association of the PSI with the CBCL Total problem score over controlling for number of siblings.....	61
5.5.5	The association of the PSI with the CBCL Internalizing scores over time, controlling for number of siblings.....	63
5.5.6	Model 3: The Association of the PSI with the CBCL Externalizing score over time, controlling for number of siblings.....	64
<b>6</b>	<b>Discussion</b>	<b>67</b>
6.1	Main results.....	67
6.1.1	Communicative development: The MCDI at follow-up.....	67
6.1.2	Relationship between communicative development (MCDI) and social-emotional problems and competencies (BITSEA) and behavioural problems (CBCL).....	70
6.1.3	Associations between maternal stress and children's behaviour.....	73
6.1.4	Limitations.....	73
6.2	Conclusions: Implications and future research.....	75
	<b>References</b>	<b>77</b>
	<b>Original Publications</b>	<b>89</b>

# 1 Introduction

The unique aspects of individual development have been the subject of research fascination and inquiry for decades. In particular, infant development has been the focus of scientific interest for greater than fifty years, when Kanner published his first book concerning child psychiatry in 1935 (Kanner, 1935). Since this publication, a plethora of research has sought to address the important developmental phases throughout an individual's entire life, and relate these stages to early intervention and prevention of later problematic outcomes/behaviours.

Responsive/attentive and sensitive care, especially in the earliest months of an infant's life, is critical to child development (Olds, Sadler & Kitzman 2007). A parents' ability to attend to their infant's communicative signals and respond to them accurately, is associated with an infant's enhanced behavioural and emotional adjustment later in life (Mäntymaa *et al.* 2004) and parenting interventions are therefore essential at early stages (Ayoub, Vallotton & Mastergeorge 2011, Lowell *et al.* 2011).

Piaget & Inhelder (1977), proposed that during infant development, when an infant begins to explore the world by grabbing objects (i.e., the age of four to six months), represents a purposeful primary step to approach the environment and other persons. Understanding and becoming aware of communication and interaction is one basis to form relationships with others, and therefore the sense of a core self and core others later becomes integrated into the infant's world (Stern, 1985).

The first eighteen months, include, perhaps the most important phases of cognitive, intellectual and perceptual development (Piaget & Inhelder 1977). An individual's development involves a series of age-related stages (Elkind & Weiner 1978) whereby certain abilities/competencies (e.g., intentional gestures predicting early communicative acts) evolve during certain phases.

Response to joint attention between child-caregiver and infant has been related to language development at 18-months (Markus *et al.* 2000). Stern (1985), has proposed that an infants' subjective experience of self and others forms at approximately 15–18 months, concurrently with his/her ability to point with the index finger, an important developmental achievement. In order to enhance communicative skills, infants must have developed an interactive and close relationship to their caregiver; otherwise it becomes extremely difficult to develop meaningful social-emotional communication and language (Stern 1985).

Language development and social interaction are related to emotional expressions. Research suggests that children, who start to speak earlier, also have more discernible emotional expressions before their first clear words (Bloom & Tinker 2001). Moreover, parents who rate their infants as having low communication skills, as measured by the MacArthur Communicative Development Inventories, MCDI (Fenson *et al.* 1994) and low social-emotional competence, as measured by the Brief Infant Toddler Social-Emotional Assessment, BITSEA (Briggs-Gowan & Carter, 2006), are more likely to report worry about their children's development (Briggs-Gowan *et al.* 2004).

Children with delayed verbal comprehension and general language delays may manifest more behavioural problems than do children with communication development within the "normative" range (Silva, Williams & McGee 1987, Rescorla, Ross & McClure 2007a). Irwin, Carter & Briggs-Gowan (2002), reported that "late-talkers" were observed to be more withdrawn and/or depressed and moreover were less interested in playing with same aged peers, than were controls with normative language development. The parent-child interaction was also affected; specifically mothers of "late-talkers" reported higher parenting stress on the Parent-Child Dysfunction scale of the Parenting Stress Index (PSI, Abidin 1999). Based on the abovementioned research findings, parent-child dyads seem to be highly interactive and therefore it would be appropriate for early interventions to target both children and parents.

## 2 Review of literature

### 2.1 Early interaction and communicative development from 8 to 36 months

Language development and social competence are acquired within parent-child interactions with language acquisition beginning very early in the first months of a child's life. More important than vocabulary or grammar is the communicative functionality of affective expressions in early care giving situations (Överlund, 1996). Early symbolic actions and communicative gestures have relatively strong positive relationship with later language comprehension (Laakso *et al.* 1999, Paavola 2006a, Reilly *et al.* 2006).

In children's communicative development, there is a dramatic change between 8 and 12 months, when in particular gestural skills are developing and improving. Preverbal communication, such as gestures and positions (e.g., pointing, imitating the actions of others) manifest early in an infant's life, typically by the age of nine months (Wetherby & Pritzand 1999). Moreover, research suggests, that joint attention skills, when a child starts to confirm pointing with an alternating gaze, are positively associated with later language development (Carpenter, Nagell & Tomasello 1998, Morales *et al.* 2000).

Pointing is one of the notable joint attention skills emerging at this phase of a child's communication development. This phase (from eight to twelve months of age) also includes large changes in prelinguistic (nonverbal communicative actions such as holding hands up as a request to be held in someone's arms) and linguistic (first words) communication skills (Reilly *et al.* 2006). Further, gestures such as reaching for something and giving/handing objects to others can be demonstrative of interactive meaning even before a child's speech production, which typically develops at the age of nine months (Jakkula 2002). Other nonverbal communicative gestures and appropriate actions with objects in social situations at the age of 14 months have been shown to predict early vocabulary comprehension at age of 18 months (Laakso *et al.* 1999).

Early language acquisition is an integral part of a child's cognitive and social-communicative development (Tomasello 2000). Children's receptive development of language (comprehension) begins much earlier than productive language development (verbal expressions and vocabulary) and infants younger than one year of age understand a great number of words and communicative expressions (Lyyt-

inen, 1999a). Initially, young children learn to differentiate the phonemes of their native language, and become aware of the formula of the words (Laalo 1998).

Maternal sensitivity (appropriate reactions and actions to infant's needs) has also been associated with early intentional communication (Carpenter, Nagell & Tomasello, 1998, Paavola *et al.* 2006b) and infant behaviour (Kivijärvi *et al.* 2001, Dollberg, Feldman & Keren, 2010). In addition, mothers' postpartum stress has been associated with general cognitive and language development in young children (Ayoub, Vallotton & Mastergeorge 2011, Keirn *et al.* 2011).

In Finland, children are typically producing their first words at the age of nine to 13 months (Kunnari, 1997), and it is very typical that the first words are nominal (Laalo 1998). At the age of two years, children's language development starts to have more complex meanings; for example the word "cup" means the exact word, noun and quality, and the situation is very strongly related to the interpretation of the word (Parre 1994). In the study of Reilly *et al.* (2008), two-year-old-children in the U. S. scored an average of 260 points in the *MacArthur Communicative Development Inventories* (MCDI, Fenson *et al.* 1994, Finnish version by Lyytinen 1999a), likewise in the study by Laakso and colleagues (1999), Finnish children, on average, scored  $M = 265.1$  points on the MCDI *Total* scores. The morphological communicative development appears to peak at the age of two and a half years in Finnish children (Toivainen 1994b).

Early linguistic communication skills may be related to gender. According to a study by Paavola and colleagues (2006a), girls are more likely to produce comprehensible words than are boys by the age of 12 months. Reilly *et al.* (2007), found that male gender and problematic family history were strongly associated with poorer outcomes in communication skills. Similarly, Mauridsen & Hauschild (2010), found that boys may be more vulnerable than girls to different environmental stressors (e.g., low socioeconomic status), and that male sex hormones may be implicated in the aetiology of developmental language disorder in clinically assessed children from cohort sample (Mauridsen & Hauschild 2010).

Because language comprehension and the maturity of verbal communication are developing in cooperative situations, play is one important part of this emerging relationship. Naming toys and playing with those specific items are central aspects of verbalizing these actions (Tolonen, 1996). Children's symbolic play at the age of 14 months has longitudinally predicted vocabulary production at the age of 24 months; however, studies suggest that after 24 months, the contribution of symbolic play is no longer statistically predictive (Lyytinen *et al.* 1999b).



In general, children with slow motor development or familial risk for dyslexia have smaller vocabularies and produce shorter sentences than other children without similar risk factors (Viholainen *et al.* 2002). By the age of three, children are extending their vocabulary by naming objects. Children who are trained to name objects and to categorize them, have shown dramatic changes in novel language acquisition, a skill that may transfer to naming new objects outside of those learned during the training sessions (Smith *et al.* 2002).

Although genetic factors play a significant role in early language development (Plomin & Dale, 2000, Mauridsen & Hauschild, 2010), there are many other factors influencing the development of communication. For example, children's language and communication development is related to differentiation of emotional expressions (Bloom & Tinker, 2001) and thus more differentiated emotional expressions may be related to greater communicative action.

Language development also may influence building peer-relationships; longitudinal analysis of early language difficulties has predicted decreased friendship quality later in adolescence (Durkin & Conti-Ramsden 2007). Of interest, Kokkonen and colleagues (2003) found, that the ability to speak (at the age of 1 year) is negatively associated with symptoms of alexithymia 30 years later (cohort-study in Northern-Finland).

In addition, family background may effect to children's communicative development (Stein *et al.* 2008); maternal education is significantly associated with two-year-old children's maximum sentence length as well as cognitive development (Lyytinen *et al.* 1999b). Moreover, research suggests the influential role of socio-economic status (SES) on infant language growth (Hutterlocher *et al.* 2010, Reilly *et al.* 2007, 2010, Lung *et al.* 2010).

### **2.1.1 Evaluative assessments and interventions**

Interactional and communication skills are most observable in the home environment (Lyytinen, 1999a). Most of children's developmental questionnaires assess interaction, language and behaviour (for example the Child Behaviour Checklist, CBCL, Achenbach & Rescorla 2001, Finnish version by Almqvist 2001), and therefore comprehensive and productive language skills are observable in a variety of situations with different methods of measurement (e.g., via the MCDI). Parents' knowledge of their children's skills is essential and is many times adequate, as parents are able to monitor their child in everyday situations in life (Bricker *et al.* 1988, Lyytinen *et al.* 1999a, Charman *et al.* 2003).

When examining developmental or language disorders, it is important to employ methodologies and questionnaires that reliably assess areas of targeted need. For example, sensorimotor development (Bernabei *et al.* 2003) or semi-structured play-interaction (Trillingsgaard *et al.* 2005) can be identifiable areas to differentiate autism from other developmental disorders. Also both communicative acts and joint attention can be important indicators when screening children with possible Autism Spectrum Disorders (ASDs), since delayed use of gestures and interactive play are commonly identified by the age of 12 months in children later diagnosed as ASD (Mitchell *et al.* 2006).

In both, research and clinical settings, it is important to utilize empirically valid evaluative tools to identify deviations in language development early enough, so that intervention and rehabilitation (e.g., speech therapy) can be initiated in the early stages. Buschmann and colleagues (2008), found that if a child's reported vocabulary is lower than 50 words at the age of two years, further investigation and possible early intervention is suggested. In Finland, the punctual healthcare system observes child language development very carefully, and if a child demonstrates less than average communicative skills in productive or comprehensive language, further clinical assessment is recommended (Kiehellinen erityisvaikeus, Käypä Hoito -suositus 2010).

Although mothers' may rate their infants as more competent, regardless of their child's actual abilities to communicate (Delgado & Delgado 2002) (particularly during clinical evaluation), parent and professional agreement on developmental questionnaires are often reliable and valid (Squires, Bricker & Potter 1997, Lyytinen, 1999a, Charman *et al.* 2003).

In order to identify children at moderate-risk for interventional services, parent-report assessments are a particularly useful tool in a true multi-trait, multi-method assessment (Hughes *et al.* 1998). For example, 23–47.7% of paediatricians utilize parent-report measures in U. S. (Radecki *et al.* 2011).

Many times, parental worry is the first predictor of children's problematic development (Ellingson *et al.* 2004); however, parents of young children are often unaware that the behaviours they report are associated with concerning emotional or behavioural problems (Briggs-Gowan & Carter 2008). Thus, it is valid to suggest early screening of children to help identify young individuals with delayed development and/or social-emotional and behavioural problems, which may ultimately facilitate referral to early intervention. Based on psychometrically sound assessments of competencies (e.g., the BITSEA *Competence* subscale), interven-

tion can be targeted to certain areas of interaction development and areas of need, such as the use of gestures, play and communication skills.

Parent-directed early interventions have gained research attention and have demonstrated empirical validity for children with behavioural or developmental problems (Tonge *et al.* 2006); brief parenting programmes alone may be insufficient to prevent the development of children's externalizing problems (Bayer *et al.* 2010). However, some parent education and behaviour management interventions have shown efficacy for reducing anxiety, insomnia and somatic symptoms in the parents of autistic children (Tonge *et al.* 2006).

To date, family-based intervention studies (i.e., frequency = once a month), have failed to demonstrate efficacy in reducing children's behavioural problems; however there is evidence that enhancing the quality of the mother-infant relationship may result in a significant positive relationship (Cheng *et al.* 2007). Therefore, a carefully planned early intervention focusing on improving the quality of the mother-infant relationship may prove to be the direction of early intervention in children with communicative language deficiencies (Cheng *et al.* 2007). For example, in a study by Smith, Groen & Wynn (2000), children with Pervasive Developmental Disorder, not otherwise specified (PDD-NOS), showed a statistically significant improvement in measures of intelligence, visual-spatial skills, language and academics following intensive treatment or parent training, addressing the importance of intervention. Unfortunately, there is a paucity of empirically validated studies employing family-based early intervention for parent-child-interaction and child behavioural development, especially in Northern-Finland.

### **2.1.2 Communicative development with or without interventions in follow-up studies**

Longitudinal follow-up studies suggest that delay in language abilities, such as expressive and receptive language, are strongly associated with later emergent/diagnosed psychiatric problems (Charman *et al.* 2003, Toth *et al.* 2006). Early intervention for developing communication skills is thus essential, because of the identifiable milestones of children's developmental stages (e.g., Piaget & Inhelder, 1977). Research of early phonological and language skills also supports a strong genetic component as a statistically significant predictor of language and communicative development (e.g., reading difficulties; Puolakanaho *et al.* 2007). Follow-up studies of communication development and/or intervention are presented in table 1.

**Table 1. Communicative development with or without intervention in follow-up.**

Author(s), year of publication	Title of research article	Participants	Methods	Main results after intervention or follow-up
Best, Melvin & Williams, 1993	The effectiveness of communication groups in day nurseries	36 children, mean age 33.9 months	Free play, attention, listening, language games and action songs	Significantly more improvement of children's interactions
Carpenter, Nagell & Tomasello, 1998	Social cognition, joint attention and communicative competence from 9 to 15 months of age	24 mother-infant pairs, 9-15 month old children	Free-play video analyses and interview, MCDI	Social-cognitive skills are associated with better language skills
Smith, Groen & Wynn, 2000	Randomized trial of intensive early intervention for children with PDD	28 children (14 with autism, 14 with PDD-NOS), mean age 36 months	Parent training or intensive treatment, IQ, RDLS, CBCL, Wechsler Individualized Achievement test, Early Learning Measure	Intensive treatment group had a statistically significant benefit compared to parent training group. PDD-group scored higher than autism group
Jakkula, 2002	Giving objects; A mirror of development and a social sign anticipating language acquisition of 9–34 month old children	6 mother-child-pairs	Reynell Verbal comprehension and Verbal production, parent interview and videoanalysis	Object giving mirrors social skills and language development
Mendelsohn <i>et al.</i> 2007	Use of videotaped interactions during paediatric well-child care: impact at 33 months on parenting and on child development	99 children with developmental delay risk and with low maternal education aged 33 months	Video Interaction Project with paediatric primary care consultations	Intervention group showed lower levels of parenting stress and more likely typical cognitive development and fewer developmental delays
Puolakanaho <i>et al.</i> 2007	Very early phonological and language skills: estimating individual risk of reading disability	192 children with or without familial risk of dyslexia, mean ages 3,5 and 4,5 and 5,5	Phonological Awareness, rapid naming, short-term memory, expressive vocabulary, pseudoword repetition, letter naming and performance IQ	Familial risk status is a statistically significant predictor of reading disability

Author(s), year of publication	Title of research article	Participants	Methods	Main results after intervention or follow-up
Landry <i>et al.</i> 2008	A responsive parenting intervention: the optimal timing across early childhood for impacting maternal behaviours and child outcomes	80 babies born full-term and 86 pre-term babies and their mothers, follow-up from 30,2 months to 38,2 months of age	Peabody Picture Vocabulary Test III, Preschool Language Scales 3rd Ed., Mother-Child interaction videotape, 2 different interventions in infants and/or toddlers	Mothers' responsive behaviour increased child's skills; most improvement gained after 2 periods of intervention
Buschmann <i>et al.</i> 2009	Parent based language intervention for 2-year-old children with specific expressive language delay: a randomized controlled trial	61 children with specific expressive language delay aged 24.7 months	MCDI, Heidelberg Parent-based Language Intervention	Highly structured intervention reduced language impairment
Wake <i>et al.</i> 2011	Outcomes of population based language promotion for slow to talk toddlers at ages 2 and 3 years: let's learn language cluster randomised controlled trial	1217 children, initially assessed at 12 months and screened at follow-up at the age of 2 and 3 years. Six week intervention initiated at the age of 18 months	Child Behaviour Checklist, Preschool Language Scale, parental advice and support strategies for adaptive interaction and language enhancement tools to be used with the child	Intervention showed little evidence of improved language skills or behaviour

## 2.2 Social-emotional and behavioural problems and competences from 18 to 36 months

It is estimated, that between ten and thirty percent of children are in need of special support or rehabilitation in Southern-Finland (Hakulinen-Viitanen, Pelkonen & Haapakorva 2005). In the United States (U.S.) the prevalence estimate of one-to two-year-old children's social-emotional and behavioural problems is 10–15% (Briggs-Gowan *et al.* 2001). Gross cultural differences and different evaluation methods may explain a majority of the variability in social-emotional and behavioural problems identified in different countries; however it is critical to note that the ultimate goal of early recognition and intervention is a common goal (cf., Ellingson *et al.* 2004 and Pihlakoski *et al.* 2006, Bayer *et al.* 2010).

The most often occurring problems with small children's social-emotional and behavioural development include conflict-aggressive, self-destructive behaviours or similar features (e.g., Rubin *et al.* 2003). Many times, these symptoms as

well as behavioural withdrawal are stable over time, and predict long-term problematic behaviours (Pihlakoski *et al.* 2006). Longitudinal follow-up studies have demonstrated the stability of somatic complaints from early childhood to preadolescence as well (Pihlakoski *et al.* 2006).

Social-emotional competencies (e.g., attention skills, mastery motivation, imitation and play behaviour, emerging empathy and prosocial interactions with peers) are predictors of young children's behaviour (Ellingson *et al.* 2004). In a study by Ellingson and colleagues (2004), parents who reported low social-emotional competence of their child at the age of 11 to 39 months, also reported more behavioural problems in their child as measured by the BITSEA (Ellingson *et al.* 2004). It is important to take into account temperamental differences in young children, particularly parent-report of such differences. For example, a difficult child temperament (e.g., emotion and behavioural dysregulation and conflictual-aggressive parent-child interactions) may interact with maternal stress and expressed negativity toward a child (Rubin *et al.* 2003).

Briggs-Gowan & Carter (2008), reported that the majority of children, who exhibit significant subclinical/clinical emotional or behavioural symptoms by 12 to 36 months of age, based on parental report, are rated by teachers as exhibiting emotional/behavioural problems in early elementary school. In a study by Rubin *et al.* (2003), results indicated that especially boys' externalizing difficulties at the age of two years predicted later conflict-aggression behaviour at the age of four years.

Behavioural problems may also be associated with delayed verbal comprehension and general language delay (Silva, Williams & McGee, 1987). Irwin, Carter and Briggs-Gowan (2002), reported that withdrawn and/or depressed children are less interested in playing than are children, who are better able to express themselves verbally and have normative language development.

Early recognition of behavioural problems in young children and infants is of great research and clinical importance. Such areas of need (e.g. internalizing and externalizing behaviour, aggressive behaviour and somatic complaints) identified early in childhood have been shown to predict later long-term risk for psychopathology (e.g., anxiety disorders, depression or ADHD) in preadolescence (Pihlakoski *et al.* 2006). Externalizing behaviours, in particular, may negatively affect later relationships with caregivers (e.g., low warmth and affective enjoyment within the parent-child dyad) (Olson *et al.* 2000).

### **2.3 The mother-infant interaction and maternal mood**

W. Ernest Freud (1989, p. 497) said: "A mother and her baby are intended to fulfil each other," and awareness of this unique relationship involves very sensitive, influential, and reciprocal factors. Maternal expectations during pregnancy (e.g., negative emotions toward pregnancy) may affect the parent-infant relationship (Fowles & Walker, 2009) and pregnancy-related anxiety may be one variable related to postnatal parental stress (Saisto *et al.* 2008).

The parent-infant relationship, which develops from the very beginning of an infant's life, may be vulnerable to developmental distress or instability; mothers that are experiencing clinical levels of depression and life difficulties/psychosocial stressors, report more behavioural problems with their children compared to mothers without such difficulties (Prior *et al.* 2008, Mothander & Moe, 2010). Research also has demonstrated that low social engagement, less mature regulatory behaviours, and more negative emotionality are common features in one year old infants of depressed mothers (Feldman *et al.* 2009, Keirn *et al.* 2011). In the recent study by Dollberg and colleagues (2010), results suggest a link between maternal representations and infant psychiatric problems or mastery motivation (Sparks *et al.* 2011), and underline the mother-child relationship as influenced by pervasive, negative emotional tone in mother-infant interactions.

A mothers' ability to attend to infant communication skills is especially important when interacting with a child with developmental delays; sensitive mothers can adjust their own communication so that it is appropriate and consistent with their child's communication and reflective of particular task demands (Guralnick *et al.* 2008). Additionally, maternal sensitivity has been associated with early intentional communication (Paavola *et al.* 2006b) and infant behaviour (Kivijärvi *et al.* 2001). The mother-infant relationship has its own unique features and there is a paucity of longitudinal research focused on the mother-infant relationship in Finnish culture. Some of Finnish mother-infant studies are presented in table 2.

**Table 2. Finnish studies of mother-infant relationships and infant behaviour and communication.**

Author(s), year of publication	Title of research article	Participants	Methods	Results
Tamminen, 1990	Postnatal depression, breastfeeding and mother- infant interaction	279 first-time mothers	NPI,PSE,EPDS	Depression negatively effects breastfeeding and interaction with the baby
Pajulo <i>et al.</i> 2001	An early report on the mother-baby interactive capacity of substance- abusing mothers	12 mothers with alcohol or drug abuse and 12 mothers with no alcohol or drug abuse and their babies	PC-ERA, EPDS	Mothers with substance- abuse have more problems in their interactive behaviours with their children
Kivijärvi <i>et al.</i> 2005	A contented baby has a sensitive mother	57 non-risk mothers and low-risk mothers	PC-ERA, R-ITQ	A positive association emerged between more sensitive mothers and more contented babies
Savonlahti <i>et al.</i> 2005	Interactive skills of infants with their high-risk mothers	14 high-risk mothers and 12 low-risk mothers	PC-ERA	High-risk mothers demonstrated more interactive deficiencies than low risk mothers
Mäntymaa <i>et al.</i> 2004	Early Mother-Infant Interaction	131 mothers and infants	PSI, ICQ, CBCL	Mother-infant relationship is associated with infants' wellbeing
Paavola, 2006a	Maternal responsiveness, characteristics and relations to child early communicative and linguistic development	27 mothers and their children at the age of 10, 12 and 30 months	PC-ERA, MCDI, Care-Index	Sensitive responsiveness to interaction predicts communicative skills at infants age of 12 months
Kemppinen <i>et al.</i> 2007	Early maternal sensitivity: Continuity and related risk factors	5 infant-mother dyads, 78- 74 well-baby clinic sample and 27 linguistic sample	PC-ERA, Care- Index, MCDI, EPDS, GHQ	Continuity of problems in early interaction and in mother-infant dyads warrants early identification and intervention

NPI = Neonatal Perception Inventory, Broussard & Hartner, 1971, PSE= Present State Examination, Wing *et al.* 1974, EPDS = Edinburgh Postnatal Depression Scale, Cox, Holden & Sagovsky, 1987, PC-ERA = Parent-Child Early Relational Assessment; Clark, 1985, GHQ = General Health Questionnaire, Goldberg *et al.* 1997, R-ITQ = Revised-Infant Temperament Questionnaire, Carey & McDevitt, 1978, ICQ=Infant Characteristics Questionnaire, Bates *et al.* 1979, CARE-Index = assessment method for adult-infant interaction, Crittenden, 1998.



Mother-infant relationships may be reciprocal in nature; specifically, it may be more challenging for a mother to respond sensitively to a high reactive infant, and in turn, diminished maternal sensitivity may increase crying and high reactivity and decrease emotion regulation in the infant (Kagan, 1997, Mäntymaa *et al.* 2004). Mothers of infants who engage in more crying and fussing and less contented behaviour appear less sensitive in and less attentive to their behavioural interactions with their infants (Kivijärvi *et al.* 2004). Moreover, depressed mothers may misinterpret or may negatively bias their child's messages and needs (Kemppinen *et al.* 2007). These effects may be particularly salient in early breastfeeding interactions, including preterm infants (Tamminen, 1990, Meyer *et al.* 1994).

Research suggests, that maternal anxiety is associated with exaggerated behaviour (e.g., high reactivity and emotional expressions) toward infant in free play and teaching situations (Kaitz *et al.* 2010) This trend appears consistent when investigating the association of mothers' postpartum stress and general cognitive and language development in infants (Keirn *et al.* 2011). In addition, toddler temperament has predicted both parental stress (Hirschfeld *et al.* 2004, Kagan, 1997) and problematic routine care-taking (Saisto *et al.* 2008).

Yamada *et al.* (2007), and Schieve *et al.* (2008), report that emotional stress levels are higher among mothers who are caring for children with various developmental problems (e.g., PDD, *Pervasive Developmental Disorder*). There may also be an association between mothers' psychological distress and infant sleep; Baird and colleagues (2009), suggest that women experiencing psychological distress prior to conceiving a child, are more likely to have babies with sleep disturbances. Furthermore, familial variables and chronic psychosocial stressors (e.g., low parental education, socioeconomic disadvantage, non-intact home environments, and parental mental health difficulties), may negatively impact children's development, and be associated with increased behavioural problems in infancy and throughout childhood (Lung *et al.* 2010, Sperlich *et al.* 2011). Specifically, Yaman and colleagues (2010) noted that an increase of family stress is related to higher rates of externalizing behaviours in toddlers.



### **3 Aims of the study**

This longitudinal follow-up study focuses on communicative development and its associations to social-emotional and behavioural problems and competences of typically developed children from eight to 36 months. With this study, we also wanted to examine the properties of the BITSEA in a pilot sample in Northern-Finland. Additional goals were to better understand and characterize possible associations between maternal stress and children's behaviour, due to a hypothesized reciprocal nature of mother-infant dyad.

The specific aims were as follows:

1. To follow-up children's communicative development (the MCDI) from 8 to 36 months, and to study the efficacy of family-based enrichment-intervention to children's communicative development (Study I).
2. To follow-up children's development via parent-report questionnaires (the BITSEA and the CBCL) and examine their associations (Studies II and III).
3. To characterize the relationship between maternal stress (PSI) and children's behavioural development (CBCL) from 8 to 36 months (Study IV).



## 4 Subjects and methods

### 4.1 Procedure, participants and study design

This follow-up study was approved in 2006 by the Ethical Committee of the Northern Ostrobothnia Hospital District, the University Hospital of Oulu and the Municipal Board of the Social and Health Care Unit of the City of Oulu. The author of this thesis and principal researcher Helena Haapsamo (H.H.), designed this follow-up study with a research-team and each five-month enrichment-intervention was completed by H.H. She also videotaped families for the Parent-Child Early Relational Assessment-analysis and delivered the questionnaires to Health Care Centres. All original articles (I, II, III and IV), presentations, reports and methods have been prepared by the author in collaboration with the study-group.

#### *Sample collection*

At the beginning of the study (year 2006), all children's health care centres in the city of Oulu were informed about the research, and children's health care nurses were provided with details about the project design and -protocol. Five Children's Health Care Centres for this study were chosen, to best represent the distribution of social classes in Oulu.

Initial questionnaires in the beginning of the study were: the MCDI, a demographic information form and the PSI-short form. Families in 2006 ( $n = 31$ ) were randomly chosen to family-based intervention ( $n = 14$ ), which is detailed below in the methods-section. The demographic information form, MCDI and PSI - questionnaires were given to all families, whose typically developing infant had his/her six month well-baby check up by a Children's Health Care nurse. All children of participating parents were born at full gestational term and without any previously diagnosed abnormalities (e.g., hearing loss or Down syndrome), which could impact normal child development.

Parents were asked to complete the packet of questionnaires when their infant was eight months old (at the age of 8 months there is no scheduled well-baby check-up). Subsequently, a study researcher (H.H.) visited participants' homes in order to conduct and video-tape the Parent-Child Early Relational Assessment (PC-ERA; Clark, 1981, 1985, Finnish version Ahlqvist, 2003). For this study, the


mother-child-dyads were first assessed with the PC-ERA when the infant was between 9 and 12 months old, and dyads were examined with the PC-ERA for a second time when the children were between the ages of 15 to 18 months. At both assessment points, mothers were informed to feed and interact with their child consistent with their typical daily routine.

The overall sample size increased (and was collected) during the Fall of 2007 (n = 19); however, these participants did not have PC-ERA -video sessions or family-based enrichment intervention, in order to obtain comparative "no intervention" data. Follow-up questionnaires were delivered altogether 103 families (at year 2006 and 2007) and 50 families returned the evaluative questionnaires.

### *The follow-up*

Children's development in our sample was evaluated in four different chronological phases: Eight, 18, 24 and 36 months of age. Children's social-emotional and behavioural development was evaluated via parent report at the toddlers' age of 18 months and 36 months with the Brief Infant Toddler Social Emotional Assessment (BITSEA, Carter *et al.* 2001) and the Child Behaviour Checklist (Achenbach, 2001, Finnish version Almqvist, 2001). In this study, our goal was to examine children's communicative development and its associations to social-emotional and behavioural development (both typical and problematic), over time via the MCDI, BITSEA, and CBCL questionnaires. The questionnaires (the BITSEA and CBCL) were mailed to all participants from the original samples (year 2006 and 2007, N = 50) when children were ages 18 and 36 months, and the communicative development was evaluated by parents when children were eight months and then followed up at the ages of 18, 24 and 36 months. The design of this longitudinal follow-up study is illustrated in Figure 1.

**A Follow-up study of children's communicative development and its associations to social-emotional and behavioural development from 8 to 36 months**



**Fig. 1. Study Design Flow Chart.**

## 4.2 Measures and assessments

### 4.2.1 MacArthur Communicative Development Inventories (MCDI)

The MCDI for 8 to 16 months is a 468-item, continuous, positive scale, which assesses communication development. The MCDI can be divided into 2 subscales: 1) *First words* (412 items) and 2) *Communicative actions and gestures* (56 items). The *First Words* subscale can be further divided into 4 subscales: *First signs of comprehension*, *Comprehension of questions and instructions*, *First words*, and *Vocabulary*. The *Communicative actions and gestures* subscale can be further divided into 6 subscales: *Communicative gestures and actions*, *Play*, *Communicative action with objects*, *Imitating play of parenthood*, *Imitating actions*, and *Object replacement in play*.

The MCDI scale for 16 to 30 month old children includes 619 items and can be divided into 2 subscales: 1) *Words* (600 items) consisting subscales of *Vocabulary* (20 categories, 595 words) and *Pragmatics* (5 items), and 2) *Grammar and sentences* (16 items) consisting of subscales including *Plural and suffixes* (9 items), *Verbs* (7 items) and *Combination of words* (3 items, which are counted from the child's longest utterance's morphemes and then divided by three). The Finnish version of the MCDI has been translated and examined by Lyytinen (1999a), and it has demonstrated good psychometric properties. Evidence supports the use of the MCDI as an easily administered, reliable and valid method for screening infant and toddler communication development.

The Finnish versions of the MCDI have been used for over two decades (Lyytinen 1999a) for clinical purposes by speech and language therapists. According to Finnish studies (e.g., Lyytinen, 1999a, 1999b, Paavola *et al.* 2006a, 2006b and Stolt *et al.* 2007), the MCDIs are comparable to other linguistic measures, such as the Reynell Developmental Language Scales (RDLS III, Edwards *et al.* 1997, Finnish version 2001) in evaluating communication and language development in young children.

The MCDI uses a method where parents observe their child's communication and language development. Items are rated in both (for younger and older children) versions:

0 = no understanding or vocabulary production or communicative actions,

1 = understands and produces the vocabulary or communicative actions.

Thus, higher scores indicate better communication skills.



For this study, we selected 17 MCDI questions as a revised and short version to study children's communication development (vocabulary and comprehension) at the ages of eight and 18 months. This brief version was determined via clinicians' consensus from early vocabulary and understanding. Because the range of the results (vocabulary and comprehension) at the age of eight month varied so greatly ( $M = 6.4$   $SD = 2.83$ , range = 1–12), we employed the same 17 questions to examine change over time at the 18-month-old follow-up assessment. Also, our family-based intervention included supportive aspects to the MCDI's *First communicative actions and gestures* and *Communicative actions with objects* subscales, as we sought to compare outcomes on the MCDI pre- and post-intervention. A brief version (99 items) was also used by Guiberson, Rodriguez & Dale (2011); the authors found that the shorter version yielded highly satisfactory classification accuracy on expressive language delays. In clinical situations, it might be more appropriate to use the original version of the MCDI, but due to time constraints and in order to reduce participant fatigue, we decided to utilize the shortened version of the MCDI for this study.

At the age of eight and 18 months, we selected items from the *First Words* subscale: 1) reacts, when calling the child's name, by turning to or looking for the caller, 2) reacts, when saying "No" by stopping the action for at least a short period of time, and 3) reacts to a familiar sentence, such as "Where is the mom/dad?" by looking for that person. The selected items from the *Communicative actions and gestures* subscale were: 1) reaches and shows an object, that he/she has in his/her hand, 2) gives a parent a toy or an object, after parent request 3) points (by finger or hand) to an interesting object or action, 4) waves hand, when leaving ("bye bye"), 5) reaches hands up to communicate a desire to be picked up or held in a caregiver's arms, 6) shakes head for "no", 7) nods head for "yes", 8) puts finger to lips for "shh" (quiet), 9) blows a kiss, 10) smacks lips, when something tastes good. The selected 10 questions from the *Communicative actions with objects* subscale were: 1) tries to eat with spoon or fork, 2) drinks from a cup independently, 3) places a hat on his/her head, 4) throws a ball. We selected questions that exemplified the most common vocabulary expressions and signs of comprehension in everyday situations.

At the age of 24 months we used the original Finnish version (including all 619 items) of the MCDI for 16- to 30- month old children (Lyytinen, 1999a). Because to date, there are no Finnish parent-report questionnaires of communicative development for children aged 36 months, and the variability of our results at earlier stages was quite apparent, we chose to use the MCDI for 16- to 30- month

old children (Lyytinen, 1999a) for the 36-month olds in our sample. The typical variation of communicative skills of the children's age group (at all ages) falls between 25 to 75% of the whole sample (Lyytinen, 1999a). The mean of the typical variation is used as a comparison value for individuals' communicative development.

#### **4.2.2 Parent-Child Early-Relational Assessment (PC-ERA)**

The PC-ERA is a video-taped, structured (researcher-observed) method to evaluate parent-infant relationship. The PC-ERA includes 64 items, which are scored on a five-point Likert-type scale (scores 1–2 = area of concern, 3 = some concern, 4–5 = area of strength). Thus, lower PC-ERA scores are indicative of more problematic early parent-infant interactions. The PC-ERA includes three variables: 1) *Parental variables* (29 items), 2) *Infant variables* (27 items) and 3) *Dyadic variables* (8 items).

The video sessions can be recorded at home in feeding situations (five minutes) according to the manual of the PC-ERA. The PC-ERA has been used in several international studies (Harel *et al.* 2002, Kempainen *et al.* 2005, Kivijärvi *et al.* 2005, Korja *et al.* 2007).

#### **4.2.3 Brief Infant/Toddler Social Emotional Assessment (BITSEA)**

The BITSEA is a screening tool for identifying social-emotional and behavioural problems and/or delays or deficits as well as social-emotional competence in children ages from 12 months to 35 months and 30 days (Briggs-Gowan *et al.* 2001). The BITSEA includes a total of 42 items that address 1) externalizing problems (six questions, addressing difficulties with activity/impulsivity, aggression/defiance and peer aggression), 2) internalizing problems (eight questions addressing fearfulness, worry, nervousness and distress upon separation, anxiety and social withdrawal), 3) dysregulation problems (eight questions addressing negative emotionality, sleep, eating, and sensory sensitivities problems), 4) competence items (11 questions addressing positive features of behaviour including attention skills, mastery motivation, imitation/play behaviour, prosocial interaction with peers and emerging empathy), 5) and autism spectrum disorder (ASD) items (17 questions addressing repetitive behaviours, social competence and joint attention). In addition, there are red flag concerns (14 questions addressing clinically significant problems, such as *does not react when hurt, hurts self on purpose*

and *gags or chokes on food*), which are important to healthcare professionals as an indicator to follow the infant's development more closely. Red flag and ASD items partially overlap with the other subscales (Briggs-Gowan & Carter, 2006).

The BITSEA response format is: 0 = Not true/rarely, 1 = Somewhat true/Sometimes, and 2 = Very True/Often, thus higher scores reflect a greater number of identified problems. The BITSEA can be used as a parent-administered questionnaire or as a structured interview (Briggs-Gowan & Carter, 2006). In this study the BITSEA was used solely as a parent-administered questionnaire.

In the United States, the cut-off scores for the BITSEA *Problem Total score* scale (i.e, the sum of the *Externalizing*, *Internalizing* and *Dysregulation* scales) correspond approximately to the 25th  $\leq$  percentile at the ages of 12 months and 35 months and 30 days. The cut-off scores vary between genders; the cut score values of the BITSEA *Problem Total score* for girls are 12 at the age of 18 months and 13 at the age of 35 months, 30 days, and for boys 14 at the age of 18 months and 15 at the age of 35 months and 30 days. The BITSEA *Competence* scores (15th percentile ranking or less) also vary between genders; the cut-off is 13 points for girls and 15 points for boys at the age of 18 months, and 12 points for girls and 14 for boys at the age of 35 months and 30 days. There are no published cut-off scores for the specific subscales of the BITSEA (e.g., *ASD* or *Red flag* items), therefore results for these scores are interpreted from the *Problem total* scores and *Competence* scores. Because the BITSEA has not yet been examined or validated in a Finnish population, we utilized the mean cut-off scores in our sample, following the instructions of the BITSEA-manual (Briggs-Gowan & Carter, 2006). Also, for this study, we used age group named 36 months instead of 35 months and 30 days for more concise expression.

In a large birth cohort study by Briggs-Gowan (2001), the BITSEA demonstrated good psychometric properties as an early screening tool for assessing social-emotional/behavioural problems and delays in social-emotional competence. The BITSEA (Briggs-Gowan & Carter, 2006) was translated into Finnish by Educational Psychologist, Varpu Penninkilampi-Kerola, Ph.D. and back-translated by an official translator. The original and the back-translated versions were subsequently compared by a native English speaking clinician, Professor Alice Carter, Ph.D. (also an author of the BITSEA), and minor changes were made to the Finnish version. Our study represents the first research, employing the BITSEA in a Finnish sample.

#### **4.2.4 Child Behaviour Checklist (CBCL)**

The CBCL for ages 1½ to 5 years (preschool form) is part of the Achenbach System of Empirically Based Assessment (ASEBA, Achenbach & Rescorla, 2000) and evaluates the psychiatric problems of toddlers and young children. In this study, we used the Finnish version of the CBCL. In the CBCL-questionnaire, the items are rated: “not true” (0), “somewhat or sometimes true” (1) and “very true or often true” (2). The CBCL consists of problem items and the sum of all 100 CBCL items forms the *Total Problems* scale, which can be further divided into *Internalizing*, *Externalizing* and *Sleep Problems* subscales. The *Internalizing* subscale can be further divided into four subscales (i.e., *Emotionally reactive*, *Anxious/Depressed*, *Somatic complaints* and *Withdrawn*), and the *Externalizing* subscale can be subdivided into two subscales (*Attention problems* and *Aggressive behaviour*). In study IV, we report only the *CBCL Total Problems*, *Externalizing* and *Internalizing* scores results, when examining the association between the CBCL and the parenting stress, PSI. This was done in order to 1) maximize the power of our analyses, and 2) avoid the role of multicollinearity from the overlapping items in the many subscales of the CBCL.

The CBCL for ages 1½-5 years has been used widely and internationally as a psychiatric assessment of children in both clinical and research pursuits (Achenbach, 2001). In the U.S., the clinical cut-off score for the *Total problems* scale is 93% of the sample (T-score of 65); however, the cut-off scores vary cross-culturally (Rescorla *et al.* 2007b). Since Finland lacks normative data on the use of the ASEBA methodology, the American norms provided by the ASEBA-manual and were used in this study (ASEBA, Achenbach & Rescorla, 2000).

#### **4.2.5 Parenting Stress Index Short-Form (PSI)**

The PSI is a 36 item, Likert-type parent self-report scale (i.e., rated from 1 = I fully agree to 5 = I fully disagree) employed to evaluate parental stress (Abidin, 1999). The PSI *Total stress* score is a sum score of all 36 questions (minimum = 36 points, maximum = 180 points). The PSI includes three subscales: *Parental Distress* (12 items), *Parent-Child Dysfunctional Interaction* (12 items) and *Difficult Child* (12 items). It can be used by clinicians and researchers working with parents and children to identify stressors that are most commonly associated with dysfunctional parenting. The PSI was developed for use as a screening and diagnostic assessment technique. It is appropriate for parents of children as young as

one month old and has been used in several empirical studies concerning parent-child relationships (e.g., Browne & Talmi, 2005, Mäntymaa *et al.* 2006, Saisto *et al.* 2008). In this study, we evaluated all PSI scales.

#### **4.2.6 Family based enrichment-intervention**

Fourteen families from phase I (year 2006) had the opportunity to participate in a family-based intervention, which gathered every second week for five months. Three groups of children with deficient or enhanced communication skills measured via parent-report MCDI were established, and consisted of 4–5 child-parent-pairs (the total intervention sample included 14 parent-child-pairs, 4 boys and 10 girls). Each session involving active singing and playing lasted from 22 to 35 minutes; ten to 20 minutes were available for the use of free conversation. Parents often shared their experiences about parenthood and their children in these free situations. New plays and songs were practised together at two sessions and there were five different variables of interest observed and recorded in each of the ten sessions: *eye contact, eye hand coordination and motor movements, joint attention, memory skills and visual-spatial awareness*. Every session began and ended with same routines: singing a song and introducing participants' names at the beginning of each session, and farewell song at the end of each session. Hand-clapping and play was included in each session.

The intervention programme was designed from speech and language therapists' practical work and along with earlier studies of children's developmental phases and milestones (see e.g., Piaget & Inhelder 1977, Howlin, 1984, Tomasello, 1988, Best, Melvin & Williams 1993, Rossetti, 1996, Överlund, 1996, Tolonen, 1996, Carpenter, 1998, Nurkkala, 1998, Laakso *et al.* 1999, Lyytinen, 1999a, Markus *et al.* 2000, Morales *et al.* 2000, Smith, Groen & Wynn, 2000, Bloom & Tinker, 2001, Kivijärvi *et al.* 2001, Briggs-Gowan & Carter, 2006) in support of children's normative developmental goals.

The primary aim of the intervention was to maintain and develop parent-child-interaction via familiar action songs and plays (where child and parent interact via singing and playing), with a research focus on interactive shared and joint attention (parent and child singing and playing together) and rhythmical cooperation (clapping songs, plays and kinaesthetic coordination) of the parent-child dyad. Previous studies of infant social and motor skills and symbolic play development (e.g., Best, Melvin & Williams, 1993, Lyytinen *et al.* 1999b, Viholainen *et al.* 2002, Reilly *et al.* 2006) were considered when planning the inter-

vention program. Many traditional Finnish songs and plays combine elements of eye-hand coordination (especially hand-movements crossing the body midline) and spatial awareness; therefore, practice using these tools may facilitate infant interaction and communication in the parent-child-relationship.

#### **4.2.7 Statistical methods**

All statistical analyses were performed using the SPSS 17.0 statistical software programme for Windows and for the Macintosh. When variables were not normally distributed, non-parametric tests of significances (i.e., Mann-Whitney U-test, Kruskal-Wallis t-test and Spearman's correlation coefficient) were used to evaluate associations between measures of interest and age groups. Moreover, we completed heuristic analyses to consider appropriate demographic covariates (e.g., infant gender, parental education, number of siblings living in the household). When statistically appropriate, we conducted parametric tests, such as Pearson correlation coefficients, repeated measures Analyses of Variance (ANOVA) and one-way analyses of covariance (ANCOVA). All p-values are reported as two-tailed values.

## 5 Results

### 5.1 Sample

Families' education levels in this sample varied from lower secondary level ( $n = 8$ , 18.6%) and upper secondary level ( $n = 16$ , 37.2%) to tertiary levels ( $n = 19$ , 44.2%) following the typical educational trends in Finland's Oulu region (Official Statistics of Finland 2011). Seven families did not report their education. All children in this study had both Finnish-speaking parents living with them and had 0–2 siblings (average number of children per family  $M = 2.2$ ,  $SD = 0.75$ ).

Evaluative assessments (the MCDI, Fenson *et al.* 1994, Finnish version by Lyytinen, 1999a), background information questions (i.e., parental education and number of siblings) and the PSI-Short -Form (Abidin *et al.* 1999) were completed by 50 participants at the age of 8 months. The developmental questionnaires (the MCDI, the BITSEA and the CBCL) were mailed to the abovementioned participants from the 8-month old sample ( $N = 50$ , 16 boys and 34 girls), at the age of 18 months ( $M = 18.1$  months,  $SD = 0.5$ ), and 48 participants (response rate 96%, 16 boys and 32 girls) returned completed questionnaires.

All developmental questionnaires (the BITSEA, the CBCL and the MCDI) were mailed again to the parents of 50 children at the child's age of approximately 36 months ( $M = 36.7$  months,  $SD = 0.3$ ). Forty parents (response rate 80%, 14 boys and 26 girls) returned the completed follow-up questionnaires.

### 5.2 Reliability of measures

In this study, results of the BITSEA suggest acceptable internal consistency for the total sample ( $N = 48$ ). Internal consistency for the BITSEA *Problem Total* scores was acceptable at both the 18 months ( $N = 48$ ,  $\alpha = 0.68$ ) and 36 months assessment points ( $N = 40$ ,  $\alpha = 0.60$ ). For the BITSEA *Competence* scores, the internal consistency was mediocre at the age of 18 months ( $N = 48$ ,  $\alpha = 0.57$ ) and acceptable at the age of 36 months ( $\alpha = 0.73$ ). The BITSEA *Externalizing* subscale yielded fair internal consistency at both the ages of 18 and 36 months ( $\alpha = 0.66$  and  $0.67$ , respectively). The internal consistency of the BITSEA *Internalizing* and *Dysregulation* subscales varied from poor to very poor at baseline and follow-up (*Internalizing*,  $N = 48$ ,  $\alpha = 0.38$ ,  $N = 40$ ,  $\alpha = 0.16$ , respectively); (*Dysregulation*,  $N = 48$ ,  $\alpha = 0.54$ ,  $N = 40$ ,  $\alpha = 0.34$ , respectively). We also calcu-

lated the internal consistency for the BITSEA *ASD* and *Red Flag* scales despite evidence of broad variation of these items noted in the BITSEA manual (Briggs-Gowan & Carter, 2006). The Chronbach's alphas for the *ASD* scale at the ages of 18 and 36 months were 0.55 and 0.48, respectively. The internal consistency for *Red Flag* items at the age of 18 and 36 months varied from poor to very poor ( $\alpha = 0.46$  and  $\alpha = 0.37$ , respectively). Due to the poor internal consistency for many of the BITSEA subscales in our sample, we report further results below primary using the BITSEA *Problem Total* scale and *Competence* scales, but all results are reported in the tables in order to demonstrate associations between other measures.

For the CBCL in our sample, the internal consistency was excellent for the CBCL *Total problem* scores at the age of 18 months ( $N = 50$ ,  $\alpha = 0.91$ ) and good at the age of 36 months ( $N = 37$ ,  $\alpha = 0.88$ ). Internal consistency for the CBCL *Externalizing* and *Internalizing* sum scores varied from good to acceptable at the age of 18 months ( $\alpha = 0.84$ ;  $\alpha = 0.74$ ) and at the age of 36 months varied from good to fair ( $\alpha = 0.85$  and  $\alpha = 0.65$ ). For the CBCL subscales (*Emotionally reactive*, *Anxious/Depressed*, *Somatic complaints*, *Withdrawn*, *Sleep problems*, *Attention problems* and *Aggressive behaviour*), the internal consistency ranged from unacceptable to good at the age of 18 months ( $N = 50$ ,  $\alpha = 0.30$  to  $0.86$ ) and at the age of 36 months ( $N = 37$ ,  $\alpha = 0.27$  to  $0.88$ ). The CBCL *Aggressive behaviour* subscale demonstrated the highest internal consistency ( $\alpha = 0.88$ ) and *Anxious/Depressed* yielded the lowest internal consistency ( $\alpha = 0.27$ ) in both age groups (18 and 36 months).

The PSI demonstrated, excellent internal consistency for the PSI *Total Stress* score ( $N = 50$ ,  $\alpha = 0.91$ ), and good for each of its subscales (*Parental Distress*, *Parent-Child Dysfunctional interaction* and *Difficult Child*) ( $N = 50$ ,  $\alpha = 0.83$  to  $0.85$ ).

For the PC-ERA, there were two raters, who evaluated the tapes and were trained by psychologist Sari Ahlqvist Björkroth, MPsych, who is an official trainer for the PC-ERA in Finland. For 23 mother-infant -dyads, six randomly selected tapes were evaluated by two raters, in order to establish reliability for the evaluations. The mean percentage of agreement computed from all 64 variables was 85% (when collapsing the variables across the three categories: *Parent variable*, *Infant variable* and *Dyadic variable*); the agreements varied from 74–97%.



### 5.3 Communicative development (MCDI questionnaire) in children aged 8 months to 36 months (Study I)

We sought to evaluate infant communicative development with the Finnish version of the MCDI and to compare groups (intervention versus no-intervention) in a longitudinal follow-up study (i.e., from eight to 36- months).

Parents completed the MCDI questionnaires, when the child was approximately eight months ( $M = 8.2$  months,  $SD = 0.3$ ), 18- months ( $M = 18.1$  months,  $SD = 0.5$ ), 24- months ( $M = 23.8$  months  $SD = 0.3$ ), and 36- months ( $M = 36.7$  months,  $SD = 0.3$ ) old. At the age points of 18, 24 and 36- months, the MCDI was mailed to all 50 participating families from the years 2006 and 2007. At the age of 18- months, 50 participants returned the MCDI; two cases returned empty questionnaires, thus the final sample of 18-month old children was 48 (response rate 94%, 16 boys and 32 girls). At the age of 24- months ( $M = 23.8$  months  $SD = 0.3$ ), 44 families returned the MCDI (response rate 86.3%, 15 boys, 29 girls). When the child was 36- months old, 37 families completed the MCDI (response rate 72.5%, 12 boys, 25 girls) questionnaire.

When children were 8 months old, the MCDI *Total scores'* descriptive were:  $M = 6.4$ , range 1–12, (girls  $M = 6.7$  range 1–12, boys  $M = 5.7$ , range = 2–12). The total 8 month old sample had the highest scores in the *Communicative gestures and actions* and *Comprehension* subscales (both  $M = 2.4$ ,  $SD = 1.7$  and  $0.7$ ), while for the 18-month old toddlers, the highest scores for the total sample were on the *Communicative gestures and actions* ( $M = 14.8$ ,  $SD = 1.25$ ) subscale.

At the age of 24- months, there was considerable variability in children's communicative skills; the mean *Total score* of the sample was 341.5 (girls  $M = 403.3$ , range = 158–559; boys  $M = 274.4$ , range = 2–487).

At the age of 36- months, children's communicative skills (*Total scores*) varied from 349 to 605; no child reached the maximum possible score of 621 points. Further, at this age, girls' scores,  $M = 570$ , 5, ranged from 445–605 and boys' scores,  $M = 525.0$ , ranged from 349–605. The widest range between subscales and gender was in the *Plural and grammar* subscale ( $M = 16.8$ , range = 5–18, girls scores  $M = 17.6$ , range = 12–18, boys  $M = 15.4$ , range = 5–18), while lowest range was in the *Pragmatics* subscale ( $M = 9.7$  range = 7–10, girls  $M = 9.7$ , range = 9–10, boys  $M = 9.5$ , range = 7–10).

Finally, there was a strong association between children's gender (girls had higher scores), communicative development and parent's education at the age of 24 and 36 months, our sample. Gender (female) and better communicative devel-

opment was statistically significantly related  $r = .520$ ,  $p < .01$  at the age of 24-months and at the age of 36 months  $r = .421$ ,  $p < .01$ . Results of the communicative development based on the MCDI are presented in Table 3.

**Table 3. Children's communicative development via MCDI (short version at the age of 8 and 18 months and standard version at 24 and 36 months of age).**

MCDI	8 m (short version),mean, SD (min-max)	18 m (short version),mean, SD (min-max)	24 m, mean, SD (min-max)	36 m, mean, SD (min-max)
Total scores (sum of first words and comprehension)	6.4, 2.8 (1-12)	14.8, 1.25 (11-17)	341.5, 156.7 (4-559)	554.5, 72.4 (349-605)
Communicative gestures and actions	2.4, 1.7 (0-6)	7.9, 1.2 (5-10)	N/A	N/A
Communicative action with objects	1.5, 1.2 (0-4)	3.9, 0.2 (3-4)	N/A	N/A
First words	2.4, 0.7 (1-3)	3.0, 0.0 (3)	N/A	N/A
Pragmatics	N/A	N/A	7.9, 2.0 (2-10)	9.7, 0.6 (7-10)
Plural and suffixes	N/A	N/A	10.4, 6.1 (0-18)	16.8, 3.0 (5-18)
Verbs	N/A	N/A	8.4, 4.4 (1-14)	13.0, 1.6 (8-14)

m = months

N/A= subscales not included in MCDI versions for that age group

### **5.3.1 Children's communicative development and PC-ERA**

We also wanted to objectively evaluate children's interactive skills; therefore we examined the associations between the PC-ERA and the MCDI. Specifically, we analyzed all variables (*Parent variables*, *Infant variables* and *Dyadic variables*) from 23 participants, whose video material provided sufficient data (same parent) from both video assessments (first, in children eight to 12 months old, and second, in children 15 to 18 months old).

There were strong associations between the 18-month old children's MCDI results and eight- month olds' PC-ERA *Parent variable* ( $r = .595$ ,  $p < .01$ ), *Infant variable* ( $r = .551$ ,  $p < .05$ ) and *Dyadic variable* ( $r = .603$ ,  $p < .01$ ). A statistically significant negative relationship emerged between the 18-month old children's scores on the PC-ERA *Dyadic variable* and 18-month old children's scores on the

MCDI ( $r = -.468, p < .05$ ). There were no other significant associations between the MCDI and the PC-ERA for other child age groups.

Interestingly, the number of siblings in the family was significantly associated with the following eight- month old PC-ERA subscales: 1) *Parent* variables ( $r = .50, p < .05$ ), 2) *Infant* variables ( $r = .67, p < .01$ ), and 3) *Dyadic* variables ( $r = .50, p < .05$ ). Of note, infants without siblings in the family scored lower than infants with siblings on the *Infant* variables ( $M = 108.0$  vs.  $M = 125.4, p < .01$ , respectively). The association between parent education level and 36- month old children's MCDI scores approached significance ( $r = .33, p = .055$ ) as did the association between PC-ERA *Parent* variables and parent education level at the age of eight months ( $r = .45, p = .054$ ). All correlations between children's communicative development and the PC-ERA are presented in table 4.

**Table 4. Associations between children's communicative development (MCDI short version and standard version) and the PC-ERA.**

Measure	Variable	MCDI	MCDI	MCDI	MCDI
		(short version) 8 months	(short version) 18 months	24 months	36 months
PC-ERA 8 months	Parent	.296	.595**	.360	.330
	Infant	.279	.551*	.196	.146
	Dyadic	.358	.603**	.233	.211
PC-ERA 18 months	Parent	-.242	-.332	.141	.001
	Infant	-.015	-.426	-.072	-.144
	Dyadic	-.254	-.468*	-.080	.009

\*\* $p < .001, *p < .05, 2$ -tailed

**5.3.2 Communicative development and family-based intervention**

We offered a family-based intervention to a random sample of children ( $n = 14$ ), who had poorer or better communicative skills. We conducted a median split based on MCDI *Total* scores (median = 7), and grouped our sub-sample, accordingly. Results from the intervention group and no-intervention groups are presented in table 5.

**Table 5. MCDI Total scores according to family-intervention (at infant's age of 8 months) status.**

Measure	Intervention group	No-intervention group
	(n = 14-11) mean (SD) min-max	(n = 36-25) mean (SD) min-max
MCDI (short version) 8 months	7.8 (3.5) 3-12	5.5 (2.0) 2-9
MCDI (short version) 18 months	15.1 (1.3) 12-17	14.7 (1.3) 11-17
MCDI 24 months	349.5 (163.3) 26-559	337.2 (154.1) 4-533
MCDI 36 months	550.0 (50.3) 418-605	542.7 (83.4) 302-605

Based on the eight- month old MCDI *Total scores*, children from the 2006 data collection (n= 31) were divided into two groups as detailed above: 1) The “High MCDI group” consisted of children, equal to or above the median on the MCDI (4 boys and 11 girls), and 2) The “Low MCDI group” consisted of children, who scored lower than the median on the MCDI (7 boys and 9 girls). Children, who received the intervention and were in the High MCDI group, demonstrated higher MCDI scores at 18- month follow-up, than did children in the High MCDI group, who were in the no-intervention group; however, this difference was not statistically significant. Descriptive histograms based on intervention status and MCDI *Total scores* are presented in Figure 2.





Fig. 2. MCDI groups (High = good communication skills, Low = low communication skills) and intervention status (m = months).

### 5.3.3 Social-emotional skills (the BITSEA) and behavioural problems (the CBCL) by MCDI groups


We examined the association between children's communication development, as measured by the MCDI, and social-emotional and behavioural problems and competencies, as parent reported on the BITSEA and the CBCL. We, conducted a tertiary split based on mean MCDI *Total scores*, at both assessment points (18 months and 36 months): MCDI *Low* group (n = 13), MCDI *Average* group (n = 20), and MCDI *High* group (n = 13) in order to look at the relationship between *Low*, *Average* or *High* communicative skills group with social-emotional or behavioural problems. The MCDI skill groups differed statistically on the BITSEA *Competence* subscale at the age of 18 months. Specifically, the MCDI *High* (n = 13, M= 18.9) group scored higher than both the *Low* (n = 13, M = 16.6) and *Average* (n = 20, M = 18.1) MCDI skill groups ( $\chi^2 = 7.7$ , df = 2, p < .05). Moreover, children who differed according to the MCDI skill groups at 18 months, varied statistically on the CBCL *Withdrawn* subscale at 36 months. The MCDI *High* (n = 9, M = 19.11) group scored lower than did the other MCDI groups (MCDI *Average* n= 17, M = 18.94, MCDI *Low* n = 9, M = 16.3;  $\chi^2 = 6.5$ , df = 2, p < .05).

Comparisons between the MCDI groups and the BITSEA *Competence* subscale are presented in Figure 3.



**Fig. 3. MCDI groups and BITSEA Competence scores.**

The MCDI groups at 36 months, differed statistically on BITSEA *Competence* scores completed at the age of 18 months, with the *Low* MCDI group scoring lower on the BITSEA *Competence* subscale than the other two MCDI groups ( $\chi^2 = 11.9$ ,  $df = 2$ ,  $p < .01$ ). The MCDI groups at the follow-up assessment point differed on the BITSEA *Competence* and on the CBCL *Attention problems* subscales completed at 36 months. Specifically, the MCDI *Low* group scored lower on the BITSEA *Competence* subscale and higher on the CBCL *Attention problems* subscale than the other MCDI groups ( $\chi^2 = 6.8$ ,  $df = 2$ ,  $p < .05$ ;  $\chi^2 = 6.1$ ,  $df = 2$ ,  $p < .05$ , respectively). There were no additional significant differences between the MCDI skill groups on the BITSEA or the CBCL. Comparisons between the MCDI and CBCL are reported in Figure 4.



**Fig. 4. Comparisons of CBCL mean scores at ages 18 months and 36 months between the MCDI groups at both assessment points (m = months).**

#### **5.4 Social-emotional, behavioural problems and competencies in children aged 18 and 36 months (Studies II and III)**

Children's social-emotional and behavioural development was screened at the age of 18 months and 36 months with the BITSEA and the CBCL.

Ten percent of 18-month old children (3 boys and 2 girls) and five percent of 36-month old children (2 girls) scored above the original BITSEA *Problem Total* score cut-off (cut-off scores = those falling within the 75<sup>th</sup> percentile according to the BITSEA-manual, Briggs-Gowan & Carter, 2006). In addition, 10% of 18-month old children (2 girls and 3 boys) and five percent of children of 36-month old (2 girls) scored below the BITSEA *Competence* subscale cut-off (85<sup>th</sup> percentile) indicating possible deficits or delays in social-emotional competencies (Briggs-Gowan & Carter, 2006).

BITSEA descriptive results (scores, means and standard deviations and clinical cut-off points) for children ages of 18 months and 36 months are presented in table 6.

**Table 6. Descriptives (n = 48 at 18 months and n = 37 at 36 months) for the BITSEA and the CBCL *Total problem* scores and subscales.**

Measure (min-max)	Mean (SD) at 18 months (n = 48)	Min-Max scores at 18 months	Mean (SD) at 36 months (n = 37)	Min-Max scores at 36 months
BITSEA Problem Total score (0-100)	7.20 (4.75)	0-18	4.45 (3.18)	0-15
BITSEA Competence (0-22)	17.97 (2.20)	13-21	19.35 (2.51)	12-22
CBCL Total problem scores (0-200)	21.95 (12.63)	0-66	16.4 (9.90)	0-36
CBCL Internalizing (0-72)	3.57 (3.27)	0-14.8	3.05 (2.72)	0-9
CBCL Externalizing (0-48)	10.78 (5.95)	0-29	7.9 (4.62)	0-17
CBCL Emotionally reactive (0-18)	1.8 (1.5)	0-6	1.2 (1.32)	0-5
CBCL Anxious/Depressed (0-16)	0.7 (0.9)	0-4	0.60 (0.77)	0-3
CBCL Somatic complaints (0-22)	0.6 (1.39)	0-8.8	0.88 (1.36)	0-8
CBCL Withdrawn (0-16)	0.31 (0.69)	0-3	0.37 (0.84)	0-3
CBCL Sleep problems (0-14)	1.89 (2.27)	0-10	1.76 (1.67)	0-8
CBCL Attention problems (0-10)	2.04 (1.19)	0-4	1.21 (1.20)	0-4
CBCL Aggressive behaviour (0-38)	8.74 (5.09)	0-25	6.66 (4.15)	0-15

#### **5.4.1 Associations between age groups (18 months and 36 months) and the BITSEA, CBCL and MCDI (studies II and III)**

The BITSEA and the CBCL were consistent in their interpretable results concerning children's social-emotional and behavioural development. There was a statistically significant association between the BITSEA *Problem Total* scores and the CBCL *Total problem* scores as well as between the individual subscales of these measures for both age groups (see tables 7 and 8). There was a moderate association between the MCDI and the BITSEA *Competence* subscale at 18 months. Further, there was a moderate association between both age groups on the BITSEA *Problem Total* and *Competence* scores ( $r = 0.63$ ,  $p < .001$ ;  $r = 0.58$ ,  $p < .001$ , respectively). Similarly, a statistically significant relationship emerged for the BITSEA *ASD* items between children at both assessment points ( $r = 0.43$ ,  $p < 0.001$ ).



The CBCL yielded strong to moderate associations between 18-month and 36-month old children on the CBCL *Total problem* score, *Externalizing* and *Internalizing* as well as *Sleep problem* subscales ( $r = 0.74, p < .001$ ;  $r = 0.60, p < .001$ ;  $r = 0.58, p < .001$ ;  $r = 0.53, p < .001$ , respectively). There were no other statistically significant associations between age group and CBCL subscales. Finally, we noted a significant relationship between child's age group and MCDI scores ( $r = 0.40, p < .05$ ). The results revealed a considerable relationship between the MCDI *Total score* and the BITSEA *Competence* items for children aged 36 months. No other statistically significant relationships between these measures and their subscales emerged. All correlations between the BITSEA, the CBCL and the MCDI at the age of 18 months are presented in table 7 and at the age of 36 months in table 8.

**Table 7. Correlations between the BITSEA, CBCL and MCDI at 18 months.**

Measure at 18 months	BITSEA Problem		BITSEA		BITSEA		BITSEA		BITSEA ASD	BITSEA Red Flag
	Total	Externalizing	Internalizing	Dysregulation	Competence					
CBCL Total problem score	0.630**	0.440**	0.555**	0.451**	-0.151	0.357*	0.518**			
CBCL Internalizing	0.457**	0.276	0.572**	0.309*	-0.108	0.350*	0.363*			
CBCL Externalizing	0.556**	0.466**	0.390**	0.341*	-0.173	0.303*	0.496**			
CBCL Emotionally reactive	0.487**	0.417**	0.515**	0.253	-0.158	0.303*	0.377**			
CBCL Anxious/Depressed	0.393**	0.145	0.554**	0.344*	-0.115	0.283	0.253			
CBCL Somatic complaints	0.164	-0.011	0.259	0.163	0.175	0.157	0.202			
CBCL Withdrawn	0.19	0.143	0.281	0.098	-0.314	0.231	0.092			
CBCL Sleep problems	0.648**	0.303*	0.500**	0.726**	0.002	0.176	0.420**			
CBCL Attention problems	0.426**	0.450**	0.268	0.141	-0.089	0.207	0.499**			
CBCL Aggressive behaviour	0.550**	0.439**	0.393**	0.365*	-0.181	0.306*	0.463**			
MCDI Total scores	0.014	-0.023	0.051	0.033	0.379**	-0.102	0.058			

\* p < 0.05 level (2-tailed)

\*\* p < 0.01 level (2-tailed)

**Table 8. Correlations between the BITSEA, CBCL and MCDI at 36 months.**

Measure at 36 months	BITSEA Problem		BITSEA Internalizing	BITSEA Dysregulation	BITSEA Competence	BITSEA ASD	BITSEA Red Flag
	Total	Externalizing					
CBCL Total problem score	0.642**	0.427**	0.438**	0.610**	-0.191	0.267	0.307
CBCL Internalizing	0.437**	0.264	0.266	0.416**	-0.014	0.202	0.290
CBCL Externalizing	0.672**	0.517**	.408**	0.570**	-0.277	0.297	0.369*
CBCL Emotionally reactive	0.288	.000	0.208	0.373*	0.117	0.038	0.187
CBCL Anxious/Depressed	0.178	-0.095	0.304	0.318*	0.204	-0.058	0.012
CBCL Somatic complaints	0.419**	0.511**	0.023	0.263	-0.055	0.350*	0.350*
CBCL Withdrawn	0.117	0.113	0.215	0.04	-0.331*	0.472**	0.065
CBCL Sleep problems	0.581**	0.253	0.395*	0.712**	-0.095	0.099	0.183
CBCL Attention problems	0.247	0.188	0.205	0.016	-0.074	0.167	0.417**
CBCL Aggressive behaviour	0.681**	0.526**	0.402*	0.630**	-0.279	0.283	0.294
MCDI Total scores	0.005	-0.217	0.038	0.088	0.541**	-0.536**	0.173

\* p < 0.05 level (2-tailed)

\*\* p < 0.01 level (2-tailed)

Girls (n = 31) in our sample scored higher than did boys (n = 16) on the CBCL *Attention problems* subscale at the age of 18 months (M = 2.3 vs. 1.5, Z = 2.3, p < .05). Of note, there was one 18-month-old child (girl) and two 36-month-old children (both girls), who met criteria for possible deficit or delay on the CBCL *Total scores* (93% cut-off, according the manual; ASEBA, Achenbach & Rescorla, 2000). Boys (n= 14) scored higher than did girls (n = 26) on the CBCL *Withdrawn* subscale at the age of 36 months (M = 0.93 vs. .08, Z = 2.9, p < .01).

Our analyses revealed a statistically significant association between the MCDI *Total scores* and the BITSEA *Competence* items for 36- month old children. No other statistically significant relationships between these measures and their subscales emerged.

### 5.5 Maternal stress at infant age of 8 months and children's behavioural problems at the ages of 18 months and 36 months (Study IV)

The aim of this particular study was to evaluate the associations of self-reported maternal stress (the PSI in infants' aged 8 months) and children's behavioural problems (the CBCL *Total problem scores* and *Internalizing* and *Externalizing* subscales) in children's aged 18 and 36 months. The results of all measures are presented in table 9.

**Table 9. Means, standard deviations (SD) and minimum and maximum scores on the PSI and the CBCL.**

Measure	Mean	SD	Minimum	Maximum
PSI Total score (N=48) 8 months	67.16	15.39	46.00	111.00
PSI Parental Distress (N=49) 8 months	24.18	7.25	13.00	41.00
PSI Parent-Child Dysfunctional Interaction (N=50) 8 months	17.20	4.90	12.00	33.00
PSI Difficult Child (N=48) 8 months	26.04	6.20	18.00	42.00
CBCL Total problem scores (N=50) 18 months	21.95	12.63	0.00	66.00
CBCL Internalizing scores (N=50) 18 months	3.57	3.27	0.00	14.80
CBCL Externalizing scores (N=50) 18 months	10.78	5.95	0.00	29.00
CBCL Total problem scores (N=37) 36 months	16.40	9.90	0.00	36.00
CBCL Internalizing scores (N=37) 36 months	3.05	2.72	0.00	9.00
CBCL Externalizing scores (N=37) 36 months	7.90	4.62	0.00	17.00

### **5.5.1 Maternal stress level**

#### *Heuristic Analyses*

We completed heuristic analyses to consider appropriate demographic covariates (e.g., infant gender, parental education, number of siblings). Neither infant gender ( $n = 48$ ,  $r = -.003$ ,  $p = ns$ ) nor parent education ( $n = 41$ ,  $r = -.074$ ,  $p = ns$ ) was statistically significantly associated with the PSI. Number of siblings in the family, was significantly associated with the PSI ( $n = 41$ ,  $r = .309$ ,  $p < .05$ ). This relationship is consistent with other recent research studies (e.g., Yaman *et al.* 2010, Sperlich, Arnhold-Kerri & Geyer 2011).

To study maternal stress level in this sample, we included all completed questionnaires ( $N = 48$ ) and divided mothers into two groups based on PSI *Total* scores, in order to optimize statistical power in our analyses. The two groups were divided via median split, Median = 65.5. The *High* group consisted of mothers reporting a PSI score  $\geq 65.5$ , and the *Low* group consisted of mothers reporting a PSI score  $< 65.5$  in addition to mothers considered “defensive” on the PSI (i.e., those underscoring or denying their problems). PSI groups are utilized in subsequent statistical analyses to examine the effects of various levels of maternal stress on the behavioural development in children and observable parent-child-interaction style.

### **5.5.2 Associations between the PSI and the CBCL**

The associations between the PSI *Total Stress* scores and the CBCL *Total problem* scores, *Internalizing* and *Externalizing* subscales at the age of 18 months are presented in table 10 and at the age of 36 months in table 11.

**Table 10. PSI (at child's age of 8 months) and associations with the CBCL at 18 months (N = 46).**

PSI subscales	CBCL, problem subscale	Pearson correlation coeff.
Parental distress	Internalizing	.592**
	Externalizing	.603**
	Total problem scores	.654**
Parent-Child Dysfunctional Interaction	Internalizing	.274
	Externalizing	.278
	Total problem scores	.314*
Difficult Child	Internalizing	.240
	Externalizing	.369*
	Total problem scores	.363*
PSI Total scores	Internalizing	.447**
	Externalizing	.499**
	Total problem scores	.532**

\* $p < 0.05$  level (2-tailed)

\*\* $p < 0.01$  level (2-tailed)

**Table 11. PSI (at child's age of 8 months) and associations with the CBCL at 36 months (N = 37).**

PSI subscales	CBCL, problem subscale	Pearson correlation coeff.
Parental Distress	Internalizing	.354*
	Externalizing	.378*
	Total problem scores	.398*
Parent-Child Dysfunctional Interaction	Internalizing	.183
	Externalizing	.078
	Total problem scores	.123
Difficult Child	Internalizing	.043
	Externalizing	.205
	Total problem scores	.186
PSI Total scores	Internalizing	.207
	Externalizing	.256
	Total problem scores	.274

\* $p < 0.05$  level (2-tailed)

### **5.5.3 Longitudinal associations between PSI groups (High and Low) and CBCL Total problem scores, Internalizing and Externalizing subscales.**

#### *Maternal stress level and number of siblings in the family*

We examined the association of the PSI groups (*High vs Low*) ( $N = 50$ ) and the CBCL ( $N = 46$  at the age of 18 months and  $N = 37$  at the age of 36 months) over time. For each of the three primary outcome variables (*CBCL Total problem score*, *Internalizing score*, *Externalizing score*) we employed 2 x 2 Repeated Measures ANOVAs, with PSI status (*High versus Low*) as a fixed factor and CBCL assessment time (18 and 36 months) as a repeated factor.

#### *Associations of PSI subgroups and CBCL Total problem score, Internalizing and Externalizing scores and number of siblings*

There were moderately significant associations between mothers' ratings of parenting stress (PSI) when their children were 8 months old and their ratings of children's CBCL *Total problem score*, *Internalizing*, and *Externalizing* problems at 18 months ( $r = .53$ ,  $r = .45$ ,  $r = .50$ ,  $p < .01$  for all), but not at 36 months ( $r = .27$ ,  $r = .21$ ,  $r = .26$ ,  $p = \text{ns}$ ).


### **5.5.4 The association of the PSI with the CBCL Total problem score over controlling for number of siblings**

In the next three models, we used repeated measure analysis of covariance (with number of siblings as the covariate, time of assessment (18 vs. 36 months) as the within subjects effect, and PSI group (*High vs. Low*) as the between subjects factor).

For the first model, 16 participants were excluded from the final analyses due to missing data (i.e., CBCL Total problem score at the 36 month follow-up, or number of siblings). Fifteen *Low* and seventeen *High* PSI participants were included in the following analyses.

Results indicated a significant multivariate model within-subjects main effect of time, Wilks'  $\Lambda = .79$ ,  $F(1, 29) = 7.7$ ,  $p < .01$ .  $\eta_p^2 = .21$ , such that CBCL *Total problem* scores were significantly higher at the 18- versus 36-month assessment. There was also a statistically significant between-subjects effect on PSI status,

indicating that CBCL *Total problem* scores were statistically significantly higher in *High* PSI mothers when compared to *Low* PSI mothers, ( $F(1, 29) = 15.9$ ,  $p < .001$ ,  $\eta_p^2 = .36$ ), (see Figure 5). The interaction between age, CBCL assessment time and PSI group was not significant, Wilks'  $\Lambda = .90$ ,  $F(1, 29) = 0.03$ ,  $p = ns$ ,  $\eta_p^2 = .00$ . Results suggest that a higher level of maternal reported stress at 8 months after the birth of a child is associated with an increase in parent report of total problematic behaviours in their child across time, and in general parent report of child emotional and problem behaviours decreases over time regardless of mother's self-reported stress. Results of the CBCL *Total problem scores* at the age of 18 months and 36 months is presented in Figure 5.



**Fig. 5. Results from CBCL *Total problem* scores and the PSI groups at the age of 18 and 36 months (controlling for number of siblings).**

\*\*\*PSI group: High > Low,  $p < .001$

\*\* Time: 18 months > 36 months,  $p < .01$

PSI x Time = ns



### 5.5.5 The association of the PSI with the CBCL Internalizing scores over time, controlling for number of siblings.

In this model, 16 participants were excluded from the final analyses due to missing data (i.e., CBCL *Internalizing* score at the age of 36 months or number of siblings). Fifteen *Low* PSI and seventeen *High* PSI mothers were included in the second model. PSI group was subsequently examined in association to CBCL *Internalizing* scores over time. There was neither a main effect of assessment time, Wilks'  $\Lambda = .99$ ,  $F(1, 29) = .42$ ,  $p = ns.$ ,  $\eta_p^2 = .01$  suggesting that the mean CBCL scores on the *Internalizing* scale did not differ within subjects over time, nor the interaction between CBCL *Internalizing* assessment time and PSI group. As expected, a between subjects difference emerged such that *High* PSI mothers reported a greater number of child internalizing problem behaviours than did *Low* PSI mothers ( $F(1, 29) = 11.1$ ,  $p < .01$ ,  $\eta_p^2 = .23$ , (see Figure 6). These results support the hypothesis that a higher level of parent reported stress is associated with an increase in parent report of internalizing behaviours in children across time. Results are presented in Figure 6.





Fig. 6. CBCL Internalizing subscale by PSI groups at child's the age of 18 months and 36 months (controlling for number of siblings).

\*\*PSI group: High > Low,  $p < .01$

\*\*Time = ns, PSI x Time = ns

### **5.5.6 Model 3: The Association of the PSI with the CBCL Externalizing score over time, controlling for number of siblings.**

In this model 16 participants were excluded from the final analyses due to missing data (i.e., CBCL *Externalizing* score at the age of 36 months or number of siblings). In the third repeated measures ANCOVA model (with PSI group predicting to CBCL *Externalizing* scores at 18 and 36 months), sixteen *Low* PSI and seventeen *High* PSI participants were included in the analysis. Similar to the first model (CBCL *Total problem* scores), results indicated a main effect of time, Wilk's  $\Lambda = .77$ ,  $F(1, 30) = 9.1$ ,  $p < .01$ ,  $\eta^2 = .23$ , demonstrating that mothers were more likely to report a greater number of problematic childhood externalizing behaviours at 18 months versus 36 months across PSI groups. There was also a main effect of PSI status, indicating that CBCL *Externalizing* scores were significantly higher in the *High* versus *Low* PSI group regardless of assessment time  $F(1, 30) = 13.9$ ,  $p < .001$ ,  $\eta^2 = .32$ , (figure 7). The interaction between CBCL assessment time and PSI group was not significant, Wilks'  $\Lambda = .99$ ,  $F(1, 30) = 0.43$ ,  $p = ns$ ,  $\eta^2 = .01$ . Thus, findings indicate that a higher level of parent reported stress is associated with an increase in parent report of externalizing behaviours in children across time, and that in general, parent report of problematic externalizing childhood behaviours decreases over time regardless of self-reported maternal stress at eight months. Associations between PSI group and CBCL *Externalizing* scores are presented in Figure 7.



**Fig. 7. CBCL *Externalizing* subscale by PSI groups at 18 months and 36 months (controlling for number of siblings).**

\*\*\*PSI group: High > Low,  $p < .001$

\*\*Time: 18 months > 36 months,  $p < .01$

PSI x Time = ns



## 6 Discussion

### 6.1 Main results

#### 6.1.1 Communicative development: The MCDI at follow-up

In our full study sample (study I), our shorter version of the MCDI identified 15 infants (7 girls and 8 boys) with possible problems in language development at the age of 8 months, when we scaled results to the standard longer version. These same individuals were later screened out with the BITSEA and the CBCL. Results support previous findings suggesting that infants' communicative skills (vocabulary and comprehension) are closely related to overall behavioural development (Carter *et al.* 2001, Briggs-Gowan *et al.* 2004, Ellingson *et al.* 2004, Tervo, 2007).

Children followed the typical pattern of Finnish language acquisition (e.g., Kunnari, 1997, Laakso *et al.* 1999, Laalo, 1998 and Lyytinen *et al.* 1999b) and the gender differences noted in our study are consistent with earlier findings in the area of language development (e.g., Lyytinen *et al.* 1999b, Paavola *et al.* 2006a, Stolt *et al.* 2007, Mauridsen & Hauschild, 2010), suggesting that males develop communicative skills slightly later than their same-aged female counterparts.

At the age of 24 months, the variation of the children's communicative skills was quite wide on all subscales of the MCDI, which reveals the broad range of individual differences in communication development in children at this age. Some children in our sample obtained scores of 18 points on the *Plural and grammar* subscale by 24 months, whereas others received zero points (boys only) on this same subscale at the same age. Interestingly, our sample mean ( $M = 330.4$ ) on the MCDI *Vocabulary* subscale for children 24 months of age was considerably higher than the mean reported by Laakso and colleagues (1999) ( $M = 265.1$ ). The larger sample size ( $N = 171$ ) in the Laakso *et al.* 1999 study may account for such differences. Differences in vocabulary production at the age of two years in different areas may be associated with numerous explanatory factors and variables. For example, our results revealed a statistically significant association between communicative skills and parent education level and luckily we had parents from all educational levels; so comparing the educational levels was possible.

At the age of 36 months, children's communicative skills (on the MCDI *Total scores* scale) varied from 302 to 605 points. Despite that our mean sample age at this time point was older than the cut-off for the 16 to 30 month old MCDI maxi-

imum score (621 points), this ceiling score was not met by any child in our sample. Professor Lyytinen (1999a and 1999b) and her colleagues, have been studying children's language development with the MCDI in the Finnish population and results have been consistent with other linguistic measures, such as the Reynell Developmental Language Scales (RDLS). Paavola and colleagues (2006a and 2006b), and Laakso and colleagues (1999), have also used the MCDI with good reliability alongside other measures, such as the RDLS. Because the RDLS is usually scored by speech and language therapists, it may have been a useful objective measure for our study; however, we wanted to obtain parent-information data only in order to compare self-report questionnaires and validity between them. The total MCDI is quite time-consuming; for children ages 16 to 30 months, the questionnaire includes 619 items (one question includes two points from maximum sentence length, therefore maximum score is 621); it is therefore possible, that some parents in our sample had time constraints or other influencing factors (e.g., motivation, attention) to complete the entire questionnaire, thus ultimately limiting our ability to collect full follow-up information.

Study I also included a five-month family-based intervention. When planning the study, we found, that there is a lack of randomized controlled trial studies or evidence-based studies investigating the role of family centred enrichment-interventions in children's communicative development. The design of our intervention phase for this study was in part based on earlier studies of children's developmental stages (see section *Family-based enrichment-intervention*, p. 36) and targeted support focused on identified stages of communicative, motor and social skills. Although the trends of the mean MCDI scores suggest the efficacy of our intervention in young children identified as "at risk" for delays in communicative development, the difference was not statistically significant. This pilot sample was simply too small to yield the statistical power necessary to draw conclusions about the effectiveness of our particular intervention in the wider community. Our findings were, however, consistent with those found by Bayer and colleagues (2010), suggesting that a brief parenting programme may be insufficient to prevent the development of externalizing problems or enhance communicative skills in an effective way. In a recent review by Law, Garrett & Nye (2010), results of speech and language therapy interventions indicated that there were no significant differences between clinician administered intervention and intervention implemented by trained parents. They also noted that there were no apparent differences between group and individual interventions. The authors summarized, that,

in general, the use of normal language by peers in a therapeutic intervention had a positive effect on therapy outcome. Wissow and colleagues (2008), however, found that brief communication training had a clear positive impact on children's communication skills impairments, when parents are were advised to support their child by addressing the children's capability to gain skills. In our sample, it seems, that children who were more competent (better scores on the MCDI), benefitted the most from the intervention; specifically, children with better scores on MCDI before the enrichment-intervention continued to have better scores in follow-up compared to those who did not receive the enrichment-intervention. Our goal was to offer support and guidance to parents on improving their child's communicative skills. Further studies with larger samples are needed to better understand our results and to posit cause and effect from such an intervention.

The parental verbal feedback from our intervention was very positive and therefore a more qualitative approach to data analyses may have enhanced interpretability of the results. Families demonstrated significant commitment to attend the enrichment-intervention meetings; 13 of 14 families participated in six or more meetings. Also many parents commented that they gained insight into their children's social skills and were better equipped to support their child's general interactive and peer-related skills. Some mothers formed friendships and met together outside of the intervention sessions, also serving to bolster support. Sharing experiences seemed to be educational and constructive to parents, in particular, parents reported having learned an entirely new way to interact with their child via play and songs. For example, several fathers, who participated in a majority of the sessions reported that it "felt strange" to engage in such activities initially, they indicated that they recognized the importance of these interactions and really enjoyed the activities by the end of the intervention sessions. The abovementioned subjective report of parents, including novel ways to communicate and interact with their child cannot be evaluated via diagnostic assessment tools (e.g., CBCL, BITSEA and MCDI) used in this study, and therefore both descriptive and qualitative viewpoints may be critical to consider when evaluating parents, children, communicative development and social relatedness.

### **6.1.2 Relationship between communicative development (MCDI) and social-emotional problems and competencies (BITSEA) and behavioural problems (CBCL)**

The aim of this study was to present the results of communicative development (study I) and its associations to children's social-emotional areas of need and competencies, and behavioural problems between the ages of eight months and 36 months (Study II and III). Before this study, the BITSEA had not been used in research and clinical settings in Finland, and there is a strong need for early screening measures of social-emotional-problems and language development. We sought to explore the association between questionnaires using a multi-method longitudinal study design.

Our results indicated statistically significant associations of time of assessment (18 months versus 36 months) using the employed measures (i.e., the MCDI revised version for 18-month old children and the MCDI original version for children ages 16–30 months, the BITSEA and the CBCL), thus suggesting some stability over time in children's social-emotional problems and competence, behavioural problems and communicative development. Although, these associations were noted only for certain subscales of the above-mentioned measures in this small sample (e.g., BITSEA *Competence* and CBCL *Attention problems*), the developmental continuum is both empirically evident and clinically relevant; Specifically, if a child manifests deficits in communicative skills and a limited ability to be attentive, there might be more observed behavioural problems, because he/she is unable to express him/herself via typical language patterns. Furthermore, the association between certain subscales (e.g., CBCL *Externalizing*, and BITSEA subscales –with the exception of BITSEA *Competence* –and MCDI *Vocabulary and comprehension*) found in our employed measures suggests that in general, children's communicative skills may reflect aspects of their psychological well-being in addition to their social-emotional competencies. This information is important for both parents and practitioners in terms of early identification and early intervention of potential problem areas.

In study II, the BITSEA cut-off scores for different items were slightly higher than those reported in the BITSEA manual for a U.S. sample at the age of 18 months (Briggs-Gowan & Carter, 2006). The differences between BITSEA *Competence* scores in our sample versus those reported in the manual was unremarkable; however when considering the BITSEA *Problem total* score, our sample



mean was 7.2 while the U.S. sample mean for the same subscale was 9.6 (Briggs-Gowan & Carter, 2006) at the age of 18 months. The mean BITSEA *Competence* scores was 17.98 in our sample, and 17.5 in a U.S. sample (Briggs-Gowan & Carter, 2006). Further, in study II, the BITSEA *Problem total* scores identified seven children (14.6%) with possible deficit or delays (total scores) at the age of 18 months, while the CBCL identified only one child at risk for such problems. The difference between these two questionnaires (the BITSEA and the CBCL) needs more careful examination, with attention to the role of demographic and cross-cultural factors as well as to sensitivity and specificity to early screening of risk factors.

Early detection of social-emotional deficit or delay is very important, particularly when considering parental support and early rehabilitation for a child considered at risk or already manifesting deficits. We noticed in study III, that there were more social-emotional problems reported on the BITSEA in girls than in boys at the age of 36 months; however since our sample of girls was substantially larger than boys ( $n = 32$  and  $n = 16$ , respectively), results should be interpreted with caution.

According to the BITSEA, 10% to 15% of 1- to 2-year-old children in the U. S. (Briggs-Gowan & Carter 2006) and 10–30% (estimate) of children in Southern-Finland (Hakulinen-Viitanen, Pelkonen & Haapakorva, 2005) demonstrate social-emotional or behavioural problems. If the BITSEA is able to identify/screen for infant behaviours suggesting risk for behavioural problems, it will then offer a possibility to discuss (e.g., with parents) a need for intervention or closer follow-up (Briggs-Gowan & Carter, 2006). As mentioned earlier, parental worry may be a primary trigger in the initiation of contact with a provider for children with behaviour problems (Ellingson *et al.* 2004). Because of the mediocre internal consistency of many of some of the BITSEA- subscales in our small sample, further research is needed in larger Finnish samples with the BITSEA to establish the psychometrics of this questionnaire for reliable use in Finland. In general, however, despite the restricted sample size in this preliminary study, the BITSEA evidenced good psychometric properties, similar to other questionnaires that are commonly used (e.g., CBCL) in Finland to children's developmental problems.

Further research with the BITSEA in a larger and broader sample of Finnish children is also needed to determine appropriate cut-off scores. It is important to implement widely-available early screening tools to assess both problems and risk for problems in different cultural and educational settings. In clinical work, ques-

tionnaires, tests and observation results are the typical indicators for intervention or rehabilitation. The addition of early assessments may be particularly valuable in rural areas that may lack the wide range of services available to highly educated individuals or those in more densely populated locations. Ultimately, there is an overall need to further develop multi-method and empirically validated means of evaluating infant developmental stages.

Assessments such as the BITSEA, CBCL and MCDI yield information that cannot be gained in brief clinical encounters. The MCDI evaluates infants' language development and the CBCL screens for behavioural problems, while the BITSEA characterizes infants' social-emotional welfare, development, problems and competence.

Results from this study also demonstrated an association between early communicative development skills and emotional and behavioural problems. Specifically, children with lower communicative skills (based on the MCDI) expressed more withdrawn, externalized and attention problems as well as autistic-like behaviours based on the CBCL and BITSEA than did toddlers with average or high communication abilities. Children with lower communicative skills also demonstrated lower social-emotional competence (based on the BITSEA *Competence* scores) compared to the children with average or high communicative skills, which is important information when planning early intervention and/or rehabilitation. In such cases, group-intervention could be more relevant to enhance social-emotional competence.

Study II and III revealed that higher scores on the BITSEA *Competence* subscale were positively associated with higher scores on the MCDI and fewer behavioural problems on the CBCL subscales at both 18 months and 36 months, which is consistent with previous studies concerning children's communication and behavioural development (Silva, Williams & McGee 1987, Toth *et al.* 2006, Charman *et al.* 2003). The CBCL *Withdrawn* scale was negatively associated with the MCDI; a child's deficient communication skills may impact (or may alternatively be impacted by) decreased social interaction with others, and therefore may result in a decreased opportunity to engage in and/or practice developmentally appropriate communicative skills (e.g., vocabulary, peer interaction via appropriate gestures and imitation). In other words, this relationship may very well be bi-directional or may evidence a negative feedback cycle; future research will investigate this association further.

### **6.1.3 Associations between maternal stress and children's behaviour**

In study IV, we sought to evaluate the association of self-reported maternal stress and parent-reported developmental and behavioural problems in their children over time. The mother-infant dyad is highly interactive (see e.g., Tamminen, 1990, Nurkkala 1998, Markus *et al.* 2000, McCartney *et al.* 2004, Kivijärvi *et al.* 2004); results suggests that higher levels of maternal self-reported stress was associated with an increase in parent-report of problematic child behaviours on the CBCL (*Total problem scores, Internalizing, and Externalizing scales*). Interestingly, this association was limited to the child's 18-month assessment point. In other words, by 36 months, child behaviour was no longer significantly related to maternal stress 8 months post-partum. The fact that early maternal stress and parent-report of problematic behaviours in their children decreased over time, may support an adaptive response of a high stress/high anxious mothers and reciprocally, the notable impact of a change in parenting style on children's behaviour over time. These findings are consistent with prior research studies noting that parenting stress tends to increase parent-report of behavioural problems in their children (e.g., Saisto *et al.* 2008, Yaman *et al.* 2010, Keirn *et al.* 2011).

### **6.1.4 Limitations**

There were notable limitations in this study, which may affect the interpretability of our results. As we have noted several times above, the sample size of this preliminary study was relatively small (N = 50 at 8 months and N = 38 at 36-month follow-up), thus significantly limiting our power overall. The reasons why parents did not participated at later stages might be many, but one could be the fact, that many questionnaires include so many items and needs a lot of time to complete. Although our attrition rate over time is average (69%), it is worth noting that in a family with small children, time to focus on questionnaires may be limited, because a young child needs significant care and attention. The following information characterizes those who did not participate at 18-, 24 and 36 month follow-up; gender and parental education. Eighty percent of participants who were "lost" at follow-up were girls, and almost all parents had higher levels of education. At the age of 18 months, many children in Finland attend day-care and when research-assistants phoned about evaluative questionnaires, many parents stated, that due to time constraints, they were unable to complete questionnaires that

include hundreds of items. Moreover, some families had moved to other towns and were therefore unable to complete the forms at the requested time periods.

There were also limitations in our sample and data collection. For example, only one PSI questionnaire was sent to the family initially, thus we obtained information regarding parental stress from mothers only; therefore, we were unable to examine the role of father's parental stress in our analyses. Further, the video sessions for PC-ERA did not include all of the same individuals at both assessment points, pre- and post- intervention. We were subsequently limited by a small sample of 23 participants who received the same PC-ERA at both assessment points. We used also the MCDI for children aged 36 months (the BITSEA and the CBCL questionnaires were gathered at the same time), although the MCDI manual has standardized this evaluation through the age of 30 months; however, since there is no Finnish parent-reported questionnaires regarding early communication development for children at 36 months, and the range of results at earlier assessment points were quite wide, we selected to use the MCDI for toddlers aged 16 to 30 months in order to see communicative development (vocabulary and comprehension) for our sample of 36-month old children. Moreover, we wanted to maximize the reliability of parent report by utilizing the same measures over time. It is important to note that no children in our sample reached the maximum score on the MCDI scale, thus we were able to rule out any ceiling effect, and further justify the appropriateness of this measure for the age groups in our sample.

In addition, for studies III and IV, we failed to systematically examine demographic variables (e.g. gender differences or parents educational level). Although the samples attrition rate (69%) is within the appropriate range for an 18-month follow-up period, parental worry regarding their child's symptoms and development may have played a role in decreasing the participation over time (e.g., Ellingson *et al.* 2004), and we did not assess parental worry or anxiety over time. The PSI was only administered once, when the child was 8 months old.

Of note, the BITSEA and the CBCL are not validated in Finnish samples, thus precluding us from using culturally appropriate cut-off scores in this study and from assessing clinically relevant psychiatric problems specific to a Finnish population. For example, in our sample toddlers scored one to eight points higher on the CBCL subscales (i.e. *Attention problems, Aggressive behaviour, Externalizing problems and Internalizing problems*) than did toddlers in the United States as reported in the CBCL manual (Achenbach & Rescorla 2000). Our findings in this study, however, are consistent with those of Rescorla and colleagues (2007),

which showed the differences in mean CBCL scores between Finnish samples and reported mean scores in other countries (e.g. Germany, Norway, Iceland, Rescorla *et al.* 2007).

Finally, the reliability of the employed measures varied substantially as indicated in the "Reliability" section above. It will therefore be important for future studies to consider data-driven subscales using inter-item and test construction reliability analyses especially for the BITSEA subscales, *ASD* and *Red Flag*. The distribution of scores obtained in this small sample suggest, that the cut-offs derived from the sample in the United States, may set too high of a threshold for the Finnish population (Haapsamo *et al.* 2009 and 2011).

## **6.2 Conclusions: Implications and future research**

Children with enhanced active vocabulary production and comprehension appear to demonstrate better social-emotional competence than do children with poorer vocabulary production and comprehension; this suggests the importance of interaction and social relatedness in language acquisition. Furthermore, behavioural problems may be more common in toddlers with deviant communication development (Silva, Williams & McGee, 1987, Irwin, Carter & Briggs-Gowan 2002, Rescorla, Ross & McClure, 2007a), which underlines the importance of early rehabilitation of communication skills and behavioural guidance.

Children who do not manifest social-emotional competence may benefit from guided play with peers and tasks with joint and shared attention (e.g., Tomasello, 1988, 1995, Lyytinen 1999b) to possibly prevent problems such as ASD features (e.g., Gray & Tonge 2001, Landa & Garret-Mayer 2006). Therefore, family-based enrichment-intervention focusing on play and music may offer one method for early prevention. It is also important to support the parent-child relationship in a preventive manner, due to the possible reciprocal link between maternal representations and child's maladaptive behaviours (Dollberg, Feldman & Keren 2010); behavioural problems in children may result in increased parental worry, stress and anxiety, and if the child's active vocabulary is limited, parents might benefit from early intervention and guidance in order to better understand their child and his/her behaviour and serve as an role model in producing more prosocial communication and behaviours.

Early assessment and identification of maladaptive mother-child-interaction facilitate early intervention and adaptive restructuring of such interaction patterns. Therefore, in the future, we plan to examine the interactional styles (from collect-

ed PC-ERA videotapes) between all parents and children during the enrichment intervention in our study, as parental feedback was quite positive despite the modest results of the parent-child observations of the intervention group.

The children's healthcare system in Finland is vigilant and capable of providing early assessment and thus early intervention for disrupted early developing parent-child-relationships. For example in Northern Finland, in the city of Oulu, almost all children's healthcare nurses are educated to evaluate early interaction between a mother and her infant (Hakulinen-Viitanen, Pelkonen & Haapakorva 2005). Hence, it is possible to initiate preventive actions with families considered at risk. Once the BITSEA (and its subscales) becomes validated in larger and broader samples of Finnish children, this assessment may prove quite helpful in the early identification of possible developmental risks (e.g., ASD).

A final important implication from this study is, that there are no parent-report questionnaires for children's communicative skills with clear cut-offs and for older children; such assessments are invaluable since young children's interaction and communication skills are most observable in the home environment (Lyytinen 1999a).

## References

- Abidin RR (1999) Parenting Stress Index. Third Edition. Professional Manual. Psychological Assessment Resources, Inc. Pediatric Psychology Press, Charlottesville, VA, 1990.
- Achenbach TM & Rescorla LA (2000) Manual for the ASEBA preschool forms and profiles. Burlington, VT; University of Vermont, Research Centre for Children, Youth and Families
- Achenbach T (2001) Child Behaviour Checklist/1½-5. University of Vermont Department of Child Psychiatry, Burlington.
- Ahlqvist S (2003) Varhainen vanhempi-lapsi-suhteen arviointimenetelmä. Manual. Turku, Finland: University of Turku, Department of Psychology.
- Ayoub C, Vallotton CD, Masterson AM (2011) Developmental pathways to integrated social skills: the roles of parenting and early intervention. *Child Dev* 82(2): 583–600.
- Baird J, Hill CM, Kendrick T, Inskip HM & SWS Study Group (2009) Infant sleep disturbance is associated with preconceptional psychological distress: findings from the Southampton Women's Survey. *Sleep* 32(4): 566–8.
- Bates JE, Freeland CA & Lounsbury ML (1979) Measurement of infant difficultness. *Child Dev* 50: 794–803.
- Bayer JK, Hiscock H, Ukoumunne OC, Scalzo K, Wake M (2010) Three-year-old outcomes of a brief universal parenting intervention to prevent behaviour problems: randomised controlled trial. *Arc Dis Child* 95(3): 187–92.
- Bernabei P, Fenton G., Fabrizi A, Camaioni L & Perucchini P (2003) Profiles of sensorimotor development in children with autism and with developmental delay. *Perception Motor Skills* 96 (3 Pt2): 1107–16.
- Best W, Melvin D & Williams S (1993) The effectiveness of communication groups in day nurseries. *J Eur Disorders Comm* 28(2): 187–212.
- Bloom L & Tinker E (2001) The intentionality Model and Language Acquisition. Overton, W.F. (ed.) *Mon Soc Res Child Dev* 267(66): 4.
- Bricker D, Squires J, Kaminski R & Mounts L (1988) The validity, reliability and cost of a parent-completed questionnaire system to evaluate at-risk infants. *J of Pediatrics Psychol* 13(1): 55–68.
- Briggs-Gowan M, Carter A Skuban E & Horwitz S (2001) Prevalence of social-emotional and behaviour problems in a community sample of 1- and 2-year-old children. *J Am Acad Child Adol Psychiatry* 40: 811–819.
- Briggs-Gowan MJ, Carter AS, Irwin JR, Wachtel K & Cicchetti DV (2004) The Brief Infant-Toddler Social and Emotional Assessment: Screening for Social-Emotional Problems and Delays in Competence. *J of Pediatric Psychol* 29(2): 143–155.
- Briggs-Gowan MJ & Carter AS (2006) BITSEA, Brief Infant-Toddler Social Emotional Assessment. Examiner's Manual. Boston: Yale University and the University of Massachusetts.
- Briggs-Gowan MJ & Carter AS (2008) Social-emotional screening status in early childhood predicts elementary school outcomes. *Pediatrics* 121(5): 957–962.

- Broussard ER & Hartner MSS (1971) Further considerations regarding maternal perception of the first born. In: *Exceptional infant 2. studies in abnormality*. Hellmuth J, Butterworths (Ed.), London: 432–449.
- Browne JV & Talmi A (2005) Family-based intervention to enhance infant-parent relationships in the neonatal intensive care unit. *J Pediatric Psychol* 30(8):667–677.
- Buschmann A, Jooss B, Rupp A, Dockter S, Blaschikowitz H, Heggen I & Pietz J (2008) Children with developmental language delay at 24 months of age: results of a diagnostic work-up. *Dev Med & Child Neurol* 50: 223–229.
- Buschmann A, Jooss B, Rupp A, Feldhusen F, Pietz J & Phillippi H (2009) Parent based language intervention for 2-year-old children with specific expressive language delay: a randomized controlled trial. *Arch Dis Child* 94(2): 110–6.
- Carey WB & McDevitt SC (1978) Revision on the infant temperament questionnaire. *Pediatrics*, 61: 735–739.
- Carpenter M, Nagell K & Tomasello M (1998) Social cognition, joint attention, and communicative competence from 9 to 15 months of age. *Mon Soc Res Child Dev*, 255 (63): 4: 1–143.
- Carter AS, Garrity-Rokous FE, Chazan-Cohen R, Little C, & Briggs-Gowan MJ (2001) Maternal depression and co morbidity: Predicting early parenting, attachment security and toddler social-emotional problems and competencies. *J Am Acad Child Adol Psychiatry* 40(1): 18–26.
- CBCL 1½-5-vuotiaille (7-28-00 Edition) Thomas M. Achenbach 2000. Almqvist 2001 (Finnish version).
- Charman T, Drew A, Baird C & Baird G. (2003) Measuring early language development in preschool children with autism spectrum disorder using the MacArthur Communicative Development Inventory (Infant Form). *J Child Lang* 30(1): 213–36.
- Cheng S, Kondo N, Aoki Y, Kitamura Y, Takeda Y, Yamagata Z (2007) The effectiveness of early intervention and the factors related to child behavioural problems at age 2: a randomized controlled trial. *Early Hum Dev* 83(10):683–91.
- Clark R (1985) *The Parent-Child Early Relational Assessment*. Manual. Wisconsin: Department of Psychiatry University of Wisconsin Medical School.
- Cox JL, Holden JM & Sagovsky R (1987) Detection of postnatal depression, development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatr* 150: 782–786.
- Crittenden PM (1998) *CARE-Index*. Coding manual. Family Relations Institute, Miami, FL.
- Davison M & Howlin P (1997) A follow-up study of children who had attended specialist primary schools for language delay. *Eur J Disorders Comm* 32(1): 19–36.
- Delgado CEF & Delgado I (2002) Infant communication development and maternal interpretation of intentionality. Poster presentation at the International Conference on Infant Studies at Toronto, Ontario Canada.
- Dollberg D, Feldman R & Keren M (2010) Maternal representations, infant psychiatric status, and mother-child relationship in clinic-referred and non-referred infants. *Eur J Child Adol Psychiatry*, 19: 25–36.



- Durkin K & Conti-Ramsden G (2007) Language, Social behaviour, and the quality of friendships in adolescents with and without a history of specific language impairment. *Child Dev* 78(5): 1441–1457.
- Edwards S, Fletcher P, Garman A, Hughes A, Letts C & Sinka I (1997) Reynell Developmental Language Scales III, The University of Reading edition. NferNelson, Berkshire, UK. Translation and standardization of the Finnish version: *Psykologien kustannus Oy* 2001.
- Elkind D & Weiner IB (1978) *Development of the Child*. Library of Congress Cataloging in Publication Data. John Wiley & Sons, Inc.
- Ellingson KD, Briggs-Gowan MJ, Carter AS & Horwitz SM (2004) Parent Identification of Early Emerging Child Behaviour Problems. Predictors of Sharing Parental Concern With Health Providers. *Archiv Ped Adol Med* 158: 766–772.
- EU-Programs, Publications 1/00, Programs for Northern-Finland 1, 2000 -2006. [www.intermin.fi](http://www.intermin.fi). Source cited at 8.10.2011.
- Feldman R, Granat A, Pariente C, Kanety H, Kuint J & Gilboa-Schechtman E (2009) Maternal depression and anxiety across the postpartum year and infant social engagement, fear regulation and stress reactivity. *J Am Acad Child Adol Psychiatry*, 48(9): 919–27.
- Fenson L, Dale PS, Reznick JS, Thal D, Bates E, Hartung JP, Pethick S & Reilly JS (1994) *Technical Manual for the MacArthur Communicative Development Inventories*, San Diego State University. San Diego, CA, Singular Press.
- Fowles E & Walker L (2009) Maternal predictors of toddler health status. *J Sp Ped Nurs*, 14(1): 33–40.
- Freud WE (1989) Psychological aspects of neonatal intensive care. In *The Course of Life*. Vol. 1 Infancy. Ed. by S. I. Greenspan and G. H. Pollock. International Universities press, Madison, Connecticut, USA.
- Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O & Rutter C (1997) The validity of two version of the GHQ in the WHO study of mental illness in general health care. *Psychol Med* 27: 191–197.
- Gray KM & Tonge BJ (2001) Review article: Are there early features of autism in infants and preschool children? *J Ped Child Health* 37: 221–226.
- Guiberson M, Rodriquez BL, & Dale PS (2011) Classification accuracy of brief parent report measures of language development in Spanish-speaking toddlers. *Language, Speech and Hearing Services in Schools* 42(4): 536–549.
- Guralnick MJ, Neville B, Hammond MA & Connor RT (2008) Mothers' Social Communicative Adjustments to Young Children With Mild Developmental Delays. *Am J Ment Retard* 113(1): 1–18.
- Haapsamo H, Ebeling H, Soini H., Joskitt L, Larinen K, Penninkilampi-Kerola V, Carter A & Moilanen I (2009) Screening infants with social and emotional problems; A pilot study of the Brief Infant Toddler Social and Emotional Assessment (BITSEA) in Northern Finland. *Int J Circumpolar Health* 68(4): 386–393.

- Haapsamo H, Kuusikko-Gauffin S, Carter A, Pollock-Wurman R, Ebeling H, Soini H, Joskitt L, Pihlaja P & Moilanen I (2011) A Pilot longitudinal follow-up study of the Brief Infant Toddler Social Emotional Assessment (BITSEA): Examining toddler's social-emotional, behavioural and communicative development. *Early Child, Dev Care.* iFirst Article 21.10.2011, 1–16. URI: <http://www.tandfonline.com/doi/abs/10.1080/03004430.2011.622756>
- Haapsamo H, Kuusikko-Gauffin S, Ebeling H, Larinen K, Penninkilampi-Kerola V, Soini H & Moilanen I (2012) Communication development and characteristics of influencing factors: a follow-up study from 8 to 36 months. *Early Child Dev Care.* iFirst article DOI:10.1080/03004430.2012.674523
- Hakulinen-Viitanen T & Pelkonen M & Haapakorva A (2005) Äitiys- ja lastenneuvolatyö Suomessa. Sosiaali- ja terveysministeriön selvityksiä 2005:22.
- Harel J, Eshel Y, Ganor O, & Scher A (2002) Antecedents of mirror self-recognition of toddlers: emotional availability, birth order and gender. *J Inf Ment Health*, 2: 293–309.
- Hirshfeld-Becker, D. R., Biederman, J., Faraone, S. V., Robin, J. A., Friedman, D., Rosenthal, J. M., *et al.*, (2004). Pregnancy complications associated with childhood anxiety disorders. *Depression and Anxiety* 19(3): 152–62.
- Howlin P (1984) Parents as therapists: A review. In Miller D (ed.) *Remediating Children's Language*. London: Croom Helm.
- Hughes CA, Thomas KA, Iatrow AM & Bozynski ME (1998) The value of the Infant/Child Monitoring Questionnaire for identification of developmental disability among neonatal intensive care unit graduate. *J Perinatology* 18(4): 259–65.
- Hutterlocher J, Waterfall H, Vasilyeva M, Vevea J & Hedges LV (2010) Sources of variability in children's language growth. *Cogn Psychol* 6(4): 343–65.
- Iivonen A (1994a) Lapsen varhainen äänteellinen kausi. *Suomen logopedis-foniatrinen aikakauslehti* 1/94: 5–19
- Iivonen A (1994b) Paradigmaattisia ja syntagmaattisia näkökohtia lapsen foneettis-fonologisessa kehityksessä. Teoksessa *Lapsen normaali ja poikkeava kielen kehitys*, toinen painos. Iivonen A & Lieko A & Korpilahti P (ed.), Helsinki: SKS: 34–77.
- Iivonen A (1995) Lapsen fonologis-foneettinen kehitys: 3. Lapsi leksikon kynnyksellä. *Suomen logopedis-foniatrinen aikakauslehti* 15(1): 1–15.
- Irwin JR, Carter AS & Briggs-Gowan MJ (2002) The social-emotional development of "late-talking" toddlers. *J Am Acad Child Adol Psychiatry* 41(11): 1324–1332.
- Jakkula K (2002) Esineiden antaminen. Kehityksen peili ja kieltä ennakoiva sosiaalinen merkki 9–34 kuukauden iässä. Doctoral thesis. Kasvatustieteiden tiedekunta, Oulun yliopisto *Acta Universitatis Ouluensis Scientiae Rerum Socialium* E 52.
- Kagan J (1997) *Galen's Prophecy; Temperament in Human Nature*. Westview Press.
- Kaiz M, Maytal HR, Devor R, Bergman L & Mankuta D (2010) Maternal anxiety, mother-infant interactions, and infants' response to challenge. *Inf Behav Dev* 33(2): 136–48.
- Kanner L (1935) *Child Psychiatry*. Third Edition. Springfield, Illinois: Charles C Thomas.

- Keirn S, Daniels J, Dole N, Herring A, Siega-Riz A & Scheidt P (2011) A prospective study of maternal anxiety, perceived stress, and depressive symptoms in relation to infant cognitive development. *J Early Hum Dev* 87(5): 373–380.
- Kemppinen K, Kumpulainen K, Räsänen E, Moilanen I, Ebeling H, Hiltunen P, & Kuneilius A (2005) Mother-child interaction on video compared with infant observation: Is five minutes enough time for assessment? *J Infant Ment Health* 26: 69–81.
- Kemppinen K, Ebeling H, Raita-Hasu J, Toivonen-Falck A, Paavola L, Moilanen I, Kumpulainen K (2007) Early maternal sensitivity and child behaviour at toddler age: Does low maternal sensitivity hinder identification of behavioural problems? *J Reproductive Inf Psychol* 25(4): 270–284
- Kivijärvi M, Voeten MJM, Niemelä P, Räihä H, Lertola K & Piha J (2001) Maternal sensitivity behaviour and infant behaviour in early interaction. *J Infant Ment Health* 22(6): 627–640.
- Kivijärvi M, Räihä H, Virtanen S, Lertola K & Piha J (2004) Maternal sensitivity behaviour and infant crying, fussing and contented behaviour: The effects of mother's experienced social support. *Scand J Psychol* 45:239–246.
- Kivijärvi M, Räihä H, Virtanen S, Lertola K & Piha J (2005) Infant temperament and maternal sensitivity behaviour in the first year of life. *Scand J Psychol* 46(5): 421–8.
- Kokkonen P, Veijola J, Karvonen JT, Läksy K, Jokelainen J, Järvelin MR & Joukamaa M (2003) Ability to speak at the age of 1 year and alexithymia 30 years later. *J Psychosom Res* 54(5): 491–5.
- Korja R, Maunu J, Kirjavainen J, Savonlahti E, Haataja L, Lapinleimu H, Manninen H, Piha J, Lehtonen L & the PIPARI Study group (2007) Mother-infant interaction is influenced by the amount of holding in preterm infants. *Early Hum Dev*, 84: 257–267.
- Kunnari S (1997) Fonologisen kehityksen varhaisvaiheet. *Suomen logopedis-foniatrinen aikakauslehti* 17: 1–2: 33–38.
- Laakso M-L, Poikkeus A-M, Katajamäki J & Lyytinen P (1999) Early intentional communication as a predictor of language development in young toddlers. *First Language* 19: 207–231.
- Laalo K (1998) Esi- ja varhaismorfologinen vaihe suomalaislapsen kielen kehityksessä. In Heinänen K & Lehtihalmes M (Eds.), *Kielihäiriöisen lapsen tutkiminen - testaamista-ko?* (Vol. 30: 22–30). Helsinki: Suomen logopedis-foniatrinen yhdistys ry:n julkaisuja.
- Landa R & Garrett-Mayer E (2006) Development in infants with autism spectrum disorders: A prospective study. *J Child Psychol Psychiatry* 47(6): 629–638.
- Landry SH, Smith KE, Swank PR, Guttentag C (2008) A responsive parenting intervention: the optimal timing across early childhood for impacting maternal behaviours and child outcomes. *Dev Psychol* 44(5): 1335–53
- Law J Garret Z & Nye C (2010) Speech and language therapy interventions for children with primary speech and language delay or disorder. *Cochrane Database of Systematic Reviews*. CD004110. DOI:10.1002/14651858.CD004110.
- Lowell DI, Carter AS, Godoy L, Paulicin B, Briggs-Gowan MJ (2011) A randomized controlled trial of Child FIRST: a comprehensive home-based intervention translating research into early childhood practice. *Child Dev* 82(1): 193–208.

- Lung FW, Shu BC, Chiang TL, Lin SJ (2010) Maternal mental health and childrearing context in the development of children at 6, 18 and 36 months: a Taiwan birth cohort pilot study. *Child Care Health Dev* 37(2): 211–213.
- Lyytinen P (1999a) Varhaisen kommunikaation ja kielen kehityksen arviointimenetelmä. Niilo Mäki Instituutti, Jyväskylän yliopiston Lapsitutkimuskeskus. Yliopistopaino, Jyväskylä.
- Lyytinen P, Laakso M-L, Poikkeus AM & Rita N (1999b) The development and predictive relations of play and language across the second year. *Scand J Psychol* 40: 3: 177–186.
- Markus J, Mundy P, Morales M, Delgado CEF & Yale M (2000) Individual Differences in Infant Skills as Predictors of Child-Caregiver Joint Attention and Language. *Soc Dev* 9(3): 302–315.
- Mauridsen SE & Hauschild KM (2010) The sex ratio of siblings of individuals with a history of developmental language disorder. *Log Phon Voc* 35(3): 144–8.
- McCartney K, Tresch Owen M, Booth CL, Clarke-Stewart A & Lowe Wandell D (2004) Testing a maternal attachment model of behaviour problems in early childhood. *J Child Psychol Psychiatry* 45(4): 765–778.
- Mendelsohn AL, Valdez PT, Flynn V, Foley GM, Berkule SB, Tomopoulos S, Fierman AH, Tineo W & Dreyer BP (2007) Use of videotaped interactions during pediatric well-child care: impact at 33 months on parenting and on child development. *J Dev Behav Pediatr* 28(3): 206–12.
- Meyer EC, Coll CT, Lester BM, Boukydis CF, McDonough SM & Oh W (1994) Family-based intervention improves maternal psychological wellbeing and feeding interaction of preterm infants. *Pediatrics* 93(2): 241–6.
- Mitchell S, Brian J, Zwaigenbaum L, Roberts W, Szatmari P, Smith I & Bryson S (2006) Early language and communication development of infants later diagnosed with autism spectrum disorder. *J Dev Behav Ped* 27(2suppl): 69–78.
- Morales M, Mundy P, Fullmer C, Yale M, Messinger D, Neal R, & Schwartz H (2000) Responding to joint attention across the 6- to 24-month age period and early language acquisition. *J Appl Dev Psychol* 21 (3): 283–298.
- Mothander, PR & Moe RG (2010) Self-reported depressive symptoms and parental stress in mothers and fathers who bring their infants to an infant mental health clinic. *Nord J Psychiatry* 64(5): 310–6
- Mäntymaa M, Puura K, Luoma I, Salmelin R K & Tamminen T (2004) Early mother-infant interaction, parental mental health and symptoms of behavioural and emotional problems in toddlers. *J Inf Behav Dev* 27: 134–149.
- Mäntymaa M, Puura K, Luoma I, Salmelin R & Tamminen T (2006) Mother's early perception of her infant's difficult temperament, parenting stress and early mother-infant interaction. *Nord J Psychiatr* 60(5): 379–86.
- Mäntymaa M, Puura K, Luoma I, Latva R, Salmelin RK & Tamminen T (2011) Predicting internalizing and externalizing problems at five years by child and parental factors in infancy and toddlerhood. *Child Psychiatry Hum Dev*, Sep 29. Epub ahead of print.

- Nurkkala K (1998) Action rituals between mother and infant. In Heinänen K & Lehtihalmes M (eds.). Proceedings of the 7th Nordic Child Language Symposium. Oulu: University of Oulu. (pp. 32–37).
- Official Statistics of Finland (OSF) (2011): Educational structure of population [e-publication]. Helsinki: Statistics Finland [referred: 25.9.2011]. URI: [http://www.stat.fi/til/vkour/index\\_en.html](http://www.stat.fi/til/vkour/index_en.html).
- Official Statistics of Finland (OSF): Births [e-publication]. ISSN=1798-2413. 2010. Helsinki: Statistics Finland [referred: 16.1.2012]. URI: [http://www.stat.fi/til/synt/2010/synt\\_2010\\_2011-04-15\\_tie\\_001\\_en.html](http://www.stat.fi/til/synt/2010/synt_2010_2011-04-15_tie_001_en.html).
- Olds D, Sadler L & Kitzman H (2007) Programs for parents of infants and toddlers: recent evidence from randomized trial. *J Child Psychol Psych* 48 (3/4): 355–391.
- Olson SL, Bates JE, Sandy JM & Lanthier R (2000) Early developmental precursors of externalizing behaviour in middle childhood and adolescence. *J Abnorm Child Psychol* 28(2): 119–33.
- Paavola L (2006a) Maternal sensitive responsiveness, characteristics and relations to child early communicative and linguistic development. Doctoral thesis, University of Oulu. Faculty of Humanities, Department of Finnish, Information Studies and Logopedics, Department of Paediatrics. *Acta Universitatis Ouluensis B* 73.
- Paavola L, Kemppinen K, Kumpulainen K, Moilanen I and Ebeling H (2006b) Maternal sensitivity, infant co-operation and early linguistic development: Some predictive relations. *Eur J Dev Psychol* 3(1): 13–30.
- Pajulo M, Savonlahti E, Sourander A, Ahlqvist S, Helenius H & Piha J (2001) An early report on the mother-baby interactive capacity of substance-abusing mothers. *J Subst Abuse Treat* 20(2): 143–51.
- Parenting Stress Index-Short form (PSI, Abidin, 1999) Psychology Press, Taylor & Francis Group.
- Parre M (1994) Lapsen kielenkehityksen komponenteista, häiriöistä ja diagnosoinnista. In *Lapsen normaali ja poikkeava kielen kehitys*, second ed. Iivonen, A. & Lieko, A. & Korpilähti, P. (ed.), Helsinki: SKS: 206–224.
- Piaget J (1977) & Inhelder B (ed.) *Lapsen psykologia*. (Original book "La psychologie de l'enfant" (1966) , translated in Finnish by Mirja Rutanen). K.J. Gummerus, Jyväskylä.
- Pihlakoski I, Sourander A, Aromaa M, Rautava P, Helenius H & Sillanpää M (2006) The continuity of psychopathology from early childhood to preadolescence. A prospective cohort study of 3–12 –year-old children. *Eur Child Adol Psychiatry* 15(7): 409–17.
- Plomin R & Dale PS (2000) Genetics and early language development: A UK study of twins. In *Speech and Language Impairments in Children. Causes, Characteristics, Intervention and Outcome*. Bishop, D.V. M. and Leonard, L. B. (Eds.) Psychology Press. Taylor and Francis Group: 35–51.
- Prior M, Bavin E L, Cini E, Reilly S, Bretherton L, Wake M & Eadie P (2008) Influence on communicative development at 24 months of age: Child temperament, behaviour problems, and maternal factors. *Inf Behav Dev* 31(2): 270–9.

- Puolakanaho A, Ahonen T, Aro M, Eklund K, Leppänen PHT, Poikkeus A-M, Tolvanen A, Torppa M. & Lyytinen H (2007) Very early phonological and language skills: estimating individual risk of reading disability. *J Child Psychol Psych* 48(9): 923–931.
- Radecki L, Sand-Loud N, O'Connor KG, Sharp S, Olson LM (2011) Trends in the use of standardized tools for developmental screening in early childhood: 2002–2009. *Pediatrics* 128(1): 14–9.
- Reilly S, Eadie P, Bavin EL, Wake M, Prior M, Williams H, Bretherton L, Barrett Y & Ukoumunne OC (2006) Growths of infant communication between 8 and 12 months: A population study. *J Paed Child Health* 42: 764–770.
- Reilly S, Wake M, Bavin E L, Prior M, Williams J, Bretherton L, Eadie P, Barrett Y & Ukoumunne OC (2007) Predicting Language at 2 years of Age: A Prospective Community Study. *Pediatrics* 120: 6.
- Rescorla L, Ross GS, McClure S (2007a) Language delay and behavioural/emotional problems in toddlers: findings from two developmental clinics. *J Speech Lang Hear Res* 50(4): 1063–78
- Rescorla L, Achenbach TM, Ivanova MY, Dumenci L, Almqvist F, Bilenberg N, Bird H, Broberg A, Dobrean A, Döpfner M, Erol N, Forns M, Hannesdottir H, Kanbayashi Y, Lambert MC, Leung P, Minaei A, Mulatu MS, Novik TS, Oh KJ, Roussos A, Sawyer M, Simsek Z, Steinhausen HC, Weintraub S, Metzke CW, Wolanczyk T, Zilber N, Zukauskienė R and Verhulst F (2007b) Epidemiological comparisons of problems and positive qualities reported by adolescents in 24 countries. *J Consult Clin Psychology* 75(2): 351–8.
- Rossetti LM (1996) Communication intervention. Birth to three. Singular publishing Group, Inc. London, UK.
- Rubin KH, Burgess KB, Dwyer KM & Hastings PD (2003) Predicting preschoolers' externalizing behaviours from toddler temperament, conflict, and maternal negativity. *Dev Psychol* 39(1): 164–76.
- Saisto T, Salmela-Aro K, Nurmi J E & Halmesmäki E (2008) Longitudinal study on the predictors of parental stress in mothers and fathers in toddlers. *J Psychosom Obs Gyn* 29(3): 213–22.
- Savonlahti E, Pajulo M, Ahlqvist S, Helenius H, Korvenranta H, Tamminen T & Piha J (2005) Interactive skills of infants with their high-risk mothers. *Nord J Psychiatry* 59(2): 139–47.
- Schieve LA, Blumberg SJ, Rice C, Visser SN & Boyle C (2008) The Relationship Between Autism and Parenting Stress. *J Am Acad Pediatrics* 119: S1114-S121
- Silk JS, Shaw DS, Skuban EM, Oland AA & Kovacs M (2006) Emotion regulation strategies in offspring of childhood-onset depressed mothers. *J Child Psychol Psych* 47(1): 69–78.
- Silva PA, Williams S & McGee R (1987) A longitudinal study of children with developmental language delay at age three: later intelligence, reading and behaviour problems. *Dev Med Child Neurol* 29(5): 630–40.

- Sinkkonen J (1995) Lapsen varhaiset vuorovaikutussuhteet, kommunikaatio ja kielen kehitys. *Suomen logopedis-foniatriinen aikakauslehti*, 15(3–4): 65–70.
- Skovgaard AM Olsen EM, Christiansen E, Houmann T, Landorph SL, Jorgensen T & CCC 2000 Study group (2008) Predictors (0–10 months) of psychopathology at age 1½ years – a general population study in the Copenhagen Child Cohort CCC 2000. *J Child Psychol Psych* 49(5): 553–62.
- Smith T, Groen AD & Wynn JW (2000) Randomized Trial of Intensive Early Intervention for Children With Pervasive Developmental Disorder. *Am J Ment Retard* 105(4): 269–285.
- Smith LB, Jones SS, Landau B, Gershkoff-Stowe, L & Samuelson L (2002) Object name learning provides on-the-job training for attention. *Psychol Sci* 13(1): 13–19.
- Sparks TA, Hunter SK, Backman T L, Morgan GA & Ross RG (2011) Maternal parenting stress and mothers' reports of their infants' mastery motivation. *Infant Behav Development* Sep 2. Epub ahead of print.
- Sperlich S, Arnhold-Kerri S. & Geyer S (2011) What accounts for depressive symptoms among mothers? The impact of socioeconomic status, family structure and psychosocial stress. *Int J Pub Health* 56(4): 385–396.
- Squires J, Bricker D & Potter L (1997) Revision of a Parent-Completed Developmental Screening Tool: Ages and Stages Questionnaires. *J Pediatric Psychol* 2(3):313–328.
- Stakes, Indikaattoripankki SOTKANet 2005–2008. <http://www.stakes.fi> Tarkistettu 25.9.2011.
- State Provincial Office Publications, services in Southern-Finland 2004, web-pages: [www.laanhallitus.fi](http://www.laanhallitus.fi). Peruspalvelut Etelä-Suomen läänissä 2003, Etelä-Suomen lääninhallituksen julkaisuja 78, Hakapaino, Helsinki 2004. Source cited at 3.2.2009.
- Stein E, Malmberg LE, Sylva K, Barnes J, & Leach P & FCCC team (2008) The influence of maternal depression, caregiving and socioeconomic status in the post-natal year on children's language development. *Child, Care Health Dev* 34(5): 603–12.
- Stern DN (1985) *The Interpersonal World of the Infant. A View from Psychoanalysis and Developmental Psychology*. Basic Books, Inc. Publishers, New York.
- Stolt S, Klippi A, Launonen K, Munck P, Lehtonen L, Lapinleimu H, Haataja L & Pipari study group (2007) Size and composition of the lexicon in prematurely born very-low-birth-weight and full-term Finnish children at two years of age. *J Child Lang* 34(2): 283–310.
- Suomalaisen Lääkäriseuran Duodecimin ja Suomen Fysiatryhdistyksen asettama työryhmä. *Kielellinen erityisvaikeus, Käypä Hoito-suositus* (2010) Helsinki: Suomalainen Lääkäriseura Duodecim, viitattu 9.10.2011, URI: [www.kaypahoito.fi](http://www.kaypahoito.fi)
- Tamminen T (1990) Äidin masennus, imetys ja varhainen vuorovaikutus. Doctoral thesis. *Acta Universitatis Tamperensis*. Vammalan Kirjapaino Oy.
- Tervo R (2007) Language proficiency, development and behavioural difficulties in toddlers. *J Clinic Pediatrics* 46(6): 530–9.
- Toivainen J (1994a) Mistä lapsi aloittaa kielensä opettelun? Lingvistisiä lähtökohtia. *Suomen logopedis-foniatriinen aikakauslehti* 14(4): 72–80.

- Toivainen J (1994b) Lasten erilaisuus suomen kielen morfologian omaksunnassa. In Iivonen A & Lieko A & Korpilahti P (Eds.), *Lapsen normaali ja poikkeava kielen kehitys*, second ed., Helsinki: SKS:268–304.
- Tolonen K (1996) Lapsen esikielellisen kauden verbaalisen kehityksen taustaa. Tutkimus 0–18 kuukauden ikäisen lapsen orientaatiosta, jäljittelystä, vuorottelusta, kiintymyssuhteesta sekä merkityksen muodostamisesta toiminnan kontekstissa. Oulun yliopiston opettajankoulutuslaitoksen julkaisuja. Sarja C: Varhaiskasvatus 1/1996. Kajaanin opettajankoulutuslaitos, Kajaani.
- Tomasello M (1988) The role of joint attention in early language development. *Language Sci* 11: 69–88.
- Tomasello, M. (1995) Joint attention and social cognition. In C. Moore & P. Dunham (ed.) *Joint attention: It's origins and role in development*. Hillsdale, NJ: Lawrence Erlbaum Associates: 103–131.
- Tomasello M (2000) Acquiring syntax is not what you think. In *Speech and Language Impairments in Children. Causes, Characteristics, Intervention and Outcome*. Bishop, D.V. M. and Leonard, L. B. (Eds.) Psychology Press. Taylor and Francis Group: 1–15.
- Tonge B, Brereton A, Kiomall M, MacKinnon A, King N & Rinehart N (2006) Effects on Parental Mental Health of an Education and Skills Training Program for Parents of Young Children With Autism: A Randomized Controlled Trial. *J Am Acad Child Adol Psych* 45(5): 561–569.
- Toth K, Munson J, Meltzoff AN & Dawson G (2006) Early Predictors of Communication Development in Young Children with Autism Spectrum Disorder: Joint Attention, Imitation, and Toy Play. *J Aut Dev Disorders* 36: 993–1005.
- Trillingsgaard A, Ulsted Sorensen E, Nemeč G & Jorgensen M (2005) What distinguishes autism spectrum disorders from other developmental disorders before the age of four years? *Eur Child Adol Psych* 14(2): 65–72.
- Viholainen H, Ahonen T, Cantell M, Lyytinen P & Lyytinen H (2002) Development of early motor skills and language in children at risk for familial dyslexia. *J Dev Med Child Neurol* 44(11): 761–9.
- Wake M, Tobin S, Girolametto L, Ukoumunne OC, Gold L, Levickis P, Sheehan J, Goldfield S & Reilly S (2011) Outcomes of population based language promotion for slow to talk toddlers at ages 2 and 3 years: Let's Learn Language cluster randomised controlled trial. *BMJ* 343: d4741
- Wetherby AM & Pritzand BM (1999) Enhancing Language and Communication Development in Autism: Assessment and Intervention Guidelines. In Berkell Zager, D. (ed.) *Autism. Identification, Education and Treatment*. second ed. Mahwah, USA: Lawrence Erlbaum Associates Publishers: 141–174.
- Wing JK, Cooper JE & Sartorius N (1974) *The measurement and classification of psychiatric symptoms*. Cambridge University Press, London.
- Wissow, LS, Gadomski A, Roter D, Larson S, Brown J, Zachary C, Bartlett E, Horn I, Luo X & Wang MC (2008) Improving child and parent mental health in primary care: a cluster-randomized trial of communication skills training. *Pediatrics* 121(2): 266–75.



- Yamada A, Suzuki M, Kato M, Suzuki M, Tanaka S, Shindo T, Taketani K, Akechi T & Furukawa TA (2007) Emotional distress and its correlates among parents of children with pervasive developmental disorders. *J Psychiatry Clin Neurosc* 61: 651–657.
- Yaman A, Mesman J, van Ijzendoorn MH, Bakermans-Kranenburg MJ (2010) Perceived family stress, parenting efficacy, and child externalizing behaviours in second-generation immigrant mothers. *Soc Psychiatry Psychiatr Epidemiol* 45(4): 505–12.
- Överlund J (1994) Vauvan kommunikaatiotaitojen kehitys - puheen oppimisen perusta. In Iivonen A & Lieko A & Korpilahti P (Eds.), *Lapsen normaali ja poikkeava kielen kehitys*, second ed., Helsinki: SKS: 306–331.
- Överlund J (1996) Puhe ja kieli kehittyvät vuorovaikutuksessa. In Launonen K & Korpijaakko-Huuhka A-M (Eds.), *Kommunikoinnin häiriöt; syitä, ilmenemismuotoja ja kuntoutuksen perusteita*, second ed., Helsingin yliopisto, Lahden tutkimus- ja koulutuskampus: 20–38.



## Original Publications

This thesis is based on the following publications which are referred to in the text by their Roman numerals. In addition, this thesis includes some unpublished data.

- I Haapsamo H, Kuusikko-Gauffin S, Ebeling H, Larinen K, Penninkilampi-Kerola V, Soini H & Moilanen I (2012) Communication development and characteristics of influencing factors; A follow-up study from 8 to 36 months. *Early Child Dev Care*, iFirst Article 23.4.2012. <http://www.tandfonline.com/doi/abs/10.1080/03004430.2012.674523>
- II Haapsamo H, Ebeling H, Soini H, Joskitt L, Larinen K, Penninkilampi-Kerola V, Carter AS & Moilanen I (2009) Screening infants with social and emotional problems; A pilot study of the Brief Infant Toddler Social and Emotional Assessment (BITSEA) in the Northern Finland. *Int J of Circumpolar Health* 68(4): 386–393
- III Haapsamo H, Kuusikko-Gauffin S, Carter A, Pollock-Wurman R, Ebeling H, Joskitt L, Larinen K, Soini H, Pihlaja P & Moilanen I (2011). A Pilot longitudinal follow-up study of the Brief Infant Toddler Social Emotional Assessment (BITSEA): Examining toddler's social-emotional, behavioural and communicative development. *Early Child Dev Care* 182(11): 1487–1502. <http://www.tandfonline.com/doi/abs/10.1080/03004430.2011.622756>
- IV Haapsamo H, Pollock-Wurman, R, Kuusikko-Gauffin S, Ebeling H, Larinen K, Soini H & Moilanen I Maternal stress and young children's behavioural development; A prospective pilot study from 8 to 36 months in Finnish sample. Manuscript.

Reprinted with permission from International Journal of Circumpolar Health and Taylor & Francis Group.

Original publications are not included in the electronic version of the dissertation.



1163. Salo, Jarmo (2012) Long-term consequences and prevention of urinary tract infections in childhood
1164. Ronkainen, Veli-Pekka (2012) Effects of chronic hypoxia on myocardial gene expression and function
1165. Kinnula, Sohvi (2012) Hospital-associated infections and the safety of alcohol hand gels in children
1166. Niemelä, Mika (2012) Structured child-centred interventions to support families with a parent suffering from cancer : from practice-based evidence towards evidence-based practice
1167. Pohjolainen, Virva (2012) Characterization of non-collagenous extracellular matrix proteins in cardiac and aortic valve remodelling
1168. Moilanen, Anne-Mari (2012) Identification of novel drug targets for the treatment of heart failure
1169. Jounio, Ulla (2012) Oropharyngeal carriage of respiratory bacteria among military conscripts
1170. Virtanen, Katri (2012) ”Äiti, täällä on toisia samanlaisia, ku mä!” : Voimisteluseura ja kouluterveydenhuolto perheiden tukena lasten painonhallinnassa
1171. Iломäki, Essi (2012) Conduct disorder among girls: violent behaviour, suicidality and comorbidity : A study of adolescent inpatients in Northern Finland
1172. Iломäki, Risto (2012) Substance use disorders in adolescence: Comorbidity, temporality of onset and socio-demographic background : A study of adolescent psychiatric inpatients in Northern Finland
1173. Saxlin, Tuomas (2012) Periodontal infection and obesity—results of a population-based survey
1174. Lämsä, Virpi (2012) Regulation of murine hepatic *Cytochrome P450 2a5* expression by transcription factor Nuclear factor (erythroid-derived 2)-like 2
1175. Huhtakangas, Juha (2012) The influence of medication on the incidence, outcome, and recurrence of primary intracerebral hemorrhage
1176. Puljula, Jussi (2012) Alcohol-related traumatic brain injuries before and after the reduction of alcohol prices : Observations from Oulu Province and Northern Ostrobothnia
1177. Kenttä, Tuomas (2012) Dynamics of cardiac repolarization during exercise : Rate-dependence and prognostic significance

Book orders:

Granum: Virtual book store  
<http://granum.uta.fi/granum/>

S E R I E S E D I T O R S

**A**  
**SCIENTIAE RERUM NATURALIUM**

*Senior Assistant Jorma Arhippainen*

**B**  
**HUMANIORA**

*University Lecturer Santeri Palviainen*

**C**  
**TECHNICA**

*Professor Hannu Heusala*

**D**  
**MEDICA**

*Professor Olli Vuolteenaho*

**E**  
**SCIENTIAE RERUM SOCIALIUM**

*University Lecturer Hannu Heikkinen*

**F**  
**SCRIPTA ACADEMICA**

*Director Sinikka Eskelinen*

**G**  
**OECONOMICA**

*Professor Jari Juga*

**EDITOR IN CHIEF**

*Professor Olli Vuolteenaho*

**PUBLICATIONS EDITOR**

*Publications Editor Kirsti Nurkkala*

ISBN 978-951-42-9961-2 (Paperback)

ISBN 978-951-42-9962-9 (PDF)

ISSN 0355-3221 (Print)

ISSN 1796-2234 (Online)

