

Online patient education design features: are there any differences in opinion based on consumer's stage of change?

Naffisah Mohd Hassan

Faculty of Business and Management
Universiti Teknologi MARA (UiTM) Selangor
Selangor, Malaysia
Email: naffi885@puncakalam.uitm.edu.my

Khin Than Win

School of Computing and Information Technology
Faculty of Engineering and Information Science
University of Wollongong
NSW, 2522
Email: win@uow.edu.au

Harri Oinas-Kukkonen

Information Processing Department
Faculty of Information Technology and Electrical Engineering
University of Oulu
Email: harri.oinas-kukkonen@oulu.fi

Abstract

Patients' stage of change towards their disease management plays an important role in healthcare. The aim of this study is to explore whether there are any differences in the patients' perceptions toward the design features of online patient education according to the patients' stages of change. Patients and carers who use chronic disease websites from Australia were invited to participate in the study to understand their perspectives on the design features. The statistical analysis package for the social science 19.0 (SPSS 19.0) and Smart PLS were used to perform statistical analysis for answering the research questions and related hypotheses. The results indicated that the patients' perceptions tended to differ in terms of online patient education design features, interpretability and presentation of content, according to their stages of change. The patients, across different stages of change, did have differing perceptions regarding certain constructs within the online patient education design features.

Keywords

Design features, online patient education, health education, stage of change, tailoring

1 Introduction

Health education is defined as a systemic learning experience in which a combination of methods is generally used, such as the provision of information, advice and behavioural modification techniques, to influence the way that a patient promotes his or her state of health (Van den Borne 1998) (Win 2010a). With the widespread use of the Internet and the availability of health information online, there is a growing demand for effective educational material online for patients. Traditionally, health education about chronic diseases has fallen within the ambit of doctors, nurses, and other healthcare professionals. Receiving personalised advice and support-material from a healthcare professional with whom a patient has a good relationship is known to be very effective, albeit time consuming (Orleans et al. 1985). Providing Online Patient Education tailored to patients has the potential to assist in patient education. Many benefits have been claimed for online patient education but in order to confer the optimal results, online patient education sites must be designed appropriately.

2 Background

Online patient education is typically targeted towards improving health outcomes, but it can also be used to enhance patients' social support (Wise et al. 2008). Improved health outcomes, improved health education and knowledge acquisition, improved patient awareness of the disease, increased patient confidence towards treatment, improved self-care behaviour and management, reduced hospitalization and adherence to treatment have been identified as benefits in health outcomes associated with online patient education (Win et al. 2015). Additionally, improved quality of interaction with physicians, easy access to educational material, time and cost effectiveness, improved social support and improved patient emotional state and satisfaction have also been noted (Hassan et al. 2013).

The benefits of online patient education for those who suffer from chronic disease can be anticipated to be optimised if the development process is based on design features, which have evidence of their effectiveness. A review of the literature of research into design features of online patient education identified a number of relevant studies, including studies on behaviour change support systems (Hassan et al. 2013). Key design features of importance for online patient education are identified and presented in previous study (Hassan et al. 2013). These features include: patient tailored information; interactivity; credibility; presentation of content; interpretability and multimedia graphics (Win et al. 2016). The design principles of tailoring, interactivity, credibility, and presentation of content are all applicable to the behaviour change support system features of Oinas-Kukkonen and Harjuma (2009) (Matthews et al. 2016). Further details were shown in table 1 below.

| Groups | Design Features | Study |
|---|---|--|
| Patient Tailored Information (5 items) | PTI1: Information tailored to patient's symptoms | (Crutzen et al. 2009; Goessens et al. 2008; Koivunen et al. 2008) |
| | PTI2: Guidance is appropriate for patient's self-care management regime | (Clayman et al. 2008; Ferney and Marshall 2006; McKay et al. 2002) (Heinrich et al. 2012) |
| | PTI3: Advice is tailored to patient's personal treatment preferences | (Doupi and van der Lei 2003; Doupi and van der Lei 2005; Heinrich et al. 2012; Oenema et al. 2001) |
| | PTI4: Tailored feedback provided | (Clayman et al. 2008; de Vet et al. 2008; McKay et al. 2002) |
| | PTI5: Mode of delivery of treatment information | (Ferney and Marshall 2006; Smith and Weinert 2000) |
| Interactivity (6 items) | I-act1: Live chat | (McKay et al. 2002; Rezailashkajani et al. 2008; Smith and Weinert 2000) |
| | I-act2: Animations and interactive learning material | (DeGuzman and Ross 1999; Ferney and Marshall 2006; Kim et al. 2004) |

| | | |
|--------------------------------------|--|---|
| | I-act3: Email function | (Ferney and Marshall 2006; Smith and Weinert 2000) |
| | I-act4: Linked to social networks such as Facebook & Twitter | (Fisher and Clayton 2012; Yamout et al. 2011) |
| | I-act5: Patient forums | (Ferney and Marshall 2006; Rezailashkajani et al. 2008) |
| | I-act6: Interactive quizzes | (Kim et al. 2004; Tate et al. 2001) |
| Content (8 items) | Cont1: Date of content update | (Doupi and Van Der Lei 1999; Morrison et al. 2012) |
| | Cont2: Personal information is secured | (DeGuzman and Ross 1999; Doupi and van der Lei 2003) |
| | Cont3: Accredited by a recognized healthcare organization | (Doupi and Van Der Lei 1999; Eysenbach and Diepgen 2002; Thakurdesai et al. 2004) |
| | Cont4: Patient's rights are displayed on the homepage | (Kim et al. 1999; Smith et al. 2002) |
| | Cont5: Must use login information in order to see patient's personal information | (Smith et al. 2002; Win 2010b) |
| | Cont6: Accreditation by a healthcare organization is displayed | (Doupi and Van Der Lei 1999; Thakurdesai et al. 2004) |
| | Cont7: Author's name and contact information are displayed | (Doupi and Van Der Lei 1999; Eysenbach and Diepgen 1999; Kim et al. 2004) |
| | Cont8: Adherence to quality guidelines | (Kim et al. 2004; Thakurdesai et al. 2004) |
| Presentation of content (6 items) | PC1: Colour contrast-readability | (George et al. 2001; Kim et al. 2004; Smith et al. 2002) |
| | PC2: A screen-reader that vocalizes the textual messages | (Demiris et al. 2001; Kim et al. 2004; Smith et al. 2002) |
| | PC3: Multilingual functions | (Rezailashkajani et al. 2008) |
| | PC4: Descriptive text/captions | (Doupi and van der Lei 2005; Kim et al. 2004; van Weert et al. 2011) |
| | PC5: Simple and realistic pictures illustrating medical concepts | (Doupi and van der Lei 2005; Ferney and Marshall 2006; Kim et al. 2004; Smith et al. 2002; van Weert et al. 2011) |
| | PC6: Easy navigational instruction for all levels of users | (Ferney and Marshall 2006; Kim et al. 2004) |
| Interpretability (2) | I-pret1: Free of medical jargon | (Clayman et al. 2008; Demiris et al. 2001; Rezailashkajani et al. 2008) |
| | I-pret2: Glossary of medical terms provided | (Kinzie et al. 2002; Rezailashkajani et al. 2008) |

Table 1: Items in the Design Features Construct (sources were cited from Hassan et al. 2013)

2.1 Health Behaviour change

Online patient education sites are intended to improve patients' behaviour in terms of adherence to their disease management regimen. The literature indicates that characteristics of the patient are important to consider because some patients might not be ready to change their behaviour towards their healthcare.

Consumers' readiness to change or adopt the targeted behaviour can be identified by the Transtheoretical model (Wright et al. 2015). The model has been widely applied in healthcare (Prochaska et al. 2008b) and used by health professionals for informing, undertaking and evaluating interventions for patient behavioural change (Prochaska 2008). The TTM proposes that people progress through five distinct stages of behaviour change: pre-contemplation, contemplation, preparation, action, and maintenance. These stages describe how people move from being unaware, unwilling, or too discouraged to change, to considering the possibility of change, then to becoming committed and prepared to make the change, and finally taking action and sustaining the change in the longer term (Liang et al. 2006).

Studies on patient behaviours, intervention methods, medication acceptance, smoking cessation, participation in mammography screening, sun protection, reduction of dietary fat consumption and condom use have adopted a TTM framework (Prochaska 2008). It has been proposed that in employing the TTM that interventions need to be tailored according to a patient's stage of change (Prochaska 2008). In addition, the study also indicated that the maintenance stage has been a most difficult stage to sustain among all ((Prochaska et al. 2008a). Different results for health behaviour change have been reported in previous studies. Differences in outcome were noted in a study conducted by Spittaels & de Bourdeudhuij. In their intervention study, the earlier stages have notable changes in behaviour compared to the maintenance stage (Spittaels et al. 2007)). However, a study conducted by de Niet et al. 2012 indicates that providing tailored messages to consumers would optimise the maintenance stage (de Niet et al. 2012). Antypas & Wangberg 2014 also found that patients' stage of change plays an important part in effectiveness of the intervention.

Therefore, it is needed to explore which design features would affect consumers' behaviour change. Studying consumers' preferences of design features from different stages of health behaviour change would be beneficial.

Examples of studies in which stage of change has been considered include the development and design of online material for educating patients with chronic illnesses (Konda et al. 2009; Sanchez et al. 2008). Table 2 provides a summary of the Stages of Change in the TTM (Prochaska et al. 2008b).

| | |
|------------------|---|
| Precontemplation | The consumer has no intention to take action within the next 6 months and is generally unaware or under-aware of the problem. |
| Contemplation | The consumer intends to take action within the next 6 months. He or she is aware that a problem exists but has not yet made a commitment to take action |
| Preparation | The consumer intends to take action within the next 30 days and has taken some behavioural steps in this direction |
| Action | The consumer changes his or her overt behaviour for less than 6 months |
| Maintenance | The consumer changes his or her overt behaviour for more than 6 months and works to prevent relapse and consolidate the gains attained |

Table 2: Transtheoretical model Stages of Change (Prochaska 2008)

2.2 Research questions

The stages of change may differ amongst patients and this may be a key factor to consider when attempting observing the effect of online patient education design features so as to ensure that the user is conferred the maximum benefits.

The aim of this study is to investigate whether there are differences in patients' perceptions of the design features of online patient education according to the patients' stages of change for their health management. The research question posed is: "Is there a difference in the patients' perceptions toward the OPE design features, according to the patients' stages of change?"

In this study, the following hypotheses were formulated in order to answer the research question.

H: There is a significant difference in patients' preferences for online patient education design features according to the patients' stage of change.

However, since there are six (6) dimensions within the online patient education design features, it was sub-divided into six (6) sub-hypotheses.

Ha: There is a significant difference in preferences for Patient Tailored Information design features according to stage of change.

Hb: There is a significant difference in preferences for the Interactivity design features according to stage of change

Hc: There is a significant difference in preference for the credibility design features according to stage of change.

Hd: There is a significant difference in preference for the Presentation of Content design features according to stage of change.

He: There is a significant difference in preference for the Interpretability design features according to stage of change.

Hf: There is a significant difference in preference for the Multimedia graphic design features according to stage of change.




Figure 1: Relationships stage of change and OPE design features

3 Methodology

A stepwise approach was used to identify the design features. OPE design features were identified after literature review and consultation with health professionals, health informaticians and carers. The expert review of the items list was undertaken to validate the applicability of the items in the study context. A pilot study was conducted with 20 participants using semi-structured interviews and closed questions to generate constructs for the design of a quantitative survey instrument. Questionnaires were then refined based on the pilot test results. The interviews were conducted to determine the perceived importance, feasibility and content validity of the questionnaire. Subsequently, another set of questionnaires was administrated to a convenience sample of the target population via the Diabetes Australia website as another pilot test. All participants were chronic disease patients or family members of these patients, and

all had used online patient education sites. Based on the pilot test results, unreliable items from the questionnaire were removed after employing reliability and correlation analyses in order to examine the degree of internal consistency of the questions.

3.1 Studied population and sample

The survey drew upon the users of OPE sites for chronic diseases, such as diabetes and heart disease. For both groups, the data gathered related only to the participant's reaction to an online patient education site. Users of online patient education sites include carers, family members, and chronic disease sufferers; all of whom have a variety of requirements in relation to online patient education. Gathering data from these users is crucial in order to understand the process of designing and using online patient education sites effectively. Sampling: Users of OPE sites were recruited by means of several methods, such as links on various online patient education sites for chronic diseases, and also via emails and telephone calls to friends and associates who fitted the necessary parameters. All potential participants were provided with a link to Survey Gizmo, where the survey was located. Before commencing the survey, participants were given access to a Participant Information Sheet about the research study. The anonymous survey was completed online at the participant's convenience.

4 Analysis

Data were analyzed using SPSS 19.0. The Kruskal-Wallis test was conducted to compare the means of five group of patients for their stages of change against the OPE design features. Chi-square values and the degree of freedoms were tested to interpret the results.

5 Results

One hundred and forty-five (145) participants responded to the instrument during the data collection period. Four participants were excluded as they did not complete 95% of the questions leaving final sample size of n=141, outlined in Table 3.

| Characteristic | | N | % |
|---------------------------|-----------------------|-----|-------|
| Gender | Male | 44 | 31.2% |
| | Female | 97 | 68.8% |
| Age | 25 to 34 years | 14 | 9.9% |
| | 35 to 44 years | 12 | 8.5% |
| | 45 to 54 years | 23 | 16% |
| | 55 to 64 years | 31 | 24% |
| | 65 years and above | 61 | 42% |
| Computer Competency | Very competent | 90 | 64% |
| | Somewhat competent | 51 | 36% |
| Frequency of Internet Use | Once or more a day | 122 | 86.5% |
| | A few times in a week | 18 | 12.8% |
| | Hardly use Internet | 1 | 0.7% |
| Stage of Change (SOC) | Stage 1 | 16 | 11.3% |
| | Stage 2 | 10 | 7.1% |
| | Stage 3 | 23 | 16.3% |
| | Stage 4 | 29 | 20.6% |
| | Stage 5 | 63 | 44.7% |

Table 3, Participants' characteristics

Differences in Patient/Carer Perceptions of Online Patient Education Design Features according to

Patients' Stages of Change

| SOC | Patient Tailored Information | Interactivity | Credibility | Presentation of content | Multimedia graphic | Interpretability |
|-------------|------------------------------|---------------|-------------|-------------------------|--------------------|------------------|
| SOC1 | 72.94 | 89.81 | 62.97 | 88.44 | 78.97 | 83.56 |
| SOC2 | 64.10 | 64.30 | 61.00 | 45.15 | 48.80 | 33.20 |
| SOC3 | 60.59 | 59.48 | 62.80 | 58.54 | 66.26 | 70.61 |
| SOC4 | 78.90 | 74.19 | 78.53 | 79.43 | 76.81 | 85.72 |
| SOC5 | 71.77 | 70.02 | 74.15 | 71.34 | 71.56 | 67.17 |
| Chi-Square | 3.016 | 5.750 | 3.543 | 10.545 | 4.955 | 15.649 |
| df | 4 | 4 | 4 | 4 | 4 | 4 |
| Asymp. Sig. | .555 | .219 | .471 | .032 | .292 | .004 |
| Hypotheses | Reject | Reject | Reject | Accept | Reject | Accept |

TABLE 4: Differences in the Patients' Perceptions toward the Online Patient Education Design Features, according to stage of change

Despite most design features not being perceived significantly differently across the five groups of patients and their respective stages of change; two design features demonstrated significant differences. (See Table 4.) The patients' opinions toward presentation of content and interpretability differ significantly across the patients' stages of change: $\chi^2(4, N = 141) = 10.545, p < .05$ for Presentation of content; and $\chi^2(4, N = 141) = 15.649, p < .05$ for Interpretability. Post-hoc testing was used to confirm these results.

When rejecting the null hypothesis, there is a need to identify the pair(s) of groups that are significantly different. However, the Kruskal-Wallis test does not provide an embedded post-hoc test (Klein and Zhang 2005), (Field 2009). Therefore, further analyses were conducted before making the final decision. Specifically, a separate Mann-Whitney test (for each individual pair) was tested and adjustments were made in order to reduce any potential Type 1 error, by using the Bonferroni correction for the adjustment (Garamszegi 2006). A Mann-test was used for calculating the effect size. $r = z \div \sqrt{N}$

Table 5 shows in detail the significant differences between the groups of patients according to their stages of change and their effect size on the presentation of content features. The differences between r was used to assess the overall effect size, where it has been suggested that 0.02, 0.15, and 0.35 constitute small, moderate, and large effect sizes respectively (Cohen et al. 2003). Thus, for Presentation of content, the groups that represent a medium effect size are SOC1 and SOC2 ($r = -0.20$); groups SOC1 and SOC3 ($r = -0.18$); groups SOC2 and SOC4 ($r = -0.19$); groups SOC2 and SOC5 ($r = -0.17$); and groups SOC3 and SOC4 ($r = -0.15$). On the other hand, the groups that represent a small effect size are SOC1 and SOC4 ($r = -0.07$); groups SOC1 and SOC5 ($r = -0.13$); groups SOC2 and SOC3 ($r = -0.08$); groups SOC3 and SOC5 ($r = -0.12$); and groups SOC4 and SOC5 ($r = -0.08$).

| Between Group | Presentation of Content | | | Interpretability | | |
|---------------|-------------------------|---------|-----------------|------------------|---------|-----------------|
| | Z value | P value | Effect size (r) | Z value | P value | Effect size (r) |
| SOC1 and SOC2 | -2.364 | 0.018 | -0.20 | -3.034 | 0.002 | -0.26 |
| SOC1 and SOC3 | -2.124 | 0.034 | -0.18 | -1.236 | 0.216 | -0.15 |
| SOC1 and SOC4 | -0.826 | 0.409 | -0.07 | -0.375 | 0.708 | -0.03 |

| | | | | | | |
|---------------|--------|-------|-------|--------|-------|-------|
| SOC1 and SOC5 | -1.568 | 0.117 | -0.13 | -1.529 | 0.126 | -0.13 |
| SOC2 and SOC3 | -0.961 | 0.337 | -0.08 | -2.759 | 0.006 | -0.23 |
| SOC2 and SOC4 | -2.210 | 0.027 | -0.19 | -3.218 | 0.001 | -0.27 |
| SOC2 and SOC5 | -1.996 | 0.046 | -0.17 | -2.636 | 0.008 | -0.22 |
| SOC3 and SOC4 | -1.809 | 0.070 | -0.15 | -1.511 | 0.131 | -0.13 |
| SOC3 and SOC5 | -1.391 | 0.164 | -0.12 | -0.434 | 0.664 | -0.04 |
| SOC4 and SOC5 | -0.971 | 0.332 | -0.08 | -2.079 | 0.038 | -0.18 |

TABLE 5: Effect Size for Presentation of Content and Interpretability According to the Different Stages of Change

For Interpretability, the groups that evidenced a medium effect size were SOC1 and SOC2 ($r = -0.26$); groups SOC1 and SOC3 ($r = -0.15$); groups SOC2 and SOC3 ($r = -0.23$); groups SOC2 and SOC4 ($r = -0.27$); groups SOC2 and SOC5 ($r = -0.22$); and groups SOC4 and SOC5 ($r = -0.18$). Finally, the groups that evidenced a small effect size were SOC1 and SOC4 ($r = -0.03$); groups SOC3 and SOC4 ($r = -0.13$); and groups SOC3 and SOC5 ($r = -0.04$).

6 Discussion

From this study, it was noted that the patients' perceptions tend to differ in terms of online patient education design features, according to their stages of change. Consumers' opinions on having tailored information and credibility of the content do not vary much according to behaviour change stage in their disease management. However, preference on perceptions of design features on how the content are presented and, 'the interpretability,' varies. Therefore, the patients, across different stages of change, did have differing perceptions regarding certain constructs within the Online Patient Education design features. Further studies investigating patient health literacy and preferences of design features would assist in further understanding of these responses, particularly regarding the association between differences in opinion of 'the interpretability' and stage of change.

Follow-up tests were subsequently conducted in order to evaluate the pairwise differences amongst the 5 Stage of change groups. In terms of the Presentation of Content features, the earlier stages – i.e. patients in stage 1 (pre-contemplation) or stage 2 (contemplation) – had a medium effect, which was greater than the effect of those patients in the final stages: stage 4 (action) and stage 5 (maintenance).

As for the Interpretability features, patients in stage 1 (pre-contemplation) only evidenced a small effect size compared to patients within the final stages: stage 4 (action) and stage 5 (maintenance). However, the previous groups evidenced a larger effect than those patients did in stage 2 (contemplation) or stage 3 (preparation). This demonstrates that patients within the early stages tend to believe that the interpretability features are more important for the comprehension of the specific terms that are used in the OPE sites. This could be due to the fact that at these stages their understanding of health messages may be poorly developed, they are still at the stage of trying to understand about the disease or not aware of the disease. Therefore, interpretability of the information plays a major role when consumers are not aware of the problem or no intention to change.

Based on the results, few features that derived from persuasive systems design (PSD) are strongly related to benefits of online patient education. Design features from OPE that derived from primary task support category, tailoring, tunneling and self-monitoring features of PSD Model (Oinas-Kukkonen and Harjumaa 2009) are proven to be strongly beneficial to OPE websites. These results are also consistent with interactive features that derived from Computer human dialog and Social support features from PSD model. Designing features according to the consumers' stages of change will be beneficial as persons are usually at different points along a continuum in terms of being ready for initiating behaviour changes and having tailored guidance and feedback (Oenema 2005), self-monitoring activities and reporting the results to others (Antypas and Wangberg 2014) and endorsement and assistance by the person's general practitioner or health professionals (Ammann et al. 2013).

In addition to contributing new knowledge to the online patient education literature, the results obtained here may help healthcare providers, developers, and researchers in developing online patient education sites that are more helpful for patients with chronic diseases. This finding is significant for site developers in understanding users' perceptions toward OPE design features, thus facilitating the process of building better systems and websites which may confer all of the benefits that are expected by the users.

This study also demonstrates a significant difference in the perceptions toward two OPE design features, according to the patients' stages of change. Five stages of motivation to change were used to examine the different perceptions toward the design features of OPE. Developers were advised to take the patients' stages of change into consideration before developing OPE sites, because this is likely to help tremendously in terms of the desired behavioural change aspect associated with the use of OPE sites.

Having a well-defined set of design features to guide the development of OPE sites may ultimately maximize the user-benefits associated with OPE. In addition, these design features and the possible benefits of OPE were actually perceived by users in accordance with their respective stages of change. Developers unintentionally focused more on researching and developing design features for the young and the healthy users, rather than unhealthy users. The results of this study provide an important baseline-instrument in terms of understanding the perceptions of design features according to stages of change.

7 Conclusion

Patient education is to motivate patients to change their behaviour towards healthcare. Understanding patient's stage of change, that is, patient's readiness to change will assist in the health behavioural change process. It is significant for site developers in understanding users' perceptions toward OPE design features, thus facilitating the process of building better systems and websites which may confer all of the benefits that are expected by the users. This is a completed studies done in 2015 and it did not include recent technology innovations such as smart device interfaces and tactile functionality and will be included in future studies by author.

This study demonstrates a significant difference in the perceptions toward two OPE design features, according to the patients' stages of change. It is advisable to developers to take the patients' stages of change into consideration before developing OPE sites, because this is likely to help tremendously in terms of the desired behavioural change aspect associated with the use of online patient education.

8 References

- Ammann, R., Vandelanotte, C., de Vries, H., and Mummery, W. 2013. "Can a Website-Delivered Computer-Tailored Physical Activity Intervention Be Acceptable, Usable, and Effective for Older People?," *Health Education & Behavior* (40:2), pp. 160-170.
- Antypas, K., and Wangberg, S. C. 2014. "An Internet- and Mobile-Based Tailored Intervention to Enhance Maintenance of Physical Activity after Cardiac Rehabilitation: Short-Term Results of a Randomized Controlled Trial," *Journal of Medical Internet Research* (16:3), pp. 78-95.
- Clayman, M. L., Boberg, E. W., and Makoul, G. 2008. "The Use of Patient and Provider Perspectives to Develop a Patient-Oriented Website for Women Diagnosed with Breast Cancer," *Patient Education and Counseling* (72:3), pp. 429-435.
- Cohen, J. B., Cohen, P., West, S. G., and Aiken, L. S. 2003. "Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences (3rd Ed.)." Mahwah, NJ: Lawrence Erlbaum Associates.
- Crutzen, R., de Nooijer, J., Brouwer, W., Oenema, A., Brug, J., and de Vries, N. 2009. "Effectiveness of Online Word of Mouth on Exposure to an Internet-Delivered Intervention," *Psychology and Health* (24:6), pp. 651-661.

- de Niet, J., Timman, R., Bauer, S., van den Akker, E., Buijks, H., de Klerk, C., Kordy, H., and Passchier, J. 2012. "The Effect of a Short Message Service Maintenance Treatment on Body Mass Index and Psychological Well-Being in Overweight and Obese Children: A Randomized Controlled Trial," *Pediatric Obesity* (7:3), pp. 205-219.
- de Vet, E., de Nooijer, J., de Vries, N. K., and Brug, J. 2008. "Testing the Transtheoretical Model for Fruit Intake: Comparing Web-Based Tailored Stage-Matched and Stage-Mismatched Feedback," *Health Education Research* (23:2), pp. 218-227.
- DeGuzman, M. A., and Ross, M. W. 1999. "Assessing the Application of Hiv and Aids Related Education and Counselling on the Internet," *Patient Education and Counseling* (36:3), pp. 209-228.
- Demiris, G., Finkelstein, S. M., and Speedie, S. M. 2001. "Considerations for the Design of a Web-Based Clinical Monitoring and Educational System for Elderly Patients," *Journal of the American Medical Informatics Association* (8:5), pp. 468-472.
- Doupi, P., and Van Der Lei, J. 1999. "R(X) Medication Information for the Public and the Www: Quality Issues," *Medical Informatics and the Internet in Medicine* (24:3), pp. 171-179.
- Doupi, P., and van der Lei, J. 2003. "Design Considerations for a Personalised Patient Education System," *Studies in Health Technology & Informatics* (95), pp. 762-767.
- Doupi, P., and van der Lei, J. 2005. "Design and Implementation Considerations for a Personalized Patient Education System in Burn Care," *International Journal of Medical Informatics* (74:2-4), pp. 151-157.
- Eysenbach, G., and Diepgen, T. L. 1999. "Patients Looking for Information on the Internet and Seeking Teledvice: Motivation, Expectations, and Misconceptions as Expressed in E-Mails Sent to Physicians.[See Comment]," *Archives of Dermatology* (135:2), pp. 151-156.
- Eysenbach, G., and Diepgen, T. L. 2002. "The Role of E-Health and Consumer Health Informatics for Evidence-Based Patient Choice in the 21st Century," *Clinics in Dermatology* (19:1), pp. 11-17.
- Ferney, S. L., and Marshall, A. L. 2006. "Website Physical Activity Interventions: Preferences of Potential Users," *Health Education Research* (21:4), pp. 560-566.
- Field, A. 2009. *Discovering Statistics Using Spss*. Sage publications.
- Fisher, J., and Clayton, M. 2012. "Who Gives a Tweet: Assessing Patients' Interest in the Use of Social Media for Health Care," *Worldviews on Evidence-Based Nursing* (9:2), pp. 100-108.
- Garamszegi, L. Z. 2006. "Comparing Effect Sizes across Variables: Generalization without the Need for Bonferroni Correction," *Behavioral Ecology* (17:4), pp. 682-687.
- George, D., Stanley, M. F., and Stuart, M. S. 2001. "Considerations for the Design of a Web-Based Clinical Monitoring and Educational System for Elderly Patients," *JOURNAL OF THE AMERICAN MEDICAL INFORMATICS ASSOCIATION* (8:5), p. 468.
- Goessens, B. M. B., Visseren, F. L. J., de Nooijer, J., van den Borne, H. W., Algra, A., Wierdsma, J., and van der Graaf, Y. 2008. "A Pilot-Study to Identify the Feasibility of an Internet-Based Coaching Programme for Changing the Vascular Risk Profile of High-Risk Patients," *Patient Education and Counseling* (73:1), pp. 67-72.
- Hassan, N. M., Win, K. T., and Hyland, P. 2013. "Exploring Design Features and Benefits of Online Patient Education (Ope) Sites for Chronic Diseases," *System Sciences (HICSS)*, 2013 46th Hawaii International Conference on, pp. 2454-2463.
- Heinrich, E., de Nooijer, J., Schaper, N. C., Schoonus-Spit, M. H. G., Janssen, M. A. J., and de Vries, N. K. 2012. "Evaluation of the Web-Based Diabetes Interactive Education Programme (Diep) for Patients with Type 2 Diabetes," *Patient Education and Counseling* (86:2), pp. 172-178.
- Kim, P., Eng, T. R., Deering, M. J., and Maxfield, A. 1999. "Published Criteria for Evaluating Health Related Web Sites: Review," *British Medical Journal* (318:7184), pp. 647-649.

- Kim, S., Mouradian, W. E., Leggott, P. J., Schaad, D. C., and Shaul, C. 2004. "Implications for Designing Online Oral Health Resources: A Review of Fifty-Six Websites," *Journal of Dental Education* (68:6), pp. 633-643.
- Kinzie, M. B., Cohn, W. F., Julian, M. F., and Knaus, W. A. 2002. "A User-Centered Model for Web Site Design," *Journal of the American Medical Informatics Association* (9:4), pp. 320-330.
- Klein, J. P., and Zhang, M. J. 2005. *Survival Analysis, Software*. Wiley Online Library.
- Koivunen, M., Hätönen, H., and Välimäki, M. 2008. "Barriers and Facilitators Influencing the Implementation of an Interactive Internet-Portal Application for Patient Education in Psychiatric Hospitals," *Patient Education and Counseling* (70:3), pp. 412-419.
- Konda, K., Ablah, E., Konda, K. S., and Liow, K. 2009. "Health Behaviors and Conditions of Persons with Epilepsy: A Bivariate Analysis of 2006 Brfss Data," *Epilepsy & Behavior* (16:1), pp. 120-127.
- Liang, H., Xue, Y., and Berger, B. A. 2006. "Web-Based Intervention Support System for Health Promotion," *Decision Support Systems* (42:1), pp. 435-449.
- Matthews, J., Win, K., Oinas-Kukkonen, H., and Freeman, M. 2016. "Persuasive Technology in Mobile Applications Promoting Physical Activity: A Systematic Review," *Journal of medical systems* (40:3), pp. 72-72.
- McKay, H. G., Glasgow, R. E., Feil, E. G., Boles, S. M., and Barrera, M. 2002. "Internet-Based Diabetes Self-Management and Support: Initial Outcomes from the Diabetes Network Project," *Rehabilitation Psychology* (47:1), pp. 31-48.
- Morrison, L. G., Yardley, L., Powell, J., and Michie, S. 2012. "What Design Features Are Used in Effective E-Health Interventions? A Review Using Techniques from Critical Interpretive Synthesis," *Telemed J E Health* (18:2), pp. 137-144.
- Oenema, A., Brug, J., and Lechner, L. 2001. "Web-Based Tailored Nutrition Education: Results of a Randomized Controlled Trial," *Health Education Research* (16:6), pp. 647-660.
- Oinas-Kukkonen, H., and Harjumaa, M. 2009. "Persuasive Systems Design: Key Issues, Process Model, and System Features," *Communications of the Association for Information Systems* (24:1), p. 28.
- Orleans, C. T., George, L. K., Houghton, J. L., and Brodie, K. H. 1985. "Health Promotion in Primary Care: A Survey of U.S. Family Practitioners," *Preventive Medicine* (14:5), pp. 636-647.
- Prochaska, J. J., Spring, B., and Nigg, C. R. 2008a. "Multiple Health Behavior Change Research: An Introduction and Overview," *Preventive Medicine* (46:3), pp. 181-188.
- Prochaska, J. O. 2008. "Multiple Health Behavior Research Represents the Future of Preventive Medicine," *Preventive Medicine* (46:3), pp. 281-285.
- Prochaska, J. O., Butterworth, S., Redding, C. A., Burden, V., Perrin, N., Leo, M., Flaherty-Robb, M., and Prochaska, J. M. 2008b. "Initial Efficacy of Mi, Ttm Tailoring and Hri's with Multiple Behaviors for Employee Health Promotion," *Preventive Medicine* (46:3), pp. 226-231.
- Rezailashkajani, M., Roshandel, D., Ansari, S., and Zali, M. R. 2008. "A Web-Based Patient Education System and Self-Help Group in Persian Language for Inflammatory Bowel Disease Patients," *International Journal of Medical Informatics* (77:2), pp. 122-128.
- Sanchez, A., Norman, G. J., Sallis, J. F., Calfas, K. J., Rock, C., and Patrick, K. 2008. "Patterns and Correlates of Multiple Risk Behaviors in Overweight Women," *Preventive Medicine* (46:3), pp. 196-202.
- Smith, C. E., Cha, J., Puno, F., Magee, J. D., Bingham, J., and Van Gorp, M. 2002. "Quality Assurance Processes for Designing Patient Education Web Sites," *Computers Informatics Nursing* (20:5), pp. 191-200.
- Smith, L., and Weinert, C. 2000. "Telecommunication Support for Rural Women with Diabetes," *The Diabetes Educator* (26:4), pp. 645-655.

- Spittaels, H., De Bourdeaudhuij, I., Brug, J., and Vandelanotte, C. 2007. "Effectiveness of an Online Computer-Tailored Physical Activity Intervention in a Real-Life Setting," *Health Education Research* (22:3), pp. 385-396.
- Tate, D. F., Wing, R. R., and Winett, R. A. 2001. "Using Internet Technology to Deliver a Behavioral Weight Loss Program," *JAMA: The Journal of the American Medical Association* (285:9), pp. 1172-1177.
- Thakurdesai, P. A., Kole, P. L., and Pareek, R. P. 2004. "Evaluation of the Quality and Contents of Diabetes Mellitus Patient Education on Internet," *Patient Education & Counseling* (53:3), pp. 309-313.
- Van den Borne, H. W. 1998. "The Patient from Receiver of Information to Informed Decision-Maker," *Patient Education and Counseling* (34:2), pp. 89-102.
- van Weert, J., van Noort, G., Bol, N., van Dijk, L., Tates, K., and Jansen, J. 2011. "Tailored Information for Cancer Patients on the Internet: Effects of Visual Cues and Language Complexity on Information Recall and Satisfaction," *Patient Education and Counseling* (84:3), pp. 368-378.
- Win, K. T. 2010a. "Implementing Patient Accessible Health Information Site for Diabetes Management," in: *Pacific Asia Conference on Information Systems*. Taiwan: National Taiwan University: pp. 1790-1797.
- Win, K. T. 2010b. "Implementing Patient Accessible Health Information Site for Diabetes Management," p. 8.
- Win, K. T., Hassan, N., Bonney, A., and Iverson, D. 2015. "Benefits of Online Health Education: Perception from Consumers and Health Professionals," *Journal of Medical Systems* (39:3), pp. 1-8.
- Win, K. T., Hassan, N. M., Oinas-Kukkonen, H., and Probst, Y. 2016. "Online Patient Education for Chronic Disease Management: Consumer Perspectives," *J Med Syst* (40:4), p. 88.
- Wise, M., Han, J. Y., Shaw, B., McTavish, F., and Gustafson, D. H. 2008. "Effects of Using Online Narrative and Didactic Information on Healthcare Participation for Breast Cancer Patients," *Patient Education & Counseling* (70:3), pp. 348-356.
- Wright, J. A., Whiteley, J. A., Laforge, R. G., Adams, W. G., Berry, D., and Friedman, R. H. 2015. "Validation of 5 Stage-of-Change Measures for Parental Support of Healthy Eating and Activity," *Journal of Nutrition Education and Behavior* (47:2), pp. 134-142.e131.
- Yamout, S. Z., Glick, Z. A., Lind, D. S., Monson, R. A., and Glick, P. L. 2011. "Using Social Media to Enhance Surgeon and Patient Education and Communication," *Bull Am Coll Surg* (96:7), pp. 7-15.

Acknowledgements

The authors would like to thank Universiti Teknologi MARA (UiTM) Selangor, Malaysia and Faculty of Business and Management UiTM Puncak Alam for support and funding this study. The authors also would like to thank Professor Andrew Bonney for providing feedback on the paper.

Copyright

Copyright: © 2016 authors. This is an open-access article distributed under the terms of the [Creative Commons Attribution-NonCommercial 3.0 Australia License](https://creativecommons.org/licenses/by-nc/3.0/australia/), which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and ACIS are credited.