**Recovery of Rate-Pressure Product and Cardiac Mortality in Coronary Artery Disease Patients
with Type 2 Diabetes**

Antti M. Kiviniemi^a, Tuomas V. Kenttä^a, Samuli Lepojärvi^a, Juha S. Perkiömäki^a, Olli-Pekka Piira^a,
Olavi Ukkola^a, Heikki V. Huikuri^a, M. Juhani Juntila^a, Mikko P. Tulppo^a

^aResearch Unit of Internal Medicine, Medical Research Center Oulu, Oulu University Hospital and
University of Oulu, Oulu, Finland.

Corresponding author: Antti M. Kiviniemi, PhD, Research Unit of Internal Medicine, Medical
Research Center Oulu, P.O. Box 5000, 90014 University of Oulu, Oulu, Finland, email:
antti.m.kiviniemi@oulu.fi, tel: +358 8 315 4691

STRUCTURED ABSTRACT

Aims: To investigate prognostic significance of post-exercise recovery of rate-pressure product (RPP) in patients with stable coronary artery disease (CAD) and type 2 diabetes (T2D).

Methods: Patients with angiographically documented CAD and T2D (n=697) underwent symptom-limited bicycle exercise test. Exercise capacity (EC), heart rate, blood pressure and RPP responses to peak exercise and recovery (2' and 5' after cessation of exercise) were analyzed. Cardiac death was the primary and sudden cardiac death (SCD) secondary endpoint.

Results: During a median follow-up of 76 months, 49 cardiac deaths (7.0%) and 28 SCDs (4.0%) were observed. The recovery of RPP at 5' was the strongest univariate predictor of cardiac death (hazard ratio [HR]: 2.55 per SD decrease, 95%CI: 1.82-3.58, $p<0.001$) and SCD (HR: 2.34, 95%CI: 1.51-3.62, $p<0.001$). In multivariate analysis, it remained significantly associated to cardiac death and SCD without (HR: 1.66, 95%CI: 1.14-2.41, $p<0.01$ and HR: 1.75, 95%CI: 1.08-2.85, $p<0.05$, respectively) and with additional adjustment for EC and peak RPP (HR: 1.45, 95%CI: 1.09-1.92, $p<0.05$ and HR: 1.52, 95%CI: 1.01-2.27, $p<0.05$, respectively).

Conclusions: The recovery of RPP after exercise is a potent predictor of cardiac death in patients with CAD and T2D. It provides significant prognostic information beyond EC and peak RPP.

Keywords: Exercise testing; Ischemic heart disease; Autonomic nervous system; Sudden cardiac death

1. Introduction

Rate-pressure product (RPP) is a widely used method to quantify cardiac workload and hemodynamic response exercise. Peak RPP response to exercise test provides important prognostic information in patients with coronary artery disease (CAD) [1-6]. Depressed hemodynamic response to exercise is common feature in type 2 diabetes (T2D) [7, 8]. Predictive value of RPP response to exercise has been reported in patients with diabetes and known or suspected CAD, albeit not independently of other factors [9]. Delayed heart rate (HR) recovery (HRR) is associated with impaired prognosis in patients with CAD [10-12] and T2D [13, 14]. Also, high systolic blood pressure (SBP) during recovery has been associated with risk for cardiovascular morbidities without assessment of the SBP change during recovery in these studies [15-17]. Among patients referred to exercise testing, SBP recovery/peak – ratio was a potent predictor of death but not independently of HRR [18]. It is plausible to expect that RPP recovery would illustrate the combined cardiovascular and autonomic responses and also integrate prognostic significance observed with HRR and SBP recovery. Surprisingly, only Nieminen et al. have reported the outperforming independent prognostic capability of RPP recovery over peak RPP [19]. Yet, the prognostic significance of RPP recovery in patients with stable CAD has remained to be established, particularly among CAD patients with T2D having greater mortality risk and impaired autonomic function [14, 20]. The present study was designed to test the hypothesis that delayed RPP recovery would independently be associated with cardiac mortality and sudden cardiac death (SCD).

2. Materials and methods

2.1 Subjects

The study population comprised of CAD patients with T2D from the ARTEMIS (Innovation to Reduce Cardiovascular Complications of Diabetes at the Intersection, ClinicalTrials.gov identifier: NCT01426685) study database collected in the Division of Cardiology of the Oulu University Hospital

(Oulu, Finland) between August 2007 and June 2014.[14, 20] The ARTEMIS study aims to assess several traditional and novel cardiovascular risk markers as determinants of risk for SCD during 5-year follow-up of patients with stable CAD and T2D. The patients were recruited from a consecutive series of patients with (n=834) and without T2D (n=1,112) who had undergone coronary angiography 3-6 months earlier and had > 3 months from possible previous acute coronary syndrome and revascularization before enrollment and left ventricular (LV) ejection fraction (LVEF) >35%. Exclusion criteria are described elsewhere in details [14, 20]. The patients with contraindication for exercise test or technical problems during the exercise test were excluded from the present study. Therefore, the final sample included 697 CAD patients with T2D (Table 1). Supplementary analyses were performed for 1,089 CAD patients without T2D. The study was conducted according to the Declaration of Helsinki, and the local committee of research ethics of the Northern Ostrobothnia Hospital District approved the protocol, and all the subjects gave written informed consent.

2.2 Exercise testing

An incremental symptom-limited maximal exercise test on a bicycle ergometer was started at 30 W, and the work rate was increased by 15 W in men and 10 W in women every minute until voluntary exhaustion. The patients moved to a supine position within 30 s after cessation of exercise without cool-down period. Supine resting, maximal, peak exercise and recovery (2 and 5 min post-exercise) HRs (15-lead ECG, GE Healthcare, CAM-14, Freiburg, Germany) and BPs (Dinamap Procare 100, GE Medical Systems, Milwaukee, WI) were obtained. From resting ECG, QRS duration, QT interval, QTc interval, Q wave and ECG-based left ventricular hypertrophy were additionally analyzed by custom-made software [21]. EC was calculated as metabolic equivalents (METs) from the mean workload during the last minute of the test. Predicted EC ($EC_{\text{Predicted}} = 18 - [0.15 \cdot \text{age}]$ for men, $EC_{\text{Predicted}} = 14.7 - [0.13 \cdot \text{age}]$ for women) was used to obtain age- and sex-specific reference values for the calculation

of relative EC ($EC\% = 100 \cdot EC \cdot EC_{\text{Predicted}}^{-1}$). [22] Predicted EC was based on large database of treadmill tests [22] that result approximately in ~1 MET larger EC than bicycle ergometer [23]. In the calculation of relative EC, it was assumed that the effects of age and sex on EC are similar regardless of the mode of exercise. Age-adjusted (% of $220 - \text{age}$) maximal HR and chronotropic response index ($CRI = 100 \cdot (\text{maximal HR} - \text{resting HR}) \cdot (220 - \text{age} - \text{resting HR})^{-1}$) were calculated. HRR was analyzed by subtracting the post-exercise HR from the HR at peak exercise. Similarly, SBP and RPP ($=HR \cdot SBP$) responses to exercise (peak – resting) and recovery (peak – recovery) were calculated. Additionally, RPP recovery was normalized to peak RPP ($RPP \text{ recovery } [\%] = 100 \cdot RPP \text{ recovery} \cdot \text{peak RPP}^{-1}$). The incidence of angina pectoris and the level of ST-segment were also registered. Maximal T-wave alternans was detected during the exercise and recovery using modified moving average method described elsewhere in details [24].

2.3. Other measurements

All laboratory measurements and exercise tests were done after 12-hour overnight fast and the patients were instructed to avoid smoking before the tests. Venous blood samples and urine samples were obtained for the analysis of renal function, blood lipids, plasma glucose, and glycated haemoglobin levels. Those subjects without history of diabetes underwent an oral glucose tolerance test to establish their glucose metabolism status. Two-dimensional, M-mode and tissue Doppler echocardiography was performed utilizing the same ultrasound machine for all of the patients (Vivid 7, GE Healthcare, Wauwatosa, WI). LV mass index and LVEF were measured by the biplane method from 2- and 4-chamber views. The patients filled in a health questionnaire containing a question about the frequency of habitual leisure-time physical activity (LTPA). Four LTPA groups were formed by modifying a scale originally developed by Saltin and Grimby: 1) no LTPA (hardly any physical activity or only light housework; 2) LTPA irregularly (some light physical activity randomly, e.g., walking or

cycling); 3) moderate-intensity LTPA regularly two to three times weekly; and 4) moderate- or high-intensity LTPA more than three times weekly, where “time” means a period of 30 min [25].

2.4. Follow-up and end-points

End-points were determined from emergency rescue reports, hospital and physician records, autopsy data, death certificates, and interviews with next of kin. The cause and mode of death was reviewed and adjudicated by two independent investigators; and, if needed, disagreement or uncertainty was resolved in consultation with the investigators (MJJ, HVH). The primary end-point in this study was cardiac death or resuscitation from sudden cardiac arrest, whichever occurred first. SCD, including resuscitations from sudden cardiac arrests, was defined as secondary end-point. The death was defined SCD if it was witnessed and occurred within 1 hour of the onset of symptoms or, in case of unwitnessed death, if patient was seen alive and stable 24 hours prior to discovery.

2.5. Statistical analyses

The between-group differences were assessed by one-way analysis of variance, Kruskal-Wallis or chi-square followed by post hoc analyses by Bonferroni, Mann-Whitney U-test or chi-square adjusted for multiple comparisons when applicable. Linear regression with collinearity statistics was performed when applicable. The predictive powers of the risk markers were assessed by univariate Cox regression followed by adjustment for established risk markers (age, sex, CCS class, LTPA, glycated haemoglobin, LVEF). In the complementary analyses, EC and peak RPP were also included as covariates. The hazard ratios are presented as risk per standard deviation (SD) decrease with 95% CI. Interactions of exercise variables (divided into 3 groups by; $< \text{mean} - 1 \text{ SD}$, $\text{mean} \pm 1 \text{ SD}$ and $\geq \text{mean} + 1 \text{ SD}$) with sex were also studied by Cox regression. Competing risks survival analyses were also conducted using approaches of cardiac death vs. non-cardiac death and SCD vs. other cause of death.

The discrimination abilities of the exercise test variables were assessed by the C-index and the integrated discrimination index (IDI). Continuous net reclassification index (NRI) was also analyzed based on the individual predicted risks by multivariate Cox regression. Kaplan-Meier analysis was used to illustrate survival curves of different groups of risk. The data were analyzed using SPSS software (IBM SPSS Statistics 21, IBM Corp., New York, USA) and R Statistics (3.3.1, The R Foundation for Statistical Computing, Vienna, Austria). A p-value <0.05 was considered as statistically significant.

3. Results

During the median follow-up of 76 months (in survivors; 1st-3rd quartile: 58-96 months), there were 49 cardiac deaths (7.0%), 43 non-cardiac deaths (6.2%) and 28 SCDs (4.0%). All the exercise test variables were significantly associated with cardiac death and SCD in univariate Cox regression, except EC for SCD (Table 2). One SD lower RPP_{Peak} involved 124% and 105% higher risk for cardiac death and SCD, respectively, whereas corresponding values with $RPP_{Recovery5'}$ were 155% and 134% (Table 2, Figure 1). After adjustment for potential confounders, only $HR_{Recovery5'}$, $SBP_{Recovery5'}$ and $RPP_{Recovery5'}$ were associated with cardiac death and SCD (Table 2), the $RPP_{Recovery5'}$ being the most significant and involved 66% and 75% higher risk per 1 SD decrease in $RPP_{Recovery5'}$ for cardiac death and SCD, respectively. With additional adjustment for EC, $RPP_{Recovery5'}$ remained significant factor underlying risk for cardiac death (hazard ratio: 1.61, 95%CI: 1.10-2.38, $p=0.015$) and SCD (hazard ratio: 1.89, 95%CI: 1.13-3.13, $p=0.014$). No significant interactions with sex were observed.

Neither exercise ST-depression/occurrence of angina nor T-wave alternans were associated with the outcomes (Table 1). The resting QRS duration >110 ms was not associated to cardiac death (hazard ratio: 1.56, 95%CI: 0.84-2.89, $p=0.162$) but did so for SCD (hazard ratio: 2.91, 95%CI: 1.37-6.15, $p=0.005$) in univariate analysis. However, it did not remain significant predictor of SCD after adjustments (hazard ratio: 1.36, 95%CI: 0.59-3.17, $p=0.474$). The abnormal QTc interval was

associated with cardiac death (hazard ratio: 2.21, 95%CI: 1.24-3.96, $p=0.007$) and SCD (hazard ratio: 3.23, 95%CI: 1.54-6.81, $p=0.002$) but did not remain significant after adjustments (hazard ratio: 1.05, 95%CI: 0.55-2.00, $p=0.890$ for cardiac death and hazard ratio: 1.47, 95%CI: 0.63-3.43, $p=0.374$ for SCD).

$RPP_{\text{Recovery5'}}$ correlated significantly with RPP_{Peak} ($r=0.903$, $p<0.001$) and had significant collinearity (variance inflation factor [VIF] = 5.4). Normalization of $RPP_{\text{Recovery5'}}$ to RPP_{Peak} decreased correlation ($r=0.651$, $p<0.001$) and collinearity to acceptable level (VIF = 1.7) between $RPP_{\text{Recovery5'}}$ and RPP_{Peak} . In the multivariate Cox regression, relative $RPP_{\text{Recovery5'}}$ remained significantly associated to cardiac death (hazard ratio: 1.45, 95%CI: 1.09-1.92, $p=0.011$) and SCD (hazard ratio: 1.52, 95%CI: 1.01-2.27, $p=0.045$) when also EC and RPP_{Peak} were included as covariates.

$RPP_{\text{Recovery5'}}$ was associated with cardiac death regardless of competing risk for non-cardiac death in univariate (hazard ratio: 2.38, 95%CI: 1.69-3.33, $p<0.001$) and multivariate analyses (hazard ratio: 1.67, 95%CI: 1.12-2.44, $p=0.011$, adjusted also for EC). Similarly, $RPP_{\text{Recovery5'}}$ was associated with SCD regardless of competing risk for other types of death in univariate (hazard ratio: 2.17, 95%CI: 1.37-3.45, $p<0.001$) and multivariate analyses (hazard ratio: 1.82, 95%CI: 1.11-2.94, $p=0.017$, adjusted also for EC).

Addition of $RPP_{\text{Recovery5'}}$ to the established model without EC improved discrimination of the patients regarding cardiac death (IDI: 0.032, 95%CI: 0.002-0.077, $p=0.020$; C-index from 0.763, 95%CI: 0.678-0.849 to 0.773, 95%CI: 0.691-0.856) and tended to improve classification of the patients (NRI: 0.164, 95%CI: -0.013-0.343, $p=0.086$). When EC was included in the established model, the addition of $RPP_{\text{Recovery5'}}$ did improve neither discrimination nor classification. For the SCD, the $RPP_{\text{Recovery5'}}$ did not improve discrimination (IDI: 0.020, 95%CI: -0.007-0.075, $p=0.100$; C-index from 0.676, 95%CI: 0.537-0.814 to 0.714, 95%CI: 0.571-0.858) or classification (NRI: 0.167, 95%CI: -0.108-0.389, $p=0.126$).

In the CAD patients without T2D, all exercise test variables, except for SBP_{Recovery2'}, were associated with cardiac mortality in the univariate analysis (Table 3). After adjustments for potential confounders, only EC, CRI and HR_{Recovery2'} remained significant factors underlying the risk for cardiac death. The RPP variables were not associated to cardiac death or SCD in the multivariate analysis in CAD patients without T2D.

4. Discussion

The main finding of the present study was that among exercise test variables post-exercise recovery of RPP was the most powerful factor related to the risk for cardiac death and SCD in CAD patients with T2D. Notably, it provided significant prognostic information beyond EC and RPP at peak exercise and outperformed the prognostic significance of standard resting and exercise ECG markers. However, it provided limited improvement in discrimination and reclassification analyses when added to the established risk model.

The present study confirmed our hypothesis that RPP recovery provides significant prognostic value in patients with stable CAD and T2D. It was associated with risk for cardiac death and SCD independently of established risk markers and competing risks for other types of death. Notably, the predictive value of HRR and recovery of SBP and RPP, measured at 5 min post-exercise, turned out to be greater than measures quantifying hemodynamic responses to peak exercise that were not associated with cardiac death or SCD in multivariate analyses. Evidently, prognostic significances of HRR and SBP recovery were combined with RPP recovery. Therefore, as readily available index, RPP recovery is a potent marker for risk stratification in patients with CAD and T2D in clinical practice. However, it yielded borderline improvements in discrimination and classification of the patients and that is why confirmatory studies are warranted.

The recovery of RPP, as absolute value, was strongly related to RPP_{Peak} which compromised multivariate analysis with both variables included. Therefore, RPP recovery was also normalized to RPP_{Peak} , which considerably decreased correlation and collinearity between these markers. Subsequently, it was also observed that relative RPP recovery was associated with the outcomes independently of RPP_{Peak} as well as EC. This underscores the greater predictive value of RPP recovery compared to RPP_{Peak} and suggests the importance of post-exercise monitoring of hemodynamics in exercise testing. While prognostic significance of RPP response to peak exercise has been promising in many populations [1-6, 26], it has provided limited additional value to EC and HR responses to exercise in patients referred to exercise test and without CAD [27]. Also, in small sample of patients with T2D, RPP response to exercise was not associated with the presence of silent myocardial ischemia independently of EC and HR responses to exercise and recovery [28]. Taken together, RPP recovery might be more potent and independent prognostic marker for cardiac death and SCD in patients with stable CAD and T2D. The latter notion is also-supported by the findings by Nieminen et al. among patients referred to exercise testing [19].

Our supplementary analyses in CAD patients without T2D showed significant univariate associations of exercise and recovery variables to cardiac death and SCD. However, only HRR and HR response to exercise predicted outcome in the multivariate analyses. The risk ratios regarding these variables were close to those observed in CAD patients with T2D and the small number of endpoints in CAD patients without T2D most likely explains the lack of significant association of RPP recovery in these sub-analyses. Therefore, it is plausible to consider that exercise and recovery responses in central hemodynamics are significant risk factors in CAD patients regardless of glycemic status.

The main hemodynamic response contributor to exercise and recovery is autonomic nervous system which plays an important part in pathophysiology of cardiac death and SCD [29-32]. Impaired exercise responses in HR and SBP are mostly explained by depressed vagal and augmented

sympathetic activity at rest accompanied with modest vagal withdrawal and sympathetic response to exercise that additionally limit the EC [29]. Pain and ischemia may also decrease these responses [33]. During the post-exercise recovery, there is faster vagal reactivation followed by slower sympathetic withdrawal [34]. In the present study, recovery indexes measured at 5 min after exercise were more predictive than those measured at 2 min. Therefore, impaired sympathetic response to recovery may be the main culprit in the present observations. Accumulation of metabolites in active muscle opposes vascular sympathetic responses – a phenomenon called sympatholysis [35]. In turn, sympathetic vasoconstriction persists in inactive muscles. Impaired sympatholysis with persistent vasoconstriction in the inactive muscles may, therefore, delay recovery of blood pressure regardless of cardiac autonomic responses that is opposed also by atrial reflex due to sustained venous return. As impaired sympatholysis has not been verified in T2D, it has been hypothesized that advanced T2D with endothelial dysfunction may alter this mechanism [36] that may well explain the abnormal SBP and, thus, RPP recovery in the high risk CAD patients with T2D.

The present study is partly limited by the small number of endpoints why only the most relevant covariates were allowed to be included in the established model. In turn, the use of continuous risk variables decreases the possibility for overly optimized results. Therefore, confirmatory studies are warranted. The adjudication of SCD, the present secondary endpoint, raises questions about the reliability of the outcome because death certificates have often been considered as unreliable method of adjudication. However, medicolegal autopsy is mandatory according to law in Finland whenever a death has been sudden and unexpected even when the prior cardiac condition is known which increases our confidence in adjudication of SCD.

In conclusion, the recovery of rate-pressure product after exercise is a potent predictor of cardiac death in patients with coronary artery disease and diabetes. It provides significant prognostic

information beyond exercise capacity and peak rate-pressure product and is potent candidate for risk stratification in patients with ischemic heart disease and diabetes.

Acknowledgments

The authors gratefully acknowledge the assistance of the registered nurses Pirkko Huikuri, Päivi Koski, Päivi Kastell and Sari Kaarlenkaski. No relationships to disclose.

Funding

This work was supported by Finnish Technology Development Center, TEKES, Helsinki, Finland [grant number n/a, 2007-2014]; the Academy of Finland, Helsinki, Finland [grant number #267435, 2014-2017]; the Sigrid Juselius Foundation, Helsinki, Finland [grant number n/a, 2018], the Finnish Foundation for Cardiovascular Research, Helsinki, Finland [grant number n/a, 2017] and the Paulo Foundation, Espoo, Finland [grant number n/a, 2016]. The authors appreciate the financial support received from the ARTEMIS consortium partners (Polar Electro, Kempele, Finland and Hur Oy, Kokkola, Finland).

References

- [1] Ottenhof MJ, Wai MC, Boiten HJ, Korbee RS, Valkema R, van Domburg RT, et al. 12-Year outcome after normal myocardial perfusion SPECT in patients with known coronary artery disease. *Journal of nuclear cardiology : official publication of the American Society of Nuclear Cardiology*. 2013;20:748-54.
- [2] Elhendy A, Modesto KM, Mahoney DW, Khandheria BK, Seward JB, Pellikka PA. Prediction of mortality in patients with left ventricular hypertrophy by clinical, exercise stress, and echocardiographic data. *J Am Coll Cardiol*. 2003;41:129-35.

- [3] Bigi R, Cortigiani L, Gregori D, De Chiara B, Parodi O, Fiorentini C. Exercise versus recovery electrocardiography for predicting outcome in hypertensive patients with chest pain. *J Hypertens.* 2004;22:2193-9.
- [4] Saunamaki KI, Andersen JD. Clinical significance of the ST-segment response and other early exercise test variables in uncomplicated vs complicated myocardial infarction. *Eur Heart J.* 1987;8:603-10.
- [5] Saunamaki KI, Damgaard Andersen J. Early exercise test in the assessment of long-term prognosis after acute myocardial infarction. *Acta medica Scandinavica.* 1981;209:185-91.
- [6] Villella M, Villella A, Barlera S, Franzosi MG, Maggioni AP. Prognostic significance of double product and inadequate double product response to maximal symptom-limited exercise stress testing after myocardial infarction in 6296 patients treated with thrombolytic agents. GISSI-2 Investigators. Gruppo Italiano per lo Studio della Sopravvivenza nell-Infarto Miocardico. *Am Heart J.* 1999;137:443-52.
- [7] Kahn JK, Zola B, Juni JE, Vinik AI. Decreased exercise heart rate and blood pressure response in diabetic subjects with cardiac autonomic neuropathy. *Diabetes Care.* 1986;9:389-94.
- [8] Khoshdel AR, Carney SL, White S. Disturbed hemodynamic cardiac exercise stress test response in non-smoking, normolipidemic, normotensive, diabetic subjects. *Diabetes Res Clin Pract.* 2007;75:193-9.
- [9] Innocenti F, Agresti C, Baroncini C, Caldi F, Mannucci E, Monami M, et al. Prognostic value of dobutamine stress echocardiography in diabetic patients. *The international journal of cardiovascular imaging.* 2010;26:499-507.
- [10] Kiviniemi AM, Lepojarvi S, Kentta TV, Junttila MJ, Perkiomaki JS, Piira OP, et al. Exercise capacity and heart rate responses to exercise as predictors of short-term outcome among patients with stable coronary artery disease. *Am J Cardiol.* 2015;116:1495-501.

- [11] Nissinen SI, Makikallio TH, Seppanen T, Tapanainen JM, Salo M, Tulppo MP, et al. Heart rate recovery after exercise as a predictor of mortality among survivors of acute myocardial infarction. *Am J Cardiol.* 2003;91:711-4.
- [12] Vivekananthan DP, Blackstone EH, Pothier CE, Lauer MS. Heart rate recovery after exercise is a predictor of mortality, independent of the angiographic severity of coronary disease. *J Am Coll Cardiol.* 2003;42:831-8.
- [13] Chacko KM, Bauer TA, Dale RA, Dixon JA, Schrier RW, Estacio RO. Heart rate recovery predicts mortality and cardiovascular events in patients with type 2 diabetes. *Med Sci Sports Exerc.* 2008;40:288-95.
- [14] Karjalainen JJ, Kiviniemi AM, Hautala AJ, Piira OP, Lepojarvi ES, Peltola MA, et al. Determinants and prognostic value of cardiovascular autonomic function in coronary artery disease patients with and without type 2 diabetes. *Diabetes Care.* 2014;37:286-94.
- [15] Laukkanen JA, Kurl S, Salonen R, Lakka TA, Rauramaa R, Salonen JT. Systolic blood pressure during recovery from exercise and the risk of acute myocardial infarction in middle-aged men. *Hypertension.* 2004;44:820-5.
- [16] Yosefy C, Jafari J, Klainman E, Brodtkin B, Handschumacher MD, Vaturi M. The prognostic value of post-exercise blood pressure reduction in patients with hypertensive response during exercise stress test. *Int J Cardiol.* 2006;111:352-7.
- [17] Laukkanen JA, Willeit P, Kurl S, Makikallio TH, Savonen K, Ronkainen K, et al. Elevated systolic blood pressure during recovery from exercise and the risk of sudden cardiac death. *J Hypertens.* 2014;32:659-66.
- [18] Ellis K, Pothier CE, Blackstone EH, Lauer MS. Is systolic blood pressure recovery after exercise a predictor of mortality? *Am Heart J.* 2004;147:287-92.

- [19] Nieminen T, Leino J, Maanoja J, Nikus K, Viik J, Lehtimäki T, et al. The prognostic value of haemodynamic parameters in the recovery phase of an exercise test. The Finnish Cardiovascular Study. *Journal of human hypertension*. 2008;22:537-43.
- [20] Junttila MJ, Kiviniemi AM, Lepojärvi ES, Tulppo MP, Piira OP, Kenttä TV, et al. Type 2 diabetes and coronary artery disease: preserved ejection fraction and sudden cardiac death. *Heart Rhythm*. 2018;(in press).
- [21] Holkeri A, Eranti A, Kenttä TV, Noponen K, Haukilahti MAE, Seppanen T, et al. Experiences in digitizing and digitally measuring a paper-based ECG archive. *J Electrocardiol*. 2018;51:74-81.
- [22] Kim ES, Ishwaran H, Blackstone E, Lauer MS. External prognostic validations and comparisons of age- and gender-adjusted exercise capacity predictions. *J Am Coll Cardiol*. 2007;50:1867-75.
- [23] Maeder M, Wolber T, Atefy R, Gadza M, Ammann P, Myers J, et al. Impact of the exercise mode on exercise capacity: bicycle testing revisited. *Chest*. 2005;128:2804-11.
- [24] Verrier RL, Klingenhöfen T, Malik M, El-Sherif N, Exner DV, Hohnloser SH, et al. Microvolt T-wave alternans physiological basis, methods of measurement, and clinical utility--consensus guideline by International Society for Holter and Noninvasive Electrocardiology. *J Am Coll Cardiol*. 2011;58:1309-24.
- [25] Saltin B, Grimby G. Physiological analysis of middle-aged and old former athletes. Comparison with still active athletes of the same ages. *Circulation*. 1968;38:1104-15.
- [26] Sadrzadeh Rafie AH, Sungar GW, Dewey FE, Hadley D, Myers J, Froelicher VF. Prognostic value of double product reserve. *Eur J Cardiovasc Prev Rehabil*. 2008;15:541-7.
- [27] Ahmed HM, Al-Mallah MH, McEvoy JW, Nasir K, Blumenthal RS, Jones SR, et al. Maximal Exercise Testing Variables and 10-Year Survival: Fitness Risk Score Derivation From the FIT Project. *Mayo Clinic proceedings*. 2015;90:346-55.

- [28] Yamada T, Yoshitama T, Makino K, Lee T, Saeki F. Heart rate recovery after exercise is a predictor of silent myocardial ischemia in patients with type 2 diabetes. *Diabetes Care*. 2011;34:724-6.
- [29] Brubaker PH, Kitzman DW. Chronotropic incompetence: causes, consequences, and management. *Circulation*. 2011;123:1010-20.
- [30] Imai K, Sato H, Hori M, Kusuoka H, Ozaki H, Yokoyama H, et al. Vagally mediated heart rate recovery after exercise is accelerated in athletes but blunted in patients with chronic heart failure. *J Am Coll Cardiol*. 1994;24:1529-35.
- [31] Schwartz PJ, Billman GE, Stone HL. Autonomic mechanisms in ventricular fibrillation induced by myocardial ischemia during exercise in dogs with healed myocardial infarction. An experimental preparation for sudden cardiac death. *Circulation*. 1984;69:790-800.
- [32] Huikuri HV, Castellanos A, Myerburg RJ. Sudden death due to cardiac arrhythmias. *N Engl J Med*. 2001;345:1473-82.
- [33] Bacon SL, Lavoie KL, Campbell TS, Fleet R, Arsenault A, Ditto B. The role of ischaemia and pain in the blood pressure response to exercise stress testing in patients with coronary heart disease. *Journal of human hypertension*. 2006;20:672-8.
- [34] Pecanha T, Silva-Junior ND, Forjaz CL. Heart rate recovery: autonomic determinants, methods of assessment and association with mortality and cardiovascular diseases. *Clin Physiol Funct Imaging*. 2014;34:327-39.
- [35] Saltin B, Mortensen SP. Inefficient functional sympatholysis is an overlooked cause of malperfusion in contracting skeletal muscle. *J Physiol*. 2012;590:6269-75.
- [36] Thaning P, Bune LT, Zaar M, Saltin B, Rosenmeier JB. Functional sympatholysis during exercise in patients with type 2 diabetes with intact response to acetylcholine. *Diabetes Care*. 2011;34:1186-91.

Figures

Figure 1. Kaplan-Meier survival curves of the groups according to recovery of rate-pressure product

($RPP_{Recovery5'}$) in patients with coronary artery disease and type 2 diabetes

Table 1. Characteristics of patients with coronary disease and type 2 diabetes according to specific endpoint-groups.

		Survivors	Cardiac death	Non-cardiac death
Variable		n=605	n=49	n=43
Age (years)		66 (8)	71 (7)‡	73 (8)‡
Men		424 (70%)	41 (84%)*	39 (91%)*
Body mass index (kg/m ²)		29.9 (4.6)	28.7 (4.7)	29.1 (5.0)
Resting heart rate (bpm)		64 (10)	65 (11)	65 (11)
Resting systolic blood pressure (mmHg)		148 (24)	143 (24)	143 (25)
Resting diastolic blood pressure (mmHg)		81 (12)	80 (11)	74 (12)‡§
Smokers		51 (9%)	3 (6%)	3 (7%)
Leisure-time physical activity	Inactive	72 (12%)	16 (33%)*	7 (16%)
	Irregularly active	254 (42%)	17 (35%)	20 (47%)
	Active	198 (33%)	11 (22%)	10 (23%)
	Highly active	81 (13%)	5 (10%)	6 (14%)
History of acute myocardial infarction		275 (46%)	30 (61%)	17 (40%)
History of revascularization		485 (81%)	40 (82%)	36 (84%)
CCS class	1	324 (54%)	16 (33%)*	20 (47%)

	2	225 (37%)	22 (45%)	17 (40%)
	3	56 (9%)	11 (22%)*	6 (14%)
Post-operation syntax score		1 (0-5)	5 (0-16)†	2 (0-8)
Duration of diabetes (months)		62 (24-138)	96 (24-234)	126 (21-276)
<hr/> Medication				
Beta-blockers		551 (91%)	46 (94%)	40 (93%)
Angiotensin converting enzyme inhibitors or receptor II blockers		474 (79%)	32 (85%)	34 (79%)
Calcium channel blockers		198 (33%)	14 (29%)	15 (35%)
Diuretics		268 (44%)	29 (59%)	24 (56%)
Anticholesterol agents		558 (92%)	39 (80%)*	39 (91%)
Antihyperglycemic agents		462 (76%)	32 (65%)	30 (70%)
Insulin		149 (25%)	24 (49%)*	13 (31%)
<hr/> Echogardiography				
Left ventricular ejection fraction (%)		64 (9)	57 (15)‡	65 (10)¶
Left ventricular mass index (g/m ²)		108 (27)	126 (31)‡	117 (29)

Laboratory markers

Glycated haemoglobin (%)	6.9 (1.1)	7.4 (1.8)†	7.4 (1.7)*
(mmol/mol)	52 (12)	58 (21)†	58 (19)*
Glucose (mmol/L)	7.43 (1.83)	8.61 (3.66)†	8.35 (3.28)*
Total cholesterol (mmol/L)	3.85 (0.81)	4.27 (1.12)*	3.61 (0.80)¶
(mg/dL)	149 (31)	165 (43)	140 (31)
LDL cholesterol (mmol/L)	2.19 (0.70)	2.49 (0.90)*	2.03 (0.69)
(mg/dL)	85 (27)	96 (35)	78 (27)
HDL cholesterol (mmol/L)	1.19 (0.29)	1.16 (0.35)	1.12 (0.25)
(mg/dL)	46 (11)	45 (14)	43 (10)
Triglycerides (mmol/L)	1.38 (1.00-1.91)	1.42 (1.01-2.21)	1.31 (1.07-1.82)
(mg/dL)	122 (89-169)	126 (89-196)	116 (95-161)
Estimated glomerular filtration rate (mL/min)	105 (40)	81 (35)‡	88 (33)*
Albumin/creatinine-ratio	1.0 (0.6-1.8)	1.5 (1.0-2.8)†	1.7 (0.8-8.2)†
Resting electrocardiography			
QRS duration (ms)	102 (17)	108 (18)	113 (27)†
QRS duration >110 ms	121 (20%)	14 (29%)	14 (33%)
QT interval (ms)	419 (35)	428 (36)	431 (32)

QTc interval (ms)		427 (27)	440 (28) [†]	444 (32) [‡]
QTc interval >440/460 ms (♀/♂)		124 (21%)	18 (37%)*	20 (47%)*
Q wave		102 (17%)	14 (29%)	7 (16%)
ECG-based left ventricular hypertrophy		53 (9%)	8 (16%)	2 (5%)
<hr/> Exercise test				
ST-depression	> 0.1 mV or angina	270 (45%)	24 (49%)	14 (33%)
	> 0.2 mV or angina	73 (12%)	7 (14%)	6 (14%)
T wave alternans (μV)	Exercise+Recovery	42 (22)	47 (25)	43 (18)
	Exercise	39 (22)	44 (25)	41 (20)
	Recovery	24 (14)	27 (15)	25 (12)
Exercise capacity	(METs)	5.6 (1.5)	4.7 (1.4) [‡]	4.8 (1.5) [†]
	(%)	76 (18)	68 (21)*	70 (20)
Heart rate (bpm)	Peak	123 (22)	110 (21) [‡]	111 (20) [†]
	(%)	80 (13)	74 (13) [†]	75 (12)
	Reserve	59 (21)	45 (18) [‡]	46 (18) [‡]
	(%)	67 (24)	54 (22) [†]	56 (20)*
	Recovery 2'	35 (13)	26 (13) [‡]	26 (12) [‡]

	Recovery 5'	44 (15)	33 (14)‡	35 (15)†
Systolic blood pressure (mmHg)	Peak	193 (30)	172 (40)‡	174 (29)‡
	Response	50 (30)	30 (36)‡	36 (29)*
	Recovery 2'	9 (27)	-3 (23)†	1 (24)
	Recovery 5'	31 (32)	6 (35)‡	16 (35)*
Rate-pressure product (bpm·mmHg)	Peak	24,133 (6,698)	19,220 (6991)‡	19,481 (5,742)‡
	Response	14,956 (6,407)	10,063 (6515)‡	10,567 (5,596)‡
	Recovery 2'	7,864 (4,581)	4,634 (4066)‡	5,090 (3,971)‡
	Recovery 5'	11,218 (5,714)	6,453 (5655)‡	7,592 (5,565)‡

Values are mean (SD), median (1st-3rd quartile) or n (% within group). *CCS class* Canadian Cardiovascular Society grading of angina pectoris, *LDL* low-density lipoprotein, *HDL* high-density lipoprotein. */†/‡ p<0.05/0.01/0.001 compared to survivors, §/||/¶ p<0.05/0.01/0.001 compared to patients with cardiac death.

Table 2. Predictive value of exercise test variables in patients with coronary artery disease and type 2 diabetes.

	Univariate		Multivariate	
	Cardiac death	Sudden cardiac	Cardiac death	Sudden cardiac
	(n=49)	death (n=28)	(n=49)	death (n=28)
	Hazard ratio per	Hazard ratio per	Hazard ratio per	Hazard ratio per
	+1 SD (95%CI)	+1 SD (95%CI)	+1 SD (95%CI)	+1 SD (95%CI)
Exercise capacity (%)	1.57 (1.15-2.15)†	1.41 (0.94-2.10)	1.28 (0.87-1.89)	1.01 (0.60-1.72)
Heart rate				
Peak (%)	1.66 (1.20-2.28)†	1.62 (1.07-2.47)*	1.28 (0.93-1.75)	1.35 (0.88-2.08)
Chronotropic response index (%)	1.96 (1.35-2.84)‡	1.85 (1.34-3.00)*	1.35 (0.94-1.93)	1.41 (0.87-2.31)
Recovery 2' (bpm)	2.13 (1.53-2.97)‡	1.82 (1.19-2.79)†	1.41 (0.99-2.02)	1.35 (0.86-2.12)
Recovery 5' (bpm)	2.15 (1.54-3.00)‡	2.03 (1.31-3.14)†	1.44 (1.01-2.06)*	1.54 (0.97-2.47)
Systolic blood pressure				
Peak (mmHg)	1.95 (1.47-2.50)‡	1.88 (1.32-2.68)‡	1.29 (0.97-1.71)	1.38 (0.95-2.02)
Response (mmHg)	1.83 (1.37-2.43)‡	1.81 (1.25-2.64)†	1.28 (0.93-1.77)	1.41 (0.93-2.15)
Recovery 2' (mmHg)	1.54 (1.17-2.03)†	1.68 (1.18-2.41)†	1.18 (0.87-1.60)	1.40 (0.94-2.08)

Recovery 5' (mmHg)	1.91 (1.49-2.45)‡	1.83 (1.31-2.55)‡	1.42 (1.07-1.89)*	1.50 (1.02-2.20)*
--------------------	-------------------	-------------------	-------------------	-------------------

Rate-pressure product

Peak (bpm·mmHg)	2.24 (1.61-3.13)‡	2.05 (1.33-3.15)†	1.41 (0.99-2.01)	1.49 (0.94-2.38)
-----------------	-------------------	-------------------	------------------	------------------

Response (bpm·mmHg)	2.37 (1.68-3.35)‡	2.10 (1.35-3.26)†	1.47 (1.00-2.17)	1.52 (0.93-2.49)
---------------------	-------------------	-------------------	------------------	------------------

Recovery 2' (bpm·mmHg)	2.19 (1.58-3.04)‡	2.12 (1.38-3.27)†	1.43 (1.00-2.07)	1.57 (0.99-2.51)
------------------------	-------------------	-------------------	------------------	------------------

Recovery 5' (bpm·mmHg)	2.55 (1.82-3.58)‡	2.34 (1.51-3.62)‡	1.66 (1.14-2.41)†	1.75 (1.08-2.85)*
------------------------	-------------------	-------------------	-------------------	-------------------

Adjusted for age, sex, Canadian Cardiovascular Society grading of angina pectoris, leisure-time physical activity, glycated

haemoglobin, left ventricular ejection fraction. * p<0.05, † p <0.01 and ‡ p<0.001.

Table 3. Predictive value of exercise test variables in CAD patients without diabetes (n=1089).

	Univariate		Multivariate	
	Cardiac death (n=31)	Sudden cardiac death (n=16)	Cardiac death (n=31)	Sudden cardiac death (n=16)
	Hazard ratio per +1 SD (95%CI)	Hazard ratio per +1 SD (95%CI)	Hazard ratio per +1 SD (95%CI)	Hazard ratio per +1 SD (95%CI)
Exercise capacity (%)	2.22 (1.47-3.33)‡	2.08 (1.18-3.70)*	1.92 (1.20-3.03)†	3.13 (0.87-3.13)
Heart rate				
Peak (%)	1.96 (1.33-2.86)‡	1.96 (1.18-3.33)†	1.47 (0.99-2.17)	1.67 (0.97-2.86)
Chronotropic response index (%)	2.13 (1.45-3.13)‡	2.08 (1.23-3.57)†	1.54 (1.02-2.33)*	1.79 (1.00-3.23)
Recovery 2' (bpm)	2.63 (1.96-3.57)‡	1.96 (1.25-3.13)†	2.17 (1.41-3.33)‡	1.75 (0.98-3.13)
Recovery 5' (bpm)	2.22 (1.54-3.23)‡	2.13 (1.28-3.45)†	1.52 (0.98-2.38)	1.89 (1.02-3.45)*
Systolic blood pressure				
Peak (mmHg)	1.67 (1.19-2.33)†	1.52 (0.94-2.44)	1.20 (0.83-1.75)	2.56 (0.75-2.08)
Response (mmHg)	1.47 (1.06-2.04)*	1.27 (0.79-2.00)	1.03 (0.71-1.49)	1.04 (0.62-1.75)
Recovery 2' (mmHg)	1.32 (0.92-1.85)	1.16 (0.71-1.92)	0.92 (0.61-1.37)	0.97 (0.57-1.64)

Recovery 5' (mmHg)	1.72 (1.25-2.38)*	1.73 (1.11-2.70)*	1.16 (0.79-1.69)	1.47 (0.88-2.44)
--------------------	-------------------	-------------------	------------------	------------------

Rate-pressure product

Peak (bpm·mmHg)	2.38 (1.59-3.57)‡	2.08 (1.20-3.57)†	1.56 (0.99-2.50)	1.72 (0.93-3.23)
-----------------	-------------------	-------------------	------------------	------------------

Response (bpm·mmHg)	2.44 (1.61-3.57)‡	2.08 (1.22-3.57)†	1.59 (0.99-2.56)	1.79 (0.93-3.33)
---------------------	-------------------	-------------------	------------------	------------------

Recovery 2' (bpm·mmHg)	2.22 (1.52-3.33)‡	1.67 (1.01-2.78)*	1.47 (0.92-2.38)	1.32 (0.71-2.44)
------------------------	-------------------	-------------------	------------------	------------------

Recovery 5' (bpm·mmHg)	2.33 (1.56-3.45)‡	2.17 (1.28-3.70)†	1.47 (0.93-2.33)	1.47 (0.93-2.33)
------------------------	-------------------	-------------------	------------------	------------------

Adjusted for age, sex, Canadian Cardiovascular Society grading of angina pectoris, leisure-time physical activity, glycated

haemoglobin, left ventricular ejection fraction. * $p < 0.05$, † $p < 0.01$ and ‡ $p < 0.001$.