

Leisure-time Physical Activity in Young Adults Born Preterm

Marjaana Tikanmäki, MD,^{1,2} Nina Kaseva MD, PhD,¹ Tuija Tammelin, PhD,³ Marika Sipola-Leppänen, MD, PhD,^{1,2,4} Hanna-Maria Matinolli, MHealthSci,^{1,2} Johan G Eriksson, MD, DMSc,^{1,5,6} Marjo-Riitta Järvelin, MD, PhD,^{7,8,9,10} Marja Vääräsmäki, MD, PhD^{4,11} and Eero Kajantie, MD, PhD.^{1,4,12}

Affiliations: ¹Chronic Disease Prevention Unit, Department of Health, National Institute for Health and Welfare, Oulu and Helsinki, Finland; ²Institute of Health Sciences, University of Oulu, Oulu, Finland; ³LIKES Research Centre for Physical Activity and Health, Jyväskylä, Finland; ⁴PEDEGO Research Unit (Research Unit for Pediatrics, Dermatology, Clinical Genetics, Obstetrics and Gynecology), Medical Research Center Oulu (MRC Oulu), Oulu University Hospital and University of Oulu, Oulu, Finland; ⁵Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Helsinki, Finland; ⁶Folkhälsan Research Center, Helsinki, Finland; ⁷Department of Epidemiology and Biostatistics, MRC–PHE Centre for Environment & Health, School of Public Health, Imperial College London, United Kingdom; ⁸Center for Life Course Epidemiology, Faculty of Medicine, University of Oulu, Oulu, Finland, ⁹Biocenter Oulu, Oulu, Finland; ¹⁰Unit of Primary Care, Oulu University Hospital, Oulu, Finland; ¹¹Children, Adolescents and Families Unit, Department of Welfare, National Institute for Health and Welfare, Oulu, Finland and ¹²Children's Hospital, Helsinki University Hospital, University of Helsinki, Helsinki, Finland.

Address correspondence to: Marjaana Tikanmäki, Department of Public Health Solutions, Chronic Disease Prevention Unit, National Institute for Health and Welfare, Kastelli Research Centre, PO Box 310, 90101 Oulu, Finland. [marjaana.tikanmaki@thl.fi], telephone +358-29-524 6549, fax +358-29-524 6222.

Key words: premature birth, physical activity, physical fitness

Short title: Physical Activity in Young Adults Born Preterm

Abbreviations: CI – confidence interval; GSD – Geometric standard deviation; ELBW – extremely low birth weight; EPT – early preterm; ESTER – ESTER Preterm Birth Study (Preterm Birth and Early Life Programming of Adult Health and Disease); FMBR – Finnish Medical Birth Register; LPT – late preterm, MET – metabolic equivalent; NFBC – Northern Finland Birth Cohort 1986; PA – physical activity; SGA – small for gestational age; SD – standard deviation; VLBW – very low birth weight.

Funding Sources: The ESTER study was supported by grants from the Academy of Finland (SALVE program for 2009–2012 and grants 127437, 129306, 130326, 134791 and 263924 [to EK]); the Doctoral Program for Public Health, University of Tampere (to MSL); the Emil Aaltonen Foundation (to EK), the European Commission (Framework 5 award QLG1-CT-2000-001643 to MRJ); the Finnish Foundation for Pediatric Research (to EK); the Finnish Government Special Subsidiary for Health Sciences (evo) (to JGE); Finska Läkaresällskapet (to JGE and NK); the Jalmari and Rauha Ahokas Foundation (to EK); the Juho Vainio Foundation (to EK, MT, and MV); the National Graduate School of Clinical Investigation (to MT); the Novo Nordisk Foundation (to EK and MV); the Signe and Ane Gyllenberg Foundation (to JGE, EK, and NK); the Sigrid Jusélius Foundation (to EK); and the Yrjö Jahnsson Foundation (to EK, MSL, and MV).

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflicts of Interest: The authors have no conflicts of interest to disclose.

Abstract

Objective: To evaluate the amount of self-reported physical activity in young adults born prematurely compared with those born at term.

Study Design: Unimpaired participants of the ESTER birth cohort study were studied at age 23.3 ± 1.2 (SD) years: 118 born early preterm (< 34 weeks), 210 late preterm (34 to 36 weeks), and 311 born at term (≥ 37 weeks, controls). The participants completed a validated 30-item 12-month physical activity questionnaire. The annual frequency and total volume of conditioning and non-conditioning leisure-time physical activity and commuting physical activity were calculated and analyzed the data by means of linear regression.

Results: Adults born early preterm reported 31.5% (95% CI, 17.4–43.2) lower volume of leisure-time physical activity (in metabolic equivalents, MET h/year) and had a 2.0-fold increased odds ratio (1.2–3.3) of being in the least active quintile than controls. Lower amounts of conditioning, non-conditioning, and commuting physical activity all contributed to the difference. In addition, early preterm participants undertook less vigorous physical activity (≥ 6 MET). No statistically significant differences in physical activity were found between the late preterm and control groups. Adjustments for potential early-life confounders and current mediating health characteristics did not change the results.

Conclusions: Young adults born early preterm engage less in leisure-time physical activities than peers born at term. This may in part underlie the increased risk factors of cardiometabolic and other non-communicable diseases in adults born preterm. Low physical activity is a risk factor of several non-communicable diseases and accessible to prevention.

Every year, approximately 14.9 million infants worldwide (11% of all newborns) are born preterm (< 37 weeks of gestation).¹ There is extensive evidence that preterm adults born at very low (VLBW; < 1500 g) or extremely low (ELBW; < 1000 g) birth weights have higher levels of risk factors for chronic non-communicable disease, including higher blood pressure, impaired glucose regulation, lower bone mineral density, and obstructive airflow.²⁻⁵ However, these individuals constitute only a small proportion of all preterm infants. Of all preterm infants in the United States, for example, 70% are born late preterm, between 34 and 36 weeks of gestation.⁶ Recent evidence suggests that many of these adverse consequences of preterm birth are present in those born late preterm and increase with the degree of prematurity.^{5, 7, 8}

Physical inactivity is related to increased levels of risk factors for non-communicable disease and this could in part explain them in those born preterm. Studies among children and adolescents born extremely preterm (≤ 28 weeks or ≤ 1000 g)^{9, 10} or with ELBW¹¹ or VLBW¹² suggest lower reported levels of physical activity compared with those born at term or of normal birth weight. However, some small studies among VLBW or preterm children have revealed no differences.^{13, 14} Adolescents and adults born preterm with VLBW or ELBW report substantially lower amounts of physical activity^{15, 16} and have lower levels of cardiorespiratory and muscular fitness.^{12, 17} We recently showed that lower fitness is also seen among the much larger group of early (< 34 weeks) and late preterm-born young adults.¹⁸ Whether these adults actually perform less physical activity is uncertain.

We studied self-reported physical activity in unimpaired young adults born at early or late preterm gestational ages. We hypothesized that preterm young adults would report less physical activity than young adults who were born at term. We also hypothesized that lower physical activity among those born prematurely would not be fully explained by conditions underlying preterm birth.

Methods

Participants

The ESTER (Preterm Birth and Early-Life Programming of Adult Health and Disease) Preterm Birth Study involves 1890 young adults. They were recruited through the Northern Finland Birth Cohort 1986 (NFBC; born in 1985–1986; 49.8%) and via the Finnish Medical Birth Register (FMBR; born in 1987–1989; 50.2%).⁸ In 2009–2011, 753 individuals with verified lengths of gestation participated in a clinical study at 23.3 ± 1.2 (SD) years of age.¹⁹ After exclusions (Figure 1, online), 118 participants born early preterm, 210 born late preterm, and 311 controls born at term (≥ 37 weeks) were unimpaired (no mental or physical disability), nonpregnant and had complete data on self-reported physical activity.

This study was approved by the Coordinating Ethics Committee at Helsinki and Uusimaa Hospital District. The participants provided a signed informed-consent document.

For the participants recruited through the NFBC 1986, perinatal data was collected previously.²⁰ Corresponding data from hospital and maternal welfare clinic records were obtained for those invited through the FMBR. Through these data, the length of gestation was confirmed (determined by ultrasonography in 62.7% and 53.1% of preterm infants and controls, respectively).^{8, 19} The study groups were defined by the length of gestation as early preterm (< 34 weeks), late preterm (34 to 36 weeks) and controls born at term (≥ 37 weeks). Diagnoses of maternal gestational diabetes, hypertension (gestational or chronic) or preeclampsia (including superimposed) were confirmed according to prevailing criteria.^{21, 22} Small for gestational age (SGA) was defined as birth weight more than 2 SD below the mean for gestational age.²³

The mean of three measurements was calculated for height. Body weight and composition were assessed using segmental multi-frequency bioelectrical impedance equipment (InBody 3.0, Biospace Co., Seoul, Korea). Medical history, medication, socioeconomic status and lifestyle data

were collected via questionnaires. Childhood socioeconomic status was assessed as the education level of the more highly educated parent.⁸

Self-reported Physical Activity

During the visit to the research clinic, the participants completed the modified Kuopio Ischemic Heart Disease Risk Factor Study questionnaire for detailed assessment of 12-month physical activity history.²⁴ The reproducibility and validity of the questionnaire have been confirmed.^{16, 25-28} The questionnaire comprises a 30-item list of types of physical activity, including conditioning leisure-time physical activity (20 items; e.g., running, skiing, swimming), non-conditioning leisure-time physical activity (8 items; e.g., household work, gardening, shoveling snow), physical activity from commuting to work (walking or cycling), and a category for other physical activity specified by the participant. The participants reported the monthly frequency and duration of each physical activity session for the previous 12 months and rated the average intensity of activities on a scale from 0 to 3 (0 = light, 1 = moderate, 2 = strenuous, 3 = very strenuous).

Data Analysis

The self-reported monthly frequency of physical activity was converted into times/year, and the average duration of each physical activity session was summed and converted into hours/year. The self-rated average physical activity intensities were converted into metabolic equivalents (METs) using the Compendium of Physical Activities.²⁹ An intensity of 1 MET corresponds to an energy expenditure of 1 kcal/kg/h, equivalent to the energy cost of sitting quietly. The total volume of physical activity in conditioning and non-conditioning leisure-time physical activity and commuting physical activity were calculated separately and summed as the total volume of leisure-time physical activity (MET h/year). The total amount of vigorous intensity physical activity (≥ 6 MET) was calculated per year in conditioning, non-conditioning, and commuting physical activity.

Statistical Methods

Characteristics were compared using Student's *t* test and the χ^2 test, with Yates' correction for continuity in 2 by 2 tables, and outcomes using linear and logistic regression analyses. The level of significance was set at a *p*-value of < .05. Interactions between two variables were tested (statistical significance level *p*-value < .01) by including a product term, using these variables. Non-normally distributed outcome variables, including zero values, were log-transformed ($\log_{10} [x+1]$) and mean differences reported as back-transformed percentages. Categorical covariates were entered as dummy variables with a separate dummy for missing values. Model 1 adjusted for sex, age, cohort, and season (December–February, March–May, June–August, September–November). Model 2 further adjusted for parental and early life factors: parental education, maternal smoking, gestational diabetes and hypertension, and birth weight SD score. Model 3 additionally adjusted for potential adult mediators: asthma,³⁰ height,^{15, 16} body fat percentage,³¹ and smoking.³² Analyses were rerun after replacement of the adjustment for body fat percentage with lean body mass.

Sensitivity analyses were performed to evaluate the effects of perinatal and neonatal factors on the associations. Analyses were rerun 1) after exclusion of participants born after a multiple pregnancy and 2) after exclusion of those born SGA. Further, among participants born early preterm, whether or not the outcomes could be predicted by multiple pregnancy or by supplementary oxygen treatment for more than 28 days after birth was assessed. Among all participants, prediction of outcomes by SGA status was also assessed. Analyses were performed using IBM SPSS Statistics for Windows, Version 22.0 (Armonk, NY).

Results

Gestational ages for the the early preterm, late preterm and control groups ranged from 23.9–33.9 weeks, 34–36.9 weeks and 37–43.1 weeks, respectively. The respective ranges of birth

weight were 655–3010 g, 1410–4440 g and 2310–4920 g. Participant characteristics are shown in Table I and outcomes by exposure group in Table II (online). There was no interaction between the association of sex and preterm birth with any outcomes.

The flow of participants through the study is shown in Figure 1 (online). A detailed nonparticipant analysis has been published.⁸ Among the participants in the overall study, the characteristics (listed in Table I) of those who completed the physical activity questionnaire were compared with the unimpaired nonparticipants who did not respond to the questionnaire or did not have valid questionnaire data (Figure 1, online). Similar proportions of the early and late preterm groups completed physical activity questionnaires compared with the control group. Among the late preterm group, the birth weight SD score was higher among physical activity questionnaire study participants than nonparticipants ($P = .04$), and nonparticipants were more often identified via the FMBR than the NFBC ($P = .03$). Participation in the physical activity questionnaire study was lower in the winter among the late preterm ($P = .01$) and control groups ($P = .02$) and higher in the summer among the late preterm group ($P = .01$).

The frequency and volume for the total and all types of leisure-time physical activity for women and men are detailed in Table III.

For the total amount of leisure-time physical activity, in the early preterm group, the overall frequency of physical activity was 17.2% lower and the overall total volume (MET h/wk) of physical activity was 31.5% lower compared with the control group in model 1 (adjusted for age, sex, cohort and season) (Table IV). The difference in frequency was no longer statistically significant when adjusted for socioeconomic status, pregnancy-related factors and adult characteristics in models 2 and 3, but the difference in the total volume of physical activity persisted (Figure 2, Table IV). Young adults in the early preterm group were two times more likely to be classified in the least active quintile of the population compared with young adults in the control group (odds ratio 2.0, 95% CI 1.2–3.3; model 1). The results remained similar after further

adjustments in models 2 and 3. The frequency and total volume of physical activity (Figure 2, Table IV), and classification in the least active quintile in late preterm individuals did not differ from those in the control individuals.

For conditioning leisure-time physical activity, in the early preterm group, the frequency was 31.3% lower and the total volume of conditioning physical activity was 46.6% lower than in the control group in model 1 (Table IV). These differences in frequency and total volume became slightly attenuated in models 2 and 3. The frequency and total amount of conditioning physical activity in the late preterm group did not differ from the control group.

For non-conditioning leisure-time physical activity, in the early preterm group, the total volume was 41.6% lower than in the control group in model 1 (Table IV). This difference persisted after statistical adjustments. The frequency of non-conditioning physical activity in early or late preterm groups was not different from that of the control group, and the volume of non-conditioning physical activity among late preterm individuals did not differ from that among the control individuals.

For commuting physical activity, the early preterm group reported a 46.3% lower frequency and 63.6% lower volume compared with the control group in model 1 (Table IV), but this difference was no longer statistically significant after further adjustment in models 2 and 3. The late preterm group did not differ from the control group in either frequency or volume of commuting physical activity.

For vigorous physical activity, in the early preterm group, the frequency was 43.5% lower on average and the total volume 54.3% lower than reported by the control group. Adjustments for covariates in models 2 and 3 did not change the results. The late preterm group undertook similar amounts of vigorous physical activity as the control group.

The total volume of physical activity was higher among women ($p = .02$) (Table III), older participants ($p = .02$), non-smokers ($p = .02$), and those with a lower body fat percentage ($p = .01$).

Associations with smoking were mostly in connection with conditioning and commuting physical activity ($p < .001$), and associations with lower body fat percentage with conditioning physical activity ($p = .01$). Vigorous physical activity was more common among the offspring of higher-educated parents ($p = .001$) and reported especially during winter ($p = .001$), and conditioning physical activity was reported more frequently during winter ($p = .03$) and spring ($p = .04$), while other outcomes did not differ according to season. Participants exposed to maternal gestational hypertension or chronic hypertension during pregnancy performed less vigorous physical activity ($p = .02$) (data not shown).

For the sensitivity analysis, when only singletons were included in the analyses, the differences in total, non-conditioning, and vigorous physical activity persisted after all adjustments, as did differences in conditioning physical activity in model 1, while the differences in commuting physical activity decreased to statistical non-significance. Multiple pregnancy was not associated with physical activity outcomes among the early preterm group.

Early preterm-born participants who had received supplementary oxygen for longer than 28 days after birth ($n = 17$) showed a lower frequency of total ($p = .01$) and conditioning ($p = .01$) physical activity than the other participants born early preterm. Being born SGA was not associated with physical activity outcomes among the participants, and the results were similar when those born SGA were excluded.

When adjustments were made for lean body mass instead of body fat percentage, the results were similar in all analyses.

Discussion

The main finding of this study is that unimpaired young adults born early preterm performed markedly less leisure-time physical activity than those born at term. Lower levels were seen for conditioning and non-conditioning, commuting and vigorous physical activity and resulted in a

31.5% lower total volume of physical activity. This difference was not explained by socioeconomic status or by pregnancy-related factors underlying preterm birth. The finding persisted when adjusted for adult characteristics including body size, current smoking or diagnosed asthma, suggesting that it is not mediated through these characteristics.

Previous studies focused on the smallest preterm infants, who as adolescents or adults, have shown substantially less physical activity than those born at term. Unimpaired 17-year-old survivors with birth weights ≤ 800 g have reported less frequent participation in sports.¹² Similarly, unimpaired VLBW (<1500 g) adults reported exercising less frequently, with lower intensity and shorter sessions, and performed 48.6% less conditioning leisure-time physical activity compared with those born at term.^{15, 16} This parallels our finding of 46.6% less conditioning physical activity among those born early preterm. Although the difference we found among the late preterm group was not statistically significant, the point estimate for conditioning physical activity was 15.1% less compared with those born at term, consistent with a dose-response relationship between shorter length of gestation and less physical activity.

In previous studies among ELBW or VLBW adults, lower physical activity was specifically related to less conditioning leisure-time physical activity or sports participation.^{12, 15, 16} By contrast, in the present study, low-level physical activity was observed for conditioning, non-conditioning and commuting physical activity, and also when vigorous intensity physical activity was assessed separately. The reasons for this difference remain unclear. However, our present findings suggest a general preference for a physically less active lifestyle rather than a specific aversion to conditioning physical activity sports. Adults born preterm not only undertook less conditioning and vigorous intensity activities but also reported less commuting and non-conditioning types of activity. The reported differences could not be attributed to pregnancy conditions underlying preterm birth, or to fetal growth. This suggests that mostly the observed differences in physical activity are due to postnatal events or prematurity itself rather than perinatal conditions.

Of note, findings of lower self-reported physical activity reported in this study and in prior studies have not been replicated in adults when physical activity is assessed objectively by means of accelerometry.³³ This was also the case in a subset of the present cohort who underwent accelerometry.³⁴ Objective measurements during childhood have not been able to capture differences in physical activity between those born preterm and at term,^{35, 36} except in one study of 7-year-old boys born very preterm (<32 weeks).³⁷ One reason for this discrepancy may be that self-reporting and accelerometry capture different aspects of physical activity. Self-reporting enables the assessment of a broad range of physical activities in any circumstances and provides average figures within a longer time frame, whereas objective measurement is more precise in registering the intensity of physical activity and the amount of sedentary time within a shorter period.³⁸ Correlation coefficients between self-reported and objectively measured physical activity are usually low-to-moderate, in our cohort 0.25³⁴, and according to meta-analyses approximately 0.3–0.4.^{38, 40} Additionally, in a relatively large sample with self-reported physical activity, those who are least active are well-represented, whereas physically more active participants may volunteer to participate in accelerometry.

Potential explanations for lower self-reported levels of leisure-time physical activity among preterm-born individuals include reduced muscle mass due to preterm birth,⁴¹ poorer motor skills persisting through childhood and adolescence⁴² and lower self-efficacy, leading to reduced muscular fitness in young adulthood,¹⁸ and this, combined with poorer visual acuity⁴³ and reduced pulmonary function, could discourage undertaking conditioning, non-conditioning, and commuting physical activity.⁴ Children and adults born prematurely have reduced cognitive abilities^{44, 45} and although the direction of causality has been debated, it has been shown that those with lower cognitive abilities undertake less physical activity.^{46, 47} A previous study in men showed that the association between low exercise capacity and low cognitive function was particularly strong among those born preterm,⁴⁵ suggesting that adults born preterm who have lower cognitive abilities

could be a particular risk group. We had no comprehensive assessment of cognitive abilities and could not assess this suggestion. Another explanation could be more protective parenting, which we reported in a previous study,⁴⁸ where we compared adults born preterm with VLBW with their term-born controls. However, a recent meta-analysis⁴⁸ did not reveal any systematic difference in maternal parenting behavior.⁴⁹

Low-level leisure-time physical activity among individuals born early preterm may predispose them to cardiometabolic diseases and their risk factors, including elevated blood pressure, impaired glucose regulation,^{2, 5, 8} and reduced muscular fitness.¹⁸ Low-level physical activity is also associated with other non-communicable diseases,⁵⁰ and it predicts mortality in late adulthood.⁵¹ According to current U.S. Department of Health and Human Services⁵² and World Health Organization⁵³ recommendations, 150 minutes of moderate intensity (3–6 MET) physical activity or 75 minutes of vigorous intensity (> 6 MET) physical activity per week provides substantial health benefits. For comparison, the 30% (approximately 8.8 MET h/week at mean levels) less total amount of physical activity found among young adults born early preterm corresponds to the equivalent of more than 120 minutes of brisk walking (at an intensity of 4.3 MET) per week.

The strengths of this study include assessment of self-reported 12-month leisure-time physical activity using a validated detailed questionnaire.^{25, 27} A limitation of the method is the potential for recall bias that could differ between groups. Potentially lower physical self-efficacy among those born preterm⁵⁴ may affect self-rating of physical activity. However, individuals born preterm also tend to respond to questionnaires in a socially more acceptable manner,⁵⁵ which could result in reporting of higher levels of physical activity. This would diminish rather than exaggerate the differences. The distributions of the subcomponents of physical activity, particularly commuting physical activity, were skewed even after log-transformation. However, the distribution of total physical activity, on which the main conclusions are based, was less skewed. This study population

enabled the evaluation of leisure-time physical activity across the full range of preterm birth. We had access to reliable and diverse perinatal data, including verified length of gestation. Most (86.0%) of the participants in the clinical examination phase of the ESTER study also responded to the physical activity questionnaire (Figure 1). Detailed nonparticipation analyses presented in a previous publication⁸ and the present paper raised little concern about participation bias. Also, residual confounding cannot be excluded.

The lack of physical activity noted in this study in preterm adults may contribute to a higher risk of chronic non-communicable diseases in later life and offers a target for prevention. Healthcare professionals should be encouraged to actively support preterm-born children and their families in finding physical activities suitable for each individual. Regular participation in various forms of physical activity and avoidance of a physically inactive lifestyle in childhood and adolescence are effective in decreasing risk factors of several chronic non-communicable diseases in a life-course perspective.⁵⁰ Accordingly, participation in physical activities should be strongly encouraged by pediatricians, other health professionals, and parents.

Acknowledgements

We are grateful to all the participants for their contribution in the ESTER Study. We also thank research nurses Katriina Inget, Sinikka Kursu, Hanna-Maria Matinolli, Liisa Saarikoski, Marjatta Takala, Sarianna Vaara, data manager Sigrid Rosten and research assistants Risto Karvonen, Petteri Koivurova, Antti Koskela, Sanna Mustaniemi, Heli-Kaisa Saarenpää and Marja Vanhatalo from the National Institute of Health and Welfare, Oulu and Helsinki, Finland.

References

1. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller A-B, Narwal R, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: A systematic analysis and implications. *Lancet*. 2012;379:2162-72.
2. Hovi P, Vohr B, Ment LR, Doyle LW, McGarvey L, Morrison KM, et al. Blood pressure in young adults born at very low birth weight. *Hypertension*. 2016;68:880-7.
3. Parkinson JRC, Hyde MJ, Gale C, Santhakumaran S, Modi N. Preterm birth and the metabolic syndrome in adult life: A systematic review and meta-analysis. *Pediatrics*. 2013;131:e1240-e63.
4. Saarenpää HK, Tikanmäki M, Sipola-Leppänen M, Hovi P, Wehkalampi K, Siltanen M, et al. Lung function in very low birth weight adults. *Pediatrics*. 2015;136:642-50.
5. Kajantie E, Hovi P. Is very preterm birth a risk factor for adult cardiometabolic disease? *Semin Fetal Neonatal Med*. 2014;19:112-7.
6. Martin JA, Osterman MJ. Preterm births - united states, 2006 and 2010. *MMWR Surveill Summ*. 2013;62:136-8.
7. Johansson S, Iliadou A, Bergvall N, Tuvemo T, Norman M, Cnattingius S. Risk of high blood pressure among young men increases with the degree of immaturity at birth. *Circulation*. 2005;112:3430-6.
8. Sipola-Leppänen M, Väärasmäki M, Tikanmäki M, Matinolli HM, Miettola S, Hovi P, et al. Cardiometabolic risk factors in young adults who were born preterm. *Am J Epidemiol*. 2015; :861-73.
9. Clemm H, Roksund O, Thorsen E, Eide GE, Markestad T, Halvorsen T. Aerobic capacity and exercise performance in young people born extremely preterm. *Pediatrics*. 2011;129:e97-e105.
10. Clemm HH, Vollaeter M, Roksund OD, Markestad T, Halvorsen T. Adolescents who were born extremely preterm demonstrate modest decreases in exercise capacity. *Acta Paediatr*. 2015;104:1174-81.
11. Kilbride HW, Gelatt MC, Sabath RJ. Pulmonary function and exercise capacity for ELBW survivors in preadolescence: Effect of neonatal chronic lung disease. *J Pediatr*. 2003;143:488-93.

12. Rogers M, Fay TB, Whitfield MF, Tomlinson J, Grunau RE. Aerobic capacity, strength, flexibility, and activity level in unimpaired extremely low birth weight (≤ 800 g) survivors at 17 years of age compared with term-born control subjects. *Pediatrics*. 2005;116:e58-65.
13. Winck AD, Heinzmann-Filho JP, Schumann D, Zatti H, Mattiello R, Jones MH, et al. Growth, lung function, and physical activity in schoolchildren who were very-low-birth-weight preterm infants. *J Bras Pneumol*. 2016;42:254-60.
14. Joshi S, Powell T, Watkins WJ, Drayton M, Williams EM, Kotecha S. Exercise-induced bronchoconstriction in school-aged children who had chronic lung disease in infancy. *J Pediatr*. 2013;162:813-8.e1.
15. Kajantie E, Strang-Karlsson S, Hovi P, Räikkönen K, Pesonen AK, Heinonen K, et al. Adults born at very low birth weight exercise less than their peers born at term. *J Pediatr*. 2010;157:610-6, 6 e1.
16. Kaseva N, Wehkalampi K, Strang-Karlsson S, Salonen M, Pesonen AK, Räikkönen K, et al. Lower conditioning leisure-time physical activity in young adults born preterm at very low birth weight. *PLoS One*. 2012;7:e32430.
17. Burns YR, Danks M, O'Callaghan MJ, Gray PH, Cooper D, Poulsen L, et al. Motor coordination difficulties and physical fitness of extremely-low-birthweight children. *Dev Med Child Neurol*. 2009;51:136-42.
18. Tikanmäki M, Tammelin T, Sipola-Leppänen M, Kaseva N, Matinolli HM, Miettola S, et al. Physical fitness in young adults born preterm. *Pediatrics*. 2016;137:1-10.
19. Sipola-Leppänen M, Väärasmäki M, Tikanmäki M, Hovi P, Miettola S, Ruokonen A, et al. Cardiovascular risk factors in adolescents born preterm. *Pediatrics*. 2014;134:e1072-81.
20. Järvelin MR, Hartikainen-Sorri AL, Rantakallio P. Labour induction policy in hospitals of different levels of specialisation. *Br J Obstet Gynaecol*. 1993;100:310-5.
21. Miettola S, Hartikainen AL, Väärasmäki M, Bloigu A, Ruokonen A, Järvelin MR, et al. Offspring's blood pressure and metabolic phenotype after exposure to gestational hypertension in utero. *Eur J Epidemiol*. 2013;28:87-98.

22. Vääräsmäki M, Pouta A, Elliot P, Tapanainen P, Sovio U, Ruukonen A, et al. Adolescent manifestations of metabolic syndrome among children born to women with gestational diabetes in a general-population birth cohort. *Am J Epidemiol.* 2009;169:1209-15.
23. Pihkala J, Hakala T, Voutilainen P, Raivio K. [Characteristic of recent fetal growth curves in Finland]. *Duodecim.* 1989;105:1540-6.
24. Borodullin K. Physical activity, fitness, abdominal obesity, and cardiovascular risk factors in Finnish men and women - the National FINRISK 2002 study. Helsinki: the National Public Health Institute; 2006.
25. Jacobs DR, Jr., Ainsworth BE, Hartman TJ, Leon AS. A simultaneous evaluation of 10 commonly used physical activity questionnaires. *Med Sci Sports Exerc.* 1993;25:81-91.
26. De Backer G, Kornitzer M, Sobolski J, Dramaix M, Degre S, de Marneffe M, et al. Physical activity and physical fitness levels of Belgian males aged 40-55 years. *Cardiology.* 1981;67:110-28.
27. Lakka TA, Salonen JT. Intra-person variability of various physical activity assessments in the Kuopio Ischaemic Heart Disease Risk Factor Study. *Int J Epidemiol.* 1992;21:467-72.
28. Lakka TA, Salonen JT. Moderate to high intensity conditioning leisure time physical activity and high cardiorespiratory fitness are associated with reduced plasma fibrinogen in Eastern Finnish men. *J Clin Epidemiol.* 1993;46:1119-27.
29. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Jr., Tudor-Locke C, et al. 2011 compendium of physical activities: A second update of codes and MET values. *Med Sci Sports Exerc.* 2011;43:1575-81.
30. Jaakkola JJ, Ahmed P, Ieromnimon A, Goepfert P, Laiou E, Quansah R, et al. Preterm delivery and asthma: A systematic review and meta-analysis. *J Allergy Clin Immunol.* 2006;118:823-30.
31. Sipola-Leppänen M, Vääräsmäki M, Tikanmäki M, Matinolli H-M, Miettola S, Hovi P, et al. Cardiometabolic risk factors in young adults who were born preterm. *Am J Epidemiol.* 2015;181:861-73.

32. Strang-Karlsson S, Räikkönen K, Pesonen A-K, Kajantie E, Paavonen E, Lahti J, et al. Very low birth weight and behavioral symptoms of attention deficit hyperactivity disorder in young adulthood: The Helsinki Study of very-low-birth-weight adults. *Am J Psychiatry*. 2008;165:1345-53.
33. Kaseva N, Martikainen S, Tammelin T, Hovi P, Järvenpää A-L, Andersson S, et al. Objectively measured physical activity in young adults born preterm at very low birth weight. *J Pediatr*. 2015;166:474-6.
34. Tikanmäki M, Tammelin T, Kaseva N, Sipola-Leppänen M, Matinolli H-M, Hakonen H, et al. Objectively measured physical activity and sedentary time in young adults born preterm-the ESTER study. *Pediatr Res*. 2017.
35. Lowe J, Watkins WJ, Kotecha SJ, Edwards MO, Henderson AJ, Kotecha S. Physical activity in school-age children born preterm. *J Pediatr*. 2015;166:877-83.
36. Welsh L, Kirkby J, Lum S, Odendaal D, Marlow N, Derrick G, et al. The EPICURE study: Maximal exercise and physical activity in school children born extremely preterm. *Thorax*. 2010;65:165-72.
37. Lowe J, Watkins WJ, Kotecha SJ, Kotecha S. Physical activity and sedentary behavior in preterm-born 7-year old children. *PLoS One*. 2016;11:e0155229.
38. Steene-Johannessen J, Anderssen SA, van der Ploeg HP, Hendriksen IJ, Donnelly AE, Brage S, et al. Are self-report measures able to define individuals as physically active or inactive? *Med Sci Sports Exerc*. 2016;48:235-44.
39. Prince SA, Adamo KB, Hamel ME, Hardt J, Connor Gorber S, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: A systematic review. *Int J Behav Nutr Phys Act*. 2008;5:56.
40. Sofi F, Capalbo A, Cesari F, Abbate R, Gensini GF. Physical activity during leisure time and primary prevention of coronary heart disease: An updated meta-analysis of cohort studies. *Eur J Cardiovasc Prev Rehabil*. 2008;15:247-57.
41. Brown LD. Endocrine regulation of fetal skeletal muscle growth: Impact on future metabolic health. *J Endocrinol*. 2014;221:R13-R29.

42. de Kieviet JF, Piek JP, Aarnoudse-Moens CS, Oosterlaan J. Motor development in very preterm and very low-birth-weight children from birth to adolescence: A meta-analysis. *JAMA*. 2009;302:2235-42.
43. Evensen KA, Lindqvist S, Indredavik MS, Skranes J, Brubakk AM, Vik T. Do visual impairments affect risk of motor problems in preterm and term low birth weight adolescents? *Eur J Paediatr Neurol*. 2009;13:47-56.
44. Aarnoudse-Moens CSH, Weisglas-Kuperus N, van Goudoever JB, Oosterlaan J. Meta-analysis of neurobehavioral outcomes in very preterm and/or very low birth weight children. *Pediatrics*. 2009;124:717.
45. Svedenkrans J, Kowalski J, Norman M, Bohlin K. Low exercise capacity increases the risk of low cognitive function in healthy young men born preterm: A population-based cohort study. *PLoS One*. 2016;11:e0161314.
46. Cox EP, O'Dwyer N, Cook R, Vetter M, Cheng HL, Rooney K, et al. Relationship between physical activity and cognitive function in apparently healthy young to middle-aged adults: A systematic review. *J Sci Med Sport*. 2016;19:616-28.
47. Fedewa AL, Ahn S. The effects of physical activity and physical fitness on children's achievement and cognitive outcomes. *Res Q Exerc Sport*. 2011;82:521-35.
48. Pyhälä R, Räikkönen K, Pesonen AK, Heinonen K, Lahti J, Hovi P, et al. Parental bonding after preterm birth: Child and parent perspectives in the helsinki study of very low birth weight adults. *J Pediatr*. 2011;158:251-6 e1.
49. Bilgin A, Wolke D. Maternal sensitivity in parenting preterm children: A meta-analysis. *Pediatrics*. 2015;136:e177.
50. Physical activity guidelines advisory committee report, 2008. Physical Activity Guidelines Advisory Committee. Department of Health and Human Services, Washington, DC: U.S. 2008.
51. Nocon M, Hiemann T, Müller-Riemenschneider F, Thalau F, Roll S, Willich SN. Association of physical activity with all-cause and cardiovascular mortality: A systematic review and meta-analysis. *Eur J Cardiovasc Prev Rehabil*. 2008;15:239-46.

52. Physical activity guidelines for Americans. Be active, healthy, and happy! ODPHP publication no. U0036 October 2008. US Department of Health and Human Services. U.S. Department of Health and Human Services, Washington, DC: U.S.2008.
53. WHO. Global recommendations on physical activity for health. World Health Organization. Geneva 2010. p. 58.
54. Saigal S, Stoskopf B, Boyle M, Paneth N, Pinelli J, Streiner D, et al. Comparison of current health, functional limitations, and health care use of young adults who were born with extremely low birth weight and normal birth weight. *Pediatrics*. 2007;119:e562-73.
55. Allin M, Rooney M, Cuddy M, Wyatt J, Walshe M, Rifkin L, et al. Personality in young adults who are born preterm. *Pediatrics*. 2006;117:309-16.

Figure Legends:

Figure 1. Flow chart of the study population. EPT, early preterm (< 34 gestational weeks); LPT, late preterm (34–36 gestational weeks)

Figure 2. Mean differences in physical activity (95% CIs, error bars) consisting of conditioning and non-conditioning leisure-time physical activity and commuting physical activity in preterm groups compared with controls (zero line). Model 1 was adjusted for sex, age, cohort and season. Model 2 was adjusted as for model 1 plus socioeconomic status, maternal smoking, gestational diabetes and hypertensive disorder during pregnancy, and birth weight SD score. Model 3 was adjusted as for model 2 plus diagnosed asthma, adult body size (height, body fat percentage) and smoking.

Table I. Perinatal, neonatal, and current characteristics of the young adults born preterm and the controls born at term.

	Early preterm (n = 118)			<i>P</i> ^a	Late preterm (n = 210)			<i>P</i> ^a	Controls (n = 311)			Missing (EPT/LPT / Controls)
	n	%	Mean (SD)		n	%	Mean (SD)		n	%	Mean (SD)	
Men	57	48.3		1.00	104	49.5		.73	148	47.6		0/0/0
NFBC member	44	37.3		<.001	100	47.6		.003	190	61.1		0/0/0
Peri- and neonatal												
Multiple pregnancy	30	25.4		<.001	29	13.8		<.001	3	1.0		0/0/0
Maternal hypertension ^b	17	14.4		.27	34	16.2		.08	33	10.6		0/2/2
Maternal pre-eclampsia ^c	28	23.7		<.001	24	11.4		.005	14	4.5		0/2/2
Maternal gestational diabetes ^d	4	3.4		.59	11	5.2		.07	6	1.9		21/24/8
Maternal smoking during pregnancy	17	14.4		.87	41	19.5		.40	51	16.4		7/5/4
Cesarean section	73	61.9		<.001	58	27.6		<.001	36	11.6		0/0/0
Gestational age, weeks			31.9 (1.9)	<.001			35.8 (0.8)	<.001			40.1 (1.2)	0/0/0
Birth weight, g			1786 (478)	<.001			2692 (527)	<.001			3576 (482)	0/0/0
Birth weight SD score, SD			-0.8 (1.4)	<.001			-0.6 (1.3)	<.001			0.0 (1.0)	0/0/0
SGA	20	16.9		<.001	25	11.9		<.001	5	1.6		0/0/0
Supplementary oxygen	85	72.0		<.001	78	37.1		<.001	6	1.9		0/0/0
Duration of supplementary oxygen, days			16.1 (12.9)	<.001			16.4 (69.6)	0.655			3.5 (1.6)	0/0/0 ^e
Current												
Age, y			23.1 (1.4)	.003			23.2 (1.2)	.001			23.5 (1.1)	0/0/0
Height, cm												0/0/0
Men			179 (7)	.36			177 (7)	.79			178 (7)	
Women			163 (5)	.45			165 (6)	.46			164 (6)	
BMI, kg/m ²												0/0/0
Men			24.4 (4.2)	.83			25.5 (4.8)	.03			24.3 (3.3)	
Women			24.4 (5.5)	.15			23.5 (4.1)	.62			23.2 (4.3)	
Body fat percentage, %												0/0/2
Men			17.9 (6.6)	.58			19.7 (7.7)	.01			17.4 (5.8)	
Women			30.9 (7.6)	.006			28.8 (6.9)	.26			27.8 (7.3)	
Lean body mass, kg												0/0/2
Men			63.5 (9.0)	.77			63.6 (8.6)	.65			63.1 (8.5)	
Women			44.1 (7.1)	.71			44.7 (5.7)	.66			44.4 (5.3)	
Parental education				.70				.57				
Basic or less	10	8.5			16	7.8			18	5.8		0/5/2
Secondary	71	60.2			115	56.1			189	61.0		

Lower-level tertiary	13	11.0		27	12.9		42	13.5	
Upper-level tertiary	24	20.3		47	22.4		60	19.3	
Daily smoking	32	27.1	.31	49	22.9	.77	68	21.9	0/0/0
Asthma	25	21.2	.10	34	16.2	.61	44	14.1	0/0/0
Season									0/0/0
Winter	27	22.9	.49	44	21.0	.72	60	19.3	
Spring	29	24.6	1.00	68	32.4	.05	75	24.1	
Summer	17	14.4	.19	36	17.1	.39	64	20.8	
Autumn	45	38.1	.77	62	29.5	.15	112	36.0	

Early preterm (EPT; < 34 weeks); late preterm (LPT; 34–36 weeks); controls (≥ 37 weeks); NFBC indicates Northern Finland Birth Cohort 1986; SGA, small for gestational age.

^a*P* values refer to comparisons between preterm groups and controls with the use of Student's *t* test or Pearson's χ^2 test with Yates' continuity correction for 2 by 2 tables. *P* values < 0.05 were considered statistically significant.

^bGestational or chronic hypertension.

^cIncludes superimposed preeclampsia.

^dParticipants whose mothers' gestational diabetes data are missing include those whose mothers did not undergo an oral glucose tolerance test despite risk factors and thus have uncertain gestational diabetes status.

^eOf those who received supplementary oxygen.

Table II. Geometric mean (and geometric SD) values of outcomes in early-preterm (< 34 weeks), late-preterm (34 to 36 weeks) and control (≥ 37 weeks) groups by sex.

		Early preterm (n = 118)	Late preterm (n = 210)	Controls (n = 311)
		Geometric mean and SD	Geometric mean and SD	Geometric mean and SD
Total amount of leisure-time physical activity				
Frequency, times/wk				
Men		5.6 (2.4)*	7.3 (2.0)	7.5 (2.0)
Women		8.1 (2.0)	8.4 (1.9)	8.4 (1.8)
All		6.8 (2.2)	7.8 (2.0)	8.0 (1.9)
Total volume, MET h/wk ^a				
Men		18.1 (3.4)**	28.9 (2.3)	30.8 (2.5)
Women		25.3 (2.5)	29.0 (2.2)	30.1 (2.0)
All		21.5 (2.9)**	28.9 (2.3)	30.4 (2.2)
Conditioning leisure-time physical activity				
Frequency, times/wk				
Men		1.7 (4.0)*	2.3 (3.9)	2.6 (2.7)
Women		2.0 (3.8)	2.7 (3.1)	2.6 (2.9)
All		1.8 (3.9)*	2.5 (3.5)	2.6 (2.8)
Total volume, MET h/wk ^a				
Men		7.0 (8.7)*	11.8 (8.7)	15.2 (4.3)
Women		8.7 (7.1)	13.0 (5.9)	13.1 (4.0)
All		7.8 (7.8)**	12.4 (7.2)	14.0 (4.1)
Non-conditioning leisure-time physical activity				
Frequency, times/wk				

	Men	0.9 (4.4)	1.3 (3.1)	1.3 (3.7)
	Women	1.8 (2.3)	1.7 (2.2)	1.7 (2.2)
	All	1.3 (3.5)	1.5 (2.6)	1.5 (2.9)
Total volume, MET h/wk ^a				
	Men	1.5 (19.3)*	3.0 (8.5)	3.6 (10.6)
	Women	4.8 (3.0)	5.5 (2.6)	5.7 (3.0)
	All	2.7 (9.7)*	4.1 (5.3)	4.6 (6.2)
Commuting physical activity				
Frequency, times/wk				
	Men	0.3 (14.1)*	0.7 (12.0)	0.9 (9.3)
	Women	1.2 (8.7)	1.2 (8.2)	1.4 (7.1)
	All	0.7 (11.9)	0.9 (10.1)	1.1 (8.2)
Total volume, MET h/wk ^a				
	Men	0.1 (108.5)*	0.2 (78.7)	0.3 (50.4)
	Women	0.7 (47.0)	0.7 (38.9)	0.9 (30.2)
	All	0.2 (83.0)	0.4 (57.9)	0.6 (41.7)
Vigorous ^b leisure-time physical activity				
Frequency, times/wk				
	Men	0.6 (5.0)**	1.2 (3.9)	1.2 (3.3)
	Women	0.9 (5.1)	1.3 (4.5)	1.3 (3.5)
	All	0.7 (5.1)**	1.2 (4.2)	1.3 (3.4)
Total volume, MET h/wk ^a				
	Men	3.7 (8.2)**	9.1 (5.3)	9.6 (4.7)
	Women	4.6 (7.9)	7.5 (6.5)	8.0 (4.6)
	All	4.1(8.0)***	8.3 (5.9)	8.7 (4.6)

The geometric mean is the n th root of the product of n individual values. Geometric standard deviations correspond to the percent increase in the variable corresponding to a change of one standard-deviation unit in the logarithm of the variable.^aMET h/wk indicates MET hours of physical activity per week (MET indicates metabolic equivalent).

^bPhysical activity intensity ≥ 6 MET.

Levels of statistical significance for differences between the preterm group and the control group in t tests: *** <0.001 , ** <0.01 , * <0.05 .

Table III. Physical activity in women and men presented as geometric mean (geometric standard deviation) early preterm, late preterm and term control groups combined.

	Women (n=330)	Men (n=309)
	Geometric mean and SD	Geometric mean and SD
Total amount of leisure-time physical activity		
Frequency, times/wk	8.3 (1.9)	7.0 (2.1)
Total volume, MET h/wk ^a	28.8 (2.2)	27.3 (2.6)
Conditioning leisure-time physical activity		
Frequency, times/wk	2.5 (3.2)	2.3 (3.4)
Volume, MET h/wk ^a	12.1 (5.2)	12.1 (6.5)
Non-conditioning leisure-time physical activity		
Frequency, times/week	1.7 (2.2)	1.2 (3.6)
Volume, MET h/wk ^a	5.4 (2.9)	2.9 (11.3)
Commuting physical activity		
Frequency, times/wk	1.3 (7.7)	0.7 (11.1)
Volume, MET h/wk ^a	0.8 (35.4)	0.2 (69.9)
Vigorous^b leisure-time physical activity		
Frequency, times/wk	1.2 (4.1)	1.1 (3.9)
Volume, MET h/wk ^a	7.1 (5.7)	7.9 (5.8)

The geometric mean is the nth root of the product of n individual values. Geometric standard deviations correspond to the percent increase in the variable corresponding to a change of one standard-deviation unit in the logarithm of the variable.^aMET h/wk indicates MET hours of physical activity per week (MET indicates metabolic equivalent).

^bPhysical activity intensity ≥ 6 MET.

Table IV. Outcomes in preterm groups compared with term-born controls.

	Controls (n = 311)	Model ^a	Early preterm (n = 118)	<i>P</i> ^b	Late preterm (n = 210)	<i>P</i> ^b
	Geometric mean and SD		Mean difference % (95% CI)		Mean difference % (95% CI)	
Total amount of leisure-time physical activity						
Frequency, times/wk	8.0 (1.9)					
		1	-17.2 (-28.5; -4.2)	.01	-3.0 (-14.0; 9.4)	.62
		2	-14.4 (-29.9; 0.3)	.06	-1.5 (-13.2; 11.7)	.81
		3	-13.0 (-25.7; 2.0)	.09	-0.9 (-12.6; 12.4)	.89
Total volume, MET h/wk ^c	30.4 (2.2)					
		1	-31.5 (-43.2; -17.4)	<.001	-6.5 (-19.8; 9.1)	.39
		2	-30.1 (-42.9; -14.5)	.001	-4.0 (-18.2; 12.8)	.62
		3	-28.4 (-41.6; -12.3)	.001	-2.3 (-16.8; 14.8)	.78
Conditioning leisure-time physical activity						
Frequency, times/wk	2.6 (2.8)					
		1	-31.3 (-46.7; -11.5)	.004	-8.2 (-25.5; 13.1)	.42
		2	-27.0 (-44.5; -4.1)	.02	-4.3 (-23.0; 19.0)	.69
		3	-24.6 (-42.4; -1.3)	.04	-2.9 (-21.6; 20.2)	.78
Total volume, MET h/wk ^c	14.0 (4.1)					
		1	-46.6 (-63.3; -22.2)	.001	-15.1 (-37.7; 15.7)	.30
		2	-42.6 (-61.8; -13.8)	.008	-9.6 (-34.6; 24.8)	.54
		3	-39.8 (-59.8; -9.8)	.01	-7.3 (-32.8; 27.8)	.64
Non-conditioning leisure-time physical activity						
Frequency, times/wk	1.5 (2.9)					
		1	-17.2 (-34.2; 4.1)	.11	-2.9 (-19.6; 17.3)	.76
		2	-20.4 (-37.9; 2.2)	.07	-1.7 (-19.4; 19.8)	.86
		3	-21.5 (-39.0; 1.1)	.06	-1.7 (-19.6; 20.1)	.86
Total volume, MET h/wk ^c	4.6 (6.2)					
		1	-41.6 (-60.9; -13.0)	.008	-8.7 (-34.2; 26.8)	.66
		2	-46.8 (-65.5; -18.1)	.004	-7.3 (-34.2; 30.6)	.66
		3	-48.2 (-66.6; -19.8)	.003	-7.7 (-34.8; 30.7)	.65
Commuting physical activity						
Frequency, times/wk	1.1 (8.2)					
		1	-46.3 (-66.6; -13.5)	.01	-17.5 (-44.5; 22.7)	.34
		2	-35.5 (-61.4; 7.9)	.44	-15.1 (-43.9; 28.3)	.44
		3	-28.0 (-57.3; 21.2)	.22	-14.9 (-43.7; 28.6)	.44
Total volume, MET h/wk ^c	0.6 (41.7)					
		1	-63.6 (-84.3; -15.4)	.02	-31.9 (-66.0; 36.6)	.28
		2	-48.3 (-79.3; 28.8)	.16	-27.6 (-64.9; 49.6)	.38
		3	-42.0 (-76.8; 44.8)	.24	-24.8 (-63.6; 55.4)	.44
Vigorous^d leisure-time physical activity						
Frequency, times/wk	1.3 (3.4)					
		1	-43.5 (-58.0; -24.1)	<.001	-6.2 (-26.5; 19.4)	.60
		2	-42.6 (-58.2; -21.3)	.001	-5.7 (-26.6; 21.2)	.65
		3	-41.7 (-57.6; -19.9)	.001	-5.2 (-26.3; 22.0)	.68
Total volume, MET h/wk ^c	8.7 (4.6)					
		1	-54.3 (-68.4; -33.8)	<.001	-10.3 (-33.9; 21.6)	.48
		2	-52.8 (-68.2; -30.0)	<.001	-8.2 (-32.8; 25.5)	.59
		3	-52.1 (-67.8; -28.6)	<.001	-7.8 (-32.8; 26.4)	.61

The geometric mean is the n th root of the product of n individual values. Geometric standard deviations correspond to the percent increase in the variable corresponding to a change of one standard-deviation unit in the logarithm of the variable.

^a Model 1 ($n = 639$): sex, age, cohort, season


Model 2 ($n = 639$): Model 1 plus socioeconomic status, maternal smoking, gestational diabetes and hypertension, and birth weight SD score.

Model 3 ($n = 637$): Model 2 plus diagnosed asthma, adult body size (height, body fat percentage), and smoking.

^b Level for statistical significance $P < 0.05$.

^c MET h/wk indicates MET hours of physical activity per week (MET = metabolic equivalents).

^d Physical activity intensity ≥ 6 MET.



*One person can have >1 reason for exclusion.

Figure 1 Flow chart of the study population. EPT, early preterm (< 34 gestational weeks); LPT, late preterm (34–36 gestational weeks)




Figure 2 Mean differences in physical activity (95% CIs, error bars) consisting of conditioning and non-conditioning leisure-time physical activity and commuting physical activity in preterm groups compared with controls (zero line). Model 1 was adjusted for sex, age, cohort and season. Model 2 was adjusted as for model 1 plus socioeconomic status, maternal smoking, gestational diabetes and hypertensive disorder during pregnancy, and birth weight SD score. Model 3 was adjusted as for model 2 plus diagnosed asthma, adult body size (height, body fat percentage) and smoking.