

Relation of Lipid-lowering Therapy to Need for Aortic Valve Replacement in Patients with Asymptomatic Mild to Moderate Aortic Stenosis

Anders M. Greve M.D. Ph.D. , Casper N. Bang M.D. Ph.D. ,
Kurt Boman M.D. Ph.D. , Kenneth Egstrup M.D. DMSci ,
Y. Antero Kesäniemi M.D. Ph.D. , Simon Ray M.D. ,
Terje R. Pedersen M.D. DMsci ,
Kristian Wachtell M.D. Ph.D. DMsci



PII: S0002-9149(19)31006-9
DOI: <https://doi.org/10.1016/j.amjcard.2019.08.037>
Reference: AJC 24165

To appear in: *The American Journal of Cardiology*

Received date: 14 May 2019
Revised date: 16 August 2019
Accepted date: 20 August 2019

Please cite this article as: Anders M. Greve M.D. Ph.D. , Casper N. Bang M.D. Ph.D. , Kurt Boman M.D. Ph.D. , Kenneth Egstrup M.D. DMSci , Y. Antero Kesäniemi M.D. Ph.D. , Simon Ray M.D. , Terje R. Pedersen M.D. DMsci , Kristian Wachtell M.D. Ph.D. DMsci , Relation of Lipid-lowering Therapy to Need for Aortic Valve Replacement in Patients with Asymptomatic Mild to Moderate Aortic Stenosis, *The American Journal of Cardiology* (2019), doi: <https://doi.org/10.1016/j.amjcard.2019.08.037>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Relation of Lipid-lowering Therapy to Need for Aortic Valve Replacement in Patients with Asymptomatic Mild to Moderate Aortic Stenosis.

Anders M. Greve, M.D. Ph.D.¹; Casper N. Bang, M.D. Ph.D.²; Kurt Boman, M.D. Ph.D.³; Kenneth Egstrup, M.D. DMSci⁴; Y. Antero Kesäniemi, M.D. Ph.D.⁵; Simon Ray, M.D.⁶; Terje R. Pedersen, M.D. DMSci⁷; Kristian Wachtell, M.D. Ph.D. DMSci⁸

Author Affiliations:

¹ Department of Clinical Biochemistry 3011, Rigshospitalet, Copenhagen, Denmark.

² Department of Cardiology, Rigshospitalet, Copenhagen, Denmark.

³ Department of Medicine, Institution of Public Health and Clinical Medicine, Umeå University, Skellefteå, Sweden

⁴ Medicinsk Afdeling, OUH Svendborg Sygehus, Denmark

⁵ Institute of Clinical Medicine, Department of Medicine, University of Oulu and Clinical Research center, Oulu University Hospital, Oulu, Finland

⁶ Manchester Academic Health Sciences Centre, University Hospitals of South Manchester, Manchester, United Kingdom.

⁷ Center for Preventive medicine, Oslo University Hospital, Ullevål and University of Oslo, Oslo, Norway.

⁸ Department of Cardiology, Oslo University Hospital, Oslo, Norway.

Contact information for the corresponding author:

Anders M. Greve, M.D. Ph.D.

Tel: +45 35453545

Department of clinical biochemistry 3011
Copenhagen, Denmark

Email: anders.moeller.greve@regionh.dk

Funding/Support: The Simvastatin and Ezetimibe in Aortic Stenosis (SEAS) trial was funded by Merck Sharp & Dohme Corp.

Disclosures: AMG: no disclosures, CNB: no disclosures, KB: Dr. Boman reports personal fees from MSD, during the conduct of the study, and personal fees from MSD, outside the submitted work. KE: no disclosures, YAK Dr. Kesäniemi reports grants from Merck Sharpe Dohme, during the conduct of the study. SR Dr. Ray reports other from Merck, during the conduct of the study. TRP: Dr. Pedersen reports grants and personal fees from Merck (MSD), grants and personal fees from Amgen, personal fees from AstraZeneca, grants and personal fees from Amgen, grants from Pfizer, during the conduct of the study. KW Dr. Wachtell reports grants and personal fees from Merck & Co, Inc., during the conduct of the study.

Assurances:

1. The publication of this study was approved by the SEAS steering committee, all patients had provided written informed consent, and the protocol was approved by local ethics committees.
2. Data analysis was performed by the following authors: Greve.
3. Clinicaltrials.gov identifier: NCT00092677

Number of display items: 2 Figures and 1 Table. Word Count (text, legends and references): 2,259.

Abstract

In this study, we aimed to determine if pretreatment low-density lipoprotein (LDL) levels and aortic stenosis (AS) severity alter the efficacy of lipid-lowering therapy on reducing aortic valve replacement (AVR). We used 1,687 patients with asymptomatic mild-to-moderate AS, who were randomly assigned (1:1) to 40/10 mg simvastatin/ezetimibe combination vs. placebo in the simvastatin and ezetimibe in Aortic Stenosis (SEAS) trial. Pretreatment LDL levels ($>4\text{mmol/L}$) and peak aortic jet velocity (3m/s) were used to partition study participants into 4 groups, which were followed for a primary endpoint of AVR. Cox regression with tests for interaction was used to study the effect of randomized treatment in each subgroup. During a median follow-up of 4.3 years (IQR 4.2-4.7 years; total 7,396 patient-years of follow-up), 478 (28%) patients underwent AVR and 146 (9%) died. A significant risk dependency was detected between simvastatin/ezetimibe combination, LDL levels and mild vs. moderate AS on rates of AVR ($p=0.01$ for interaction). In stratified analyses, randomized treatment, therefore, reduced the rate of AVR among patients with LDL levels $>4\text{mmol}$ and mild AS at baseline (HR 0.4; 95%CI: 0.2-0.9). There was no detectable effect of randomized treatment on the need for AVR in the 3 other participants subgroups. We conclude, that in a secondary analysis from a prospective randomized clinical trial, treatment with simvastatin/ezetimibe combination reduced the need for AVR in a subset of patients with mild AS and high pretreatment LDL levels (Unique identifier on clinicaltrials.gov: NCT00092677).

Keywords: ■ aortic stenosis ■ hyperlipidemia ■ statins ■ aortic valve replacement ■ prognosis

Hyperlipidemia is a risk factor for both atherosclerotic vascular and calcific aortic valve disease.¹ Lipid-lowering is key to primary and secondary prevention of vascular atherosclerosis, but do not translate to a similar benefit on slowing the progression of aortic stenosis (AS).² A prevailing opinion is therefore that AS is an unmodifiable disease, for which the only remedy is transcatheter or surgical aortic valve replacement (AVR). However, this longstanding notion was recently contested in a secondary analysis from the SEAS (simvastatin and ezetimibe in aortic stenosis) trial, which showed that simvastatin/ezetimibe slowed progression of peak aortic jet velocity in patients with mild AS and pretreatment low-density lipoprotein (LDL) levels >4mmol/L.³ Yet, it is unknown if LDL levels and AS severity also impact on the efficacy of lipid-lowering therapy to delay clinical progression of AS. The aim of this study was therefore to assess the effect of lipid-lowering on AVR in the SEAS trial, when stratified by LDL >4mmol/L and mild vs. moderate AS at baseline. We hypothesized that simvastatin/ezetimibe combination vs. placebo would reduce the need for AVR in participants with mild AS and pretreatment LDL >4mmol/L.

Methods

The SEAS trial was a multicenter, randomized, double-blind, placebo-controlled study, investigating whether intensive lipid lowering with simvastatin/ezetimibe combination vs. placebo could reduce hemodynamic progression of AS, the need for AVR and cardiovascular morbidity and mortality in patients (n=1,873), aged 45 to 85 years, with asymptomatic non-severe AS (defined as echocardiographic aortic valve thickening accompanied by a Doppler-measured aortic peak flow velocity ≥ 2.5 and ≤ 4.0 m/sec) and normal systolic left ventricular function. The main outcome including complete study protocol, study design, organization, clinical measures, exclusion criteria (most important systolic heart failure, diabetes and known ischemic heart disease) and baseline characteristics have been published previously.^{4,5} This post-hoc and not prespecified study tested

the hypothesis that the efficacy of lipid lowering on reducing AVR depends on pretreatment LDL levels and AS severity. We included all SEAS participants (n=1,687) with ascertained pretreatment LDL levels and at least 1 in-study echocardiographic examination of peak aortic jet velocity. The SEAS study complies with the Declaration of Helsinki, locally appointed ethics committee had approved the research protocol and informed consent was obtained from all subjects.

The echocardiographic study protocol, reading procedures and reproducibility have been published.⁶ Briefly, transthoracic echocardiograms were read blinded to the randomization and study visits at the SEAS echocardiographic core laboratory, located at Haukeland University Hospital in Bergen, Norway. Quantitative echocardiography was performed following the American Society of Echocardiography guidelines.⁷ Patients with peak aortic jet velocities <3 m/sec were classified as having mild AS and those with ≥ 3 -4 m/sec as having moderate AS.⁸

The primary endpoint in this study was AVR, which was an a priori defined endpoint in the SEAS trial.⁴ The efficacy of simvastatin/ezetimibe combination vs. placebo on reducing AVR was tested as the effect of randomization regardless of whether patients adhered to the treatment.

Data were analyzed using the statistical analytical software version 9.4 (SAS, Cary, NC). Continuous data are expressed as mean \pm SD and categorical variables as proportions. T-tests, Wilcoxon, Chi-square and Fisher's exact test was used as appropriate. Cox regression was used to examine the effect of simvastatin/ezetimibe combination vs. placebo on reducing the need for AVR. Absolute probabilities of AVR by randomized treatment were calculated by Fine and Gray competing risk regression, using all-cause mortality as a competing event.⁹ To identify a possible effect modification of pretreatment LDL levels and AS severity on the efficacy of randomized treatment, we tested for interactions between pretreatment LDL levels and baseline peak aortic jet velocity. For LDL, a cutoff of 4mmol/L was applied, which has previously been demonstrated

to be important for an effect of lipid-lowering therapy,³ and for peak velocity a cutoff of 3 m/s (mild vs. moderate AS) was used. Because of a significant third order interaction between treatment, pretreatment LDL and peak aortic jet velocity on AVR, the effect modification of pretreatment LDL levels on randomized treatment was investigated separately in patients with mild and moderate AS at baseline. For all hypothesis tested, a two-tailed $p < .05$ was regarded as statistically significant.

Results

Of the 1,873 SEAS study participants, 1,687 (90%) subjects had ascertained pretreatment LDL levels and at least 1 in-study measurement of peak aortic jet velocity. Randomization was well-preserved in the subset of participants eligible for this study, of which 855 (50.7%) were in the active arm (Table 1). Median follow-up was 4.3 years (IQR 4.2-4.7 years; total 7,396 patient-years of follow-up), during which 478 (28%) patients underwent AVR and 146 (9%) died. A significant risk dependency was detected between simvastatin/ezetimibe combination, LDL and mild vs. moderate AS on rates of AVR ($p = 0.01$ for interaction). In adjusted analyses, randomized treatment, therefore, reduced the rate of AVR in a subgroup of patients with LDL > 4 mmol and mild AS at baseline (HR 0.4; 95%CI: 0.2-0.9, Figure 1). These findings were consistent with results from competing risk regression, using death as a competing event, where active arm participants with LDL > 4 mmol/L and mild AS, had a significant lower probability of AVR (event rates 2% vs. 4% per year, respectively, Figure 2). There was no detectable impact of simvastatin/ezetimibe combination vs. placebo on AVR in the 3 other subgroups (Figure 1).

Discussion

This is the first study to show that lipid-lowering may reduce the need for AVR. This is an important extension of our previous report on lipid-lowering and slower progression in peak aortic

jet velocity per se.³ Taken together, these findings support the notion that lowering high LDL levels mitigates adverse endpoints related to hemodynamic and clinical progression of AS.

A large body of evidence supports that transcatheter or surgical AVR ameliorates the prognosis of patients with symptomatic, severe AS.¹⁰⁻¹² Conversely, the optimal surgical and/or medical management of patients with asymptomatic mild-to-moderate AS remains conjecture.⁸ The dearth of studies on medical therapies for patients with asymptomatic mild-to-moderate AS is partly explained by the frequent commodities, e.g. coronary artery disease, which precludes placebo-controlled randomized clinical trials. Current research has therefore turned to genetic epidemiology, which can be used to emulate a placebo-controlled randomized clinical trial by mendelian randomization.¹³ Importantly, such genetic studies have demonstrated a causal role of high LDL and lipoprotein(a) for calcific aortic valve disease and incident clinical AS.^{14,15} Of note, the level of lipoprotein(a) is, due to similar biochemical structures, included in LDL levels on routine laboratory assays. Thus, the genetic data corroborate a causal link between high LDL and/or lipoprotein(a) on incident AS, and have already been used as a template for clinical trials.¹⁶

Importantly, there is also evidence from non-genetic research, which used orthogonal methodologies to demonstrate that high lipoprotein(a) acts as a driver for the progression of AS.¹⁷ Furthermore, our results also agree with a report from the ASTRONOMER (Effects of Rosuvastatin on Aortic Stenosis Progression) trial, which found that the rate of hemodynamic progression in patients with AS related linearly to lipoprotein(a) level.¹⁸ Thus, it is prudent to suspect that the negative trials on randomized in patients with AS, were caused by the inclusion of too few patients with high concentrations of LDL and/or lipoprotein(a).

There are strengths and limitations to this study. The primary strength is the prospective, randomized, placebo-controlled, clinical trial design. As such, this is the only study that is powered

to explore possible effect modifications of LDL and mild vs. moderate AS on the efficacy of lipid-lowering therapy. Moreover, a study like the SEAS trial, is unlikely ever to be repeated because new lipid guidelines preclude the assignment of patients to placebo. Limitations include that we did not measure lipoprotein(a) and experiment wise type 1 error rate, which is compounded by the lack of a detectable impact on the overall effect of lipid-lowering therapy in the SEAS study.

In conclusion, in patients with mild asymptomatic AS and LDL greater than 4mmol/L, randomized treatment with simvastatin/ezetimibe combination vs. placebo reduced the need for AVR. Further studies aimed at determining the effect of lipid-lowering in asymptomatic patients with early AS are therefore warranted. Lipid-lowering is likely to be particularly beneficial in those with early AS and high levels of LDL and/or lipoprotein(a).

1. Rajamannan NM, Evans FJ, Aikawa E, Grande-Allen KJ, Demer LL, Heistad DD, Simmons CA, Masters KS, Mathieu P, O'Brien KD, Schoen FJ, Towler DA, Yoganathan AP, Otto CM. Calcific aortic valve disease: not simply a degenerative process: A review and agenda for research from the National Heart and Lung and Blood Institute Aortic Stenosis Working Group. Executive summary: Calcific aortic valve disease-2011 update. *Circulation* 2011;124:1783-1791.
2. Rossebo AB, Pedersen TR, Boman K, Brudi P, Chambers JB, Egstrup K, Gerds E, Gohlke-Barwolf C, Holme I, Kesaniemi YA, Malbecq W, Nienaber CA, Ray S, Skjaerpe T, Wachtell K, Willenheimer R. Intensive lipid lowering with simvastatin and ezetimibe in aortic stenosis. *N Engl J Med* 2008;359:1343-1356.
3. Greve AM, Bang CN, Boman K, Egstrup K, Forman JL, Kesaniemi YA, Ray S, Pedersen TR, Best P, Rajamannan NM, Wachtell K. Effect Modifications of Lipid-Lowering Therapy on Progression of Aortic Stenosis (from the Simvastatin and Ezetimibe in Aortic Stenosis [SEAS] Study). *Am J Cardiol* 2018;121:739-745.
4. Rossebo AB, Pedersen TR, Boman K, Brudi P, Chambers JB, Egstrup K, Gerds E, Gohlke-Barwolf C, Holme I, Kesaniemi YA, Malbecq W, Nienaber CA, Ray S, Skjaerpe T, Wachtell K, Willenheimer R, Investigators S. Intensive lipid lowering with simvastatin and ezetimibe in aortic stenosis. *N Engl J Med* 2008;359:1343-1356.
5. Rossebo AB, Pedersen TR, Allen C, Boman K, Chambers J, Egstrup K, Gerds E, Gohlke-Barwolf C, Holme I, Kesaniemi VA, Malbecq W, Nienaber C, Ray S, Skjaerpe T, Wachtell K, Willenheimer R. Design and baseline characteristics of the simvastatin and ezetimibe in aortic stenosis (SEAS) study. *Am J Cardiol* 2007;99:970-973.
6. Cramariuc D, Cioffi G, Rieck AE, Devereux RB, Staal EM, Ray S, Wachtell K, Gerds E. Low-flow aortic stenosis in asymptomatic patients: valvular-arterial impedance and systolic function from the SEAS Substudy. *JACC Cardiovasc Imaging* 2009;2:390-399.
7. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ, Chamber Quantification Writing G,

American Society of Echocardiography's G, Standards C, European Association of E. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2005;18:1440-1463.

8. Baumgartner H, Falk V, Bax JJ, De Bonis M, Hamm C, Holm PJ, Iung B, Lancellotti P, Lansac E, Rodriguez Munoz D, Rosenhek R, Sjogren J, Tornos Mas P, Vahanian A, Walther T, Wendler O, Windecker S, Zamorano JL, Group ESCSD. 2017 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2017;38:2739-2791.

9. Andersen PK, Geskus RB, de Witte T, Putter H. Competing risks in epidemiology: possibilities and pitfalls. *Int J Epidemiol* 2012;41:861-870.

10. Schwarz F, Baumann P, Manthey J, Hoffmann M, Schuler G, Mehmel HC, Schmitz W, Kubler W. The effect of aortic valve replacement on survival. *Circulation* 1982;66:1105-1110.

11. Smith CR, Leon MB, Mack MJ, Miller DC, Moses JW, Svensson LG, Tuzcu EM, Webb JG, Fontana GP, Makkar RR, Williams M, Dewey T, Kapadia S, Babaliaros V, Thourani VH, Corso P, Pichard AD, Bavaria JE, Herrmann HC, Akin JJ, Anderson WN, Wang D, Pocock SJ. Transcatheter versus Surgical Aortic-Valve Replacement in High-Risk Patients. *N Engl J Med* 2011;364:2187-2198.

12. Mack MJ, Leon MB, Thourani VH, Makkar R, Kodali SK, Russo M, Kapadia SR, Malaisrie SC, Cohen DJ, Pibarot P, Leipsic J, Hahn RT, Blanke P, Williams MR, McCabe JM, Brown DL, Babaliaros V, Goldman S, Szeto WY, Genereux P, Pershad A, Pocock SJ, Alu MC, Webb JG, Smith CR. Transcatheter Aortic-Valve Replacement with a Balloon-Expandable Valve in Low-Risk Patients. *N Engl J Med* 2019;380:1695-1705.

13. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Human molecular genetics* 2014;23:R89-98.

14. Smith JG, Luk K, Schulz CA, Engert JC, Do R, Hindy G, Rukh G, Dufresne L, Almgren P, Owens DS, Harris TB, Peloso GM, Kerr KF, Wong Q, Smith AV, Budoff MJ, Rotter JI, Cupples LA, Rich S, Kathiresan S, Orho-

Melander M, Gudnason V, O'Donnell CJ, Post WS, Thanassoulis G. Association of low-density lipoprotein cholesterol-related genetic variants with aortic valve calcium and incident aortic stenosis. *Jama* 2014;312:1764-1771.

15. Thanassoulis G, Campbell CY, Owens DS, Smith JG, Smith AV, Peloso GM, Kerr KF, Pechlivanis S, Budoff MJ, Harris TB, Malhotra R, O'Brien KD, Kamstrup PR, Nordestgaard BG, Tybjaerg-Hansen A, Allison MA, Aspelund T, Criqui MH, Heckbert SR, Hwang SJ, Liu Y, Sjogren M, van der Pals J, Kalsch H, Muhleisen TW, Nothen MM, Cupples LA, Caslake M, Di Angelantonio E, Danesh J, Rotter JJ, Sigurdsson S, Wong Q, Erbel R, Kathiresan S, Melander O, Gudnason V, O'Donnell CJ, Post WS. Genetic associations with valvular calcification and aortic stenosis. *N Engl J Med* 2013;368:503-512.

16. Tsimikas S. Potential Causality and Emerging Medical Therapies for Lipoprotein(a) and Its Associated Oxidized Phospholipids in Calcific Aortic Valve Stenosis. *Circulation research* 2019;124:405-415.

17. Zheng KH, Tsimikas S, Pawade T, Kroon J, Jenkins WSA, Doris MK, White AC, Timmers N, Hjortnaes J, Rogers MA, Aikawa E, Arsenault BJ, Witztum JL, Newby DE, Koschinsky ML, Fayad ZA, Stroes ESG, Boekholdt SM, Dweck MR. Lipoprotein(a) and Oxidized Phospholipids Promote Valve Calcification in Patients With Aortic Stenosis. *J Am Col Cardiol* 2019;73:2150-2162.

18. Capoulade R, Yeang C, Chan KL, Pibarot P, Tsimikas S. Association of Mild to Moderate Aortic Valve Stenosis Progression With Higher Lipoprotein(a) and Oxidized Phospholipid Levels: Secondary Analysis of a Randomized Clinical Trial. *JAMA cardiol* 2018 (epub ahead of print).

Figure legends




Figure 1. Simvastatin/Ezetimibe combination reduced the need for aortic valve replacement in SEAS trial participants with mild aortic stenosis and high lipoproteins at baseline.

Forest plot of hazard ratios for aortic valve replacement by subgroups of patients with mild vs. moderate aortic stenosis (peak jet velocity >3m/sec), and high low-density lipoprotein cholesterol (LDL >4mmol/L) at enrollment (p=0.01 for third order interaction between treatment, peak jet velocity, and LDL, p=0.02 for second-order interaction between treatment and high LDL and p=0.03 for second-order interaction between treatment and mild vs. moderate aortic stenosis when including the significant third order interaction term).




Figure 2. Cumulative incidence of aortic valve replacement in SEAS participants with mild aortic stenosis and LDL >4mmol/L at baseline.

Cumulative incidence of aortic valve replacement according to treatment in patients with mild aortic stenosis (peak jet velocity >3m/sec) and LDL >4mmol/L, during a median follow-up of 4.3 years in the SEAS trial. Fine and Gray competing risk regression, using all-cause mortality as a competing event, was used to calculate the absolute probabilities.

Table 1. Baseline characteristics in SEAS participants with ascertained peak aortic jet velocity and low-density lipoprotein levels at baseline.*

Baseline variables	Simvastatin/Ezetimibe (n=855)	Placebo (n=832)	P-value
Clinical parameters			
Age (years)	67.5±9.4	67.1±9.7	0.41
Men	62.0%	60.6%	0.55
Systolic blood pressure (mmHg)	145.5±20.1	143.9±20.1	0.09
Diastolic blood pressure (mmHg)	82.1±10.7	82.0±10.0	0.83
Body mass index (kg/m ²)	26.9±4.4	27.0±4.4	0.50
Peak aortic jet velocity (m/sec)	3.1±0.5	3.1±0.5	0.75
Tricuspid aortic valve	94.9%	93.7%	0.35
Left ventricular ejection fraction (%)	65.8±8.3	65.5±8.3	0.96
Left ventricular mass index (g/m ²)	100.3±29.1	99.3±31.0	0.53
eGFR (ml/min per 1.73m ²)	68.5±12.7	68.4±12.0	0.84
Glucose (mmol/ml)	5.3±0.8	5.3±0.8	0.46
High-density lipoprotein (mmol/l)	1.5±0.4	1.5±0.4	0.73
High-density lipoprotein (mg/dl)	57.9±15.4	57.9±15.4	0.73
Low-density lipoprotein (mmol/l)	3.6±0.9	3.6±0.9	0.82
Low-density lipoprotein (mg/dl)	139.0±34.8	139.0±34.8	0.82
Medicine			
Digoxin	3.0%	2.3%	0.33
Ca ²⁺ -blocker	16.6%	17.1%	0.80
Beta-blocker	26.8%	28.5%	0.43
Renin-Angiotensin inhibitor	25.4%	25.5%	0.96
Diuretics	22.0%	25.0%	0.14

*This baseline Table differs from the original SEAS publication by including only participants with ascertained peak aortic jet velocity and low-density lipoprotein levels at enrollment. eGFR = estimated glomerular filtration rate.