



Exchange of Microbiomes in Plant-Insect Herbivore Interactions

 A. M. Pirttilä,^a V. Brusila,^{a,b} J. J. Koskimäki,^a P. R. Wäli,^a A. L. Ruotsalainen,^a M. Mutanen,^a A. M. Markkola^a

^aEcology and Genetics, University of Oulu, Oulu, Finland

^bNatural Resources Institute Finland, Helsinki, Finland

ABSTRACT Prokaryotic and eukaryotic microbial symbiotic communities span through kingdoms. The vast microbial gene pool extends the host genome and supports adaptations to changing environmental conditions. Plants are versatile hosts for the symbionts, carrying microbes on the surface, inside tissues, and even within the cells. Insects are equally abundantly colonized by microbial symbionts on the exoskeleton, in the gut, in the hemocoel, and inside the cells. The insect gut is a prolific environment, but it is selective on the microbial species that enter with food. Plants and insects are often highly dependent on each other and frequently interact. Regardless of the accumulating evidence on the microbiomes of both organisms, it remains unclear how much they exchange and modify each other's microbiomes. In this review, we approach this question from the point of view of herbivores that feed on plants, with a special focus on the forest ecosystems. After a brief introduction to the subject, we concentrate on the plant microbiome, the overlap between plant and insect microbial communities, and how the exchange and modification of microbiomes affects the fitness of each host.

KEYWORDS herbivory, insect-microbe interactions, microbiome, plant-microbe interactions, symbiosis

The current trend of climate change provokes insect expansion to new geographic areas, providing new threats, especially in the Northern Hemisphere. In North America, the impact of insect pests and pathogens of forests has been evaluated to be equal to that of fire and forestry combined (1). Invasive forest pests can eliminate keystone species and have large negative effects on biodiversity and carbon cycling, which can eventually impair ecosystem services (2). The spread of invasive species is complex and is affected by many factors, including insect lifestyle, the presence of parasitoids and predators, and the palatability level of host tissues due to, for example, phenol-richness, toxic compounds, or hard or otherwise protected plant parts (3). An important factor that is often neglected is the presence of microbial communities, both in the insect and in the plant host tissues.

Microbial symbionts of prokaryotic and eukaryotic microorganisms exist throughout kingdoms. Microbes are short-lived and possess a huge diversity of metabolic pathways, which, together with their horizontal gene transfer systems, enable rapid evolution and environmental responses (4). In symbiosis, the huge gene pool of microbial communities extends the host genome and supports adaptations to changing environmental conditions (5). Specifically, plants are prolific hosts of microbes, as they carry microbial symbionts both on the surface, called epiphytes, as well as inside tissues, known as endophytes, and even in the cells (6). The plant microbiomes consist of bacteria, fungi, oomycetes, algae, and protozoa (7). The microbial symbionts can live in virtually any plant tissue, flower, bud, seed, stem, leaf, or root (6). The mutualistic microbes of plants are typically acquired horizontally from the environment, although a vertical transmission through seeds occurs in the most intimate symbioses (5). For example, endophytic fungi are classified into four different classes, according to their transmission, biodiversity, and host range (8). A study on a conifer, namely, the Norway spruce, showed that a tree

Editor Jacob Yount, Ohio State University

Copyright © 2023 Pirttilä et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to A. M. Pirttilä, am.pirttila@oulu.fi.

The authors declare no conflict of interest.

Published 7 March 2023

individual can host 86 to 110 fungal needle endophytes (9), whereas a single needle can be colonized by 34 different fungal species (10).

Similar to plants, insects are colonized by microbial symbionts, with their habitats being the exoskeleton, the gut, the hemocoel, and the cell interior. The insect microbiota consist of eukaryota, such as fungi and protists, as well as archaea and bacteria, and bacteriocyte symbioses are formed by intracellular microbes (11, 12). Insect microbiomes can be divided into open associations, which are invaded by external microbes, and closed associations, in which the invasion is prevented. For example, the exoskeleton and gut are open for invaders. As many as 1,000 culturable bacteria are found on the exoskeleton of the fruit fly (13). The insect gut is a prolific environment but is selective on the microbes that enter with food. The hindgut and the ileum are typically the regions with the highest counts of microbes (11). The compositions of microbial gut communities can also change along the insect development. In the gut of the African cotton leafworm (*Spodoptera littoralis* [Boisduval]), which is among the most detrimental agricultural pests worldwide, bacterial species belonging to *Enterococcus*, *Pantoea*, and *Citrobacter* are abundant in early life, and Clostridia becomes the most common class in adults (14).

Some insect species have cuticular structures, which are specialized in providing a niche for the microbial counterparts. Examples are the mycangia host fungi or bacteria that are needed for the insect's offspring (11). Among insects, intracellular microbes are common in specific groups, where they are found in cells specialized for this purpose, called bacteriocytes or mycetocytes. This is a closed association, in which the endosymbionts dominating the bacteriocyte or mycetocyte, which are considered to be primary symbionts, have no access to the environment (11). In the plant sap-feeding insects, the endosymbionts are important in providing essential amino acids for their hosts, as the phloem sap is a poor nitrogen source (15). However, it is typical for this insect group to also host secondary symbionts that are associated with the bacteriocytes or mycetocytes. The secondary symbionts are similarly vertically transmitted, but they can also transfer horizontally and reside in the insect sheath cells or in the hemolymph. The secondary symbionts can provide the host with traits, such as thermal tolerance or resistance to parasitoids or fungi (11).

The two types of higher organisms, namely, plants and insects, are often highly dependent on each other and frequently interact. Regardless of the accumulating evidence on the microbiomes of both organisms, it largely remains unclear how much their microbiomes overlap and how much they modify each other's microbiomes. In this review, we approach this question from the point of view of herbivores feeding on plants, with a special focus on the forest ecosystems. We will first take an overview on the plant-herbivore interaction and how the plant microbiome is affected by herbivory. Then, we will focus on the existing knowledge regarding the overlap between plant and herbivore microbial communities (excluding viruses). Finally, we will discuss how the modification of these microbiomes affects the fitness of each host and how they could potentially be manipulated to improve plant fitness.

PLANT DEFENSE IN HERBIVORE INTERACTIONS

Plants have developed mechanisms to sense the presence and the type of an invader. The recognition of the invader triggers signaling pathways that lead to specific responses. A number of signaling compounds and crosstalk between the pathways are required for the activation of the defense responses, which are briefly discussed below to allow for an understanding of the intricacy of the tritrophic interactions between plants, insect herbivores, and microbes (for further details, see [16, 17]).

There are two main mechanisms to identify the invaders: pattern recognition and effector molecule-based recognition. Pattern recognition is based on elicitors and pattern recognition receptors (PRRs) that can detect a range of molecular patterns. In the case of herbivory, such patterns are called herbivorous insect-associated molecular patterns (HAMP) (17). In more detail, damage-associated molecular patterns (DAMP) are endogenous elicitors that are released, for example, from the cell wall (18). Microbe-associated molecular patterns (MAMP) or pathogen-associated molecular patterns

(PAMP) are derived from microbial molecules. The recognition of the insect may also be based on the MAMP-patterns of insect endosymbionts (19). The pattern recognition initiates the PAMP-triggered immunity (PTI) (16). Besides the molecular patterns, the plant recognizes the effector molecules of pathogens by their resistance (R)-gene products, and such gene-for-gene-recognition triggers the effector-triggered immunity (ETI). For example, the components of insect saliva can serve as effectors. Pattern-based recognition occurs at the cell membrane, and the effector-based recognition is intracellular (18).

The sensing of an invader often leads to the activation of a defense pathway. The salicylic acid (SA) signaling pathway is typically triggered by biotrophic pathogens, and the jasmonic acid and ethylene (JA/ET)-mediated signaling pathway is triggered by wounds, herbivore insects, and necrotrophic pathogens. The activation of local defense responses through SA often triggers systemic acquired resistance (SAR), which requires long-distance signaling and primes the undamaged tissues (16). The association of the plant with mutualistic microbes can trigger induced systemic resistance (ISR), which also leads to priming (16, 20). Priming is an enhanced ability to respond faster and stronger to pathogen attacks (21). Basically, priming makes plants more sensitive to JA and ET, which control ISR (16). A sucking insect may also promote the plant defense against other herbivores, as the honeydew secretions of the rice brown planthopper (*Nilaparvata lugens*) contain members of the gut microbiome that induce systemic resistance on the monocot host, namely, rice (*Oryza sativa*). This further leads to the accumulation of phytoalexins and the release of volatile organic compounds that attract herbivore enemies on rice leaves (22).

The recognition of suitable and high-quality plants for feeding and reproduction is vital for herbivore insects (23), as most of them attack specific groups of plant species, with only 10% being generalists (24). Herbivorous insects can be classified in several ways, for example, based on the host taxonomy, or based on the type of consumed tissue, into algivorous (algae), frugivorous (flowers), xylophagous (wood), folivorous (leaves), granivorous (seeds), and mucivorous (sap) species, or, based on their feeding mode, into internal, external, or, alternatively, into sucking or chewing insects (25). Further, herbivores can be classified by the level of host species specialization as monophagous, oligophagous, or polyphagous feeders (i.e., those feeding on a single plant species, those that feed on several closely related species or genera, and those that feed on a wide variety of different plant groups, respectively) (26). Plants can recognize the insect species and adjust their defense responses accordingly. Overall, herbivory can trigger a cascade of community-wide interactions (27).

The defense strategy in many plants is to accumulate high quantities of compounds that are toxic to insects, whereas other plant species prefer to use their resources to minimize herbivore damage via rapid growth and development, dispersion, or choice of habitat. The selection of strategy may vary between plant genotypes from partitioning resources toward growth or defense (28, 29). Plant defense against herbivores can be divided into constitutive or induced reactions. The induced defense is often similar between plants, but the constitutive defense can change between plant species (30). The induction of the defense leads to the production of JA, systemin, oligogalacturonic acid, proteinase inhibitors, and hydrogen peroxide. The induced defense responses lack the capacity to develop a full resistance, but they can reduce the growth and survival of the insect herbivore (31). The plant defense reactions can also result in the development of physical barriers, such as lignification or resin production. Furthermore, plants typically produce volatile compounds that deter further herbivores and communicate the defense to neighboring plants (17, 30).

The range of plant defense compounds is broad with mechanisms of insect membrane disruption, suppression of metabolism, interference with signal transduction, and interference with development through imbalanced hormone regulation (30, 32). Plant defense responses may impact the settling, feeding, oviposition, fecundity, and fertility of insect herbivores (32). Herbivores attempt to overcome the plant defenses via constitutive or induced mechanisms (30), many of which are based on symbiotic microorganisms (17, 33, 34), which are discussed below. Insects that have a limited host range can lean on constitutive

adaptations to survive unfavorable plant compounds, whereas herbivores with a wide range of plant species more readily use induced adaptations to overcome plant defenses. In natural ecosystems, the majority of plant-herbivore interactions reach a stalemate, in which both partners suffer but survive (30).

PLANT-ASSOCIATED MICROBES

Even if the members of the plant microbiome are not directly associated with insects, they can play a significant role in this interaction. The microbiome can influence the nutrient status of the plant, interfere with plant-pathogen interactions, and modify the tolerance of the plant to abiotic stresses (5). On the other hand, plant pathogens and various abiotic factors can affect the plant-microbe interactions (17, 20, 35–37).

A great part of microbial interactions occurs in the plant roots, where the soil is typically the source of the microbes. A rhizobial symbiosis is formed in legumes when the plant host develops root nodules to allow for the bacterial fixation of atmospheric nitrogen (38). Rhizobia can play a role in the herbivory tolerance of legumes by inducing JA production and the development of systemic resistance (39). Another nitrogen-fixing symbiosis, called actinorhiza, is formed between the soil bacteria belonging to the genus *Frankia* and many plant species, including forest trees (40). In the red alder (*Alnus rubra* Bong), symbiosis with *Frankia* increased the herbivory of young leaves by the black slug (*Arion rufus* L.) (41). However, the accumulating evidence on actinorhizal symbiosis suggests that these symbionts can induce plant resistance, similar to rhizobia (42). There are also numerous soil bacteria living in the plant rhizosphere, which are called plant growth-promoting rhizobacteria (PGPR). The PGPR can fix nitrogen, increase the availability of nutrients, affect root growth and morphology, and promote the formation of other symbioses, such as rhizobial or mycorrhizal symbioses (43). Specifically, PGPR can induce systemic resistance of the host plant and help deter herbivores. For example, in cotton, the application of PGPR induces systemic resistance and the higher production of gossypol, which is a secondary metabolite. These responses reduced the larval feeding and development of beet armyworm (*Spodoptera exigua* Hübner) in PGPR-treated cotton plants (44). The inoculation of PGPR can also lead to the induced production of plant volatiles, which attract predatory earwigs (Dermaptera) toward the herbivore attack (45). However, the protection of the host plant is highly dependent on the combination of the PGPR and the herbivore species (46).

The most well-known fungal mutualists of plants, which are mycorrhiza that also dwell in the roots and rhizosphere, have been reported to influence plant herbivore status (47). Of all mycorrhizal types, arbuscular mycorrhiza (AM) is the most common, occurring in 74% of Angiosperm species. Orchid mycorrhiza is found in 9% of Angiosperm species, with ectomycorrhiza occurring in 2% of Angiosperm species, and the rarest mycorrhizal type, namely, ericoid mycorrhiza, occurring in only 1% of Angiosperm species (48). However, there are exceptions, as in the boreal forests, the dominant vegetation consists of ectomycorrhizal trees and ericoid mycorrhizal dwarf shrubs. The various mycorrhizal types, specifically arbuscular mycorrhiza, provide phosphate for the plant host, which delivers carbon in different forms to the fungus (49). The mycorrhizal status of the plant can affect the herbivore interaction. For example, mycorrhizal plants can carry more herbivore insects, but the herbivore survival is lower, meaning that the host plant therefore suffers less damage than do nonmycorrhizal plants (47). The result of the interaction also depends on the type of the herbivore, as phloem feeders, monophagous chewers, and oligophagous chewers thrive on mycorrhizal plants, but polyphagous chewers and mesophyll feeders are more successful on a nonmycorrhizal host. Of the various mycorrhizal types, AM fungi increase the performance of sucking insects, but the AM species *Rhizophagus irregularis* reduces chewing insect damage on the host plant more than the other mycorrhizal fungi (47). The AM fungi can induce host defense against herbivory, depending on the species of the interacting partners. For example, the AM fungi induce the defense of beggarticks (*Bidens frondosa* L.) to overcome cotton leafworm (*S. littoralis*) feeding, but they have no such

effect on the cabbage moth (*Mamestra brassicae* L.) (50). On the other hand, the AM fungi can also suppress defenses against herbivores (51).

Besides the most intimate and specialized mycorrhizal symbiosis, endophytes that reside in all plant tissues form interactions with plants, ranging from mutualism to parasitism (6, 52). Dark septate endophytic (DSE) fungi commonly occur in plant roots, but their ecological functions are currently unclear (53). Both bacterial and fungal endophytes living in plant shoots can protect the host from environmental stresses, induce plant resistance, produce secondary metabolites to repel pathogens and herbivores, and promote plant growth (6). For example, the bacterial endophytes colonizing the shoot tips of mountain birch (*Betula pubescens* subsp. *czerepanovii* [N. I. Orlova] Hämet-Ahti) may promote tree recovery through sprouting after an attack by the winter moth (*Operophtera brumata* L.) (54). Endophytes have an important and versatile role in plant-herbivore interactions (55, 56), as they participate in the microbiome interactions and exchange, which is discussed in more detail in the sections below.

INTERACTIONS BETWEEN THE MICROBIOMES OF PLANTS AND HERBIVORES

Endophytes. Many fungal endophytes can produce compounds that are toxic to herbivores (57–60). Sometimes, the toxic effects can be passed on to the second generation of herbivores, regardless of whether the herbivore is feeding on the plant that carries the endophyte (61), or even to a parasitoid, hyperparasitoid, or predator of the herbivore (62–64). In forest trees, the interaction is complicated, as tree leaves are colonized horizontally by fungal endophytes that make highly localized infections (6).

In conifers, the endophytic fungi have often been associated with the capacity to repel insect herbivores (65). As early as 1978, Carroll and Carroll (66) suggested that the foliar endophytes of conifers could be mutualistic symbionts. According to their studies on the Douglas fir (*Pseudotsuga menziesii* [Mirb.] Franco), the needle endophyte *Rhabdocline parkeri* produced a tolerance to the gall-forming needle pest *Contarinia pseudotsugae* (Condrashoff). They showed higher rates of mortality of the larvae in the galls of endophyte-infested needles (67). In further studies, a fraction of the metabolites from *R. parkeri* exhibited cytotoxicity by reducing the growth of spruce budworm (*Choristoneura fumiferana* Clemens) larvae (68). In more recent studies, the toxin rugulosin, which is produced by the fungal endophyte *Phialocephala scopiformis*, significantly reduced feeding by the spruce budworm on white spruce needles (69–72). Similarly, insecticides have been isolated and characterized from the endophytic fungi of red spruce needles (73). In another study on the Scots pine (*Pinus sylvestris* L.), Saikkonen et al. (74) showed that 5-year-old trees hosting endophytic fungi had reduced performance in 40% of the needles of the pine sawfly (*Neodiprion sertifer* Geoffroy). They found the relative growth rate of the larvae to be lower and the larval period to be longer in trees that had higher mean endophyte abundances. However, they saw no similar effects on the performance of another herbivore, namely, the aphid *Scizolachnus pineti* (Fabricius) (74).

On the other hand, when the life-history performances of the winter moth, birch leaf rolling weevil (*Deporaus betulae* L.), and birch arge (*Arge clavicornis* Fabricius) were studied with respect to endophyte infection in the deciduous tree mountain birch, Saikkonen et al. (74) found no correlation between herbivory and the presence of endophytes. They suggested that there is such seasonal and spatial variation in the colonization of tree leaves by endophytic fungi, affected by the neighboring vegetation, density of trees, weather, and topography of the ground, that consistent and effective defense against insects is not formed. Ahlholm et al. (75) further analyzed the presence of the two most common endophyte genera, namely, *Fusicladium* and *Melanconium*, along with the pathogenic rust fungus *Melampsorium botulinum*, in mountain birch at the end of the growing season. The successes of the autumnal moth (*Epirrita autumnata* L.), the leaf chewer sawfly (*Cladius compressicornis* Lepeletier), the leaf skeletonizer sawfly (*Dineura virididorsata* Schmidt and Walter), and the beetle *Deporaus betulae* L. were dependent on the fungal densities on the mountain birch leaves only in specific cases. The sawfly *D. virididorsata* was more abundant with increased fungal endophyte densities and less frequent with the presence of the rust

fungus *M. botulinum*. Similarly, the rust fungus decreased the performance of the autumnal moth on mountain birch. They suggested that fungal plant pathogens that cause premature senescence and abscission negatively affected the herbivore species that attack mountain birch just before leaf fall (75). The more consistent efficiency of endophytic fungi against insect herbivores in conifers could be explained by the fact that conifers carry needles for several years, compared to deciduous trees, which grow new leaves every year. Therefore, the interactions of endophytic fungi with conifer needles to repel insect herbivores may have developed toward higher mutualism (65).

Microbial transmission (insects as vectors). Many plant-associated microbes, especially pathogens, can use insects as vectors and are partially or totally dependent on insects for transmission. The transmitted microbes include fungi, bacteria, phytoplasmas, and protozoa (76). The vector-pathogen interaction is often specific, and the vector-borne plant pathogens can be transmitted externally or internally within the insect (76). For example, the insect can become smeared with bacteria or fungal spores (77).

Nonpersistent plant pathogens, also known as stylet-borne pathogens, are acquired and inoculated after short feeding periods, as some pathogens can live in the insect vector for only a few hours. Plant pathogens that thrive in the vector from one to four days are called semipersistent pathogens. In turn, persistent or “circulative” pathogens are accumulated internally and are released to a new plant through the insect mouthpart. Some may even multiply in the vector, and such propagative pathogens may live in the insect for the rest of their lives. In some cases, there is an incubation period between the acquisition and the transition of the plant pathogen (76), and sexual transmission between the vectors has also been observed (78). Although the plant pathogens vectoring in insects are best studied, it has been shown in grapevine (*Vitis vinifera* L.) that a phloem-feeding insect (American sap-feeding leafhopper *Scaphoideus titanus* Ball) can transfer full endophytic bacterial communities between plants. The transferred communities were most similar to those found in plant roots, even though *S. titanus* feeds on the stems of grapevines (56). This is not surprising, as a number of reports show plant root-colonizing bacteria, namely, PGPR, being transmitted to new plants via insects (79, 80).

An example of insect-mediated microbial transmission in a forest setting is the case of bark beetles (Coleoptera: Curculionidae: Scolytinae) and their fungal symbionts. The beetles feed and reproduce inside conifer bark and transfer a vast number of fungal symbionts between trees (81). Many of these bark beetle species live only in dead or decaying coniferous trees; however, there is a group that invades and kills healthy trees (82). The majority of fungal taxa vectoring between bark beetles and trees belong in the order Ophiostomatales (Ascomycota) (83). The fungal spores are transferred in the exoskeletons of the bark beetles, in mites vectored by the beetles, or in mycangia from one tree to another (77). These fungi can be pathogenic, parasitic, or commensalistic in both of the hosts (84). They may provide nutrition for the larvae of the bark beetle (85) and may help the beetles overcome tree defenses (86). For example, the fungus *Ceratocystis polonica* that vectors in the European spruce bark beetle (*Ips typographus* L.), helps the beetle invade the Norway spruce (*Picea abies* L.) via the degradation of plant defense compounds, namely, stilbenes (87). The Ophiostomatalean fungi can also produce volatiles that affect the behavior of the beetles, functioning as attractants or repellents, but the beetle behavior can vary by the fungal species (88), the developmental stage of the beetle, and the genotype of the host tree (89).

Overlap between plant and insect microbiomes. An insect or a plant host can maintain a multitude of different microbial symbionts, with each having specific functions in the host (6, 90). A microbial strain can be pathogenic in a plant host and beneficial for the insect host, whereas another microbe with positive effects in the plant host can be pathogenic to the insect (91, 92, Table 1). For example, *Beauveria bassiana* is an endophytic fungus on a wide array of plants and infects more than 700 insect species as an entomopathogen (93). In addition, the endophytes *Clonostachys rosea*, *Metarhizium anisopliae*, and *Cordyceps fumosorosea* of English oak (*Quercus robur* L.), *Metarhizium anisopliae* of Chinese yew (*Taxus chinensis* [Rehder and E.H. Wilson] Rehder), *Cordyceps farinosa* of European

beech (*Fagus sylvatica* L.), and *Ophiocordyceps sobolifera* of cacao (*Theobroma cacao* L.) can act as entomopathogens (94).

On the other hand, the plant-based diet of the insect, which consists of plant material and endophytic microbes, shapes the microbial composition of the insect gut (95–98), and this can be linked with the ability of the insect to defeat plant defenses (99, 100). In the study by López-Fernández et al. (56) on grapevines and the phloem-feeding grasshopper (*S. titanus*), the plant endophytic and insect gut communities were similar, but the plant communities were dominated by Proteobacteria, whereas the insect communities were dominated by Firmicutes (56). Similarly, in the emerald ash borer (*Agrilus planipennis* Fairmaire), which is an herbivorous pest of ash trees, the composition of the leaf microbiome was a strong predictor of the gut microbial community structure in the adult insects. The leaf microbiome of the tree explained 53% and 48% of the variation in the fungal and bacterial communities of the emerald ash borer, respectively (100). However, the bacterial endophytic communities of the roots, phloem, and bark of the pine species *Pinus arizonica* Engelm. and *P. durangensis* Martínez were significantly different from the gut communities of a bark beetle (*Dendroctonus rhizophagus* Thomas and Bright) that colonizes the saplings of the pines (101). Similarly, nine polyphagous caterpillar species feeding on five various tree species had clearly different gut microbiomes, compared to the host leaves, with respect to bacterial species. However, there were similarities between the fungal microbiomes of the insect guts and the host leaves (102). Therefore, the extent of the overlap between the plant and insect microbiomes is likely highly dependent on the combinations of species.

The plant microbiome is affected by insect herbivory, as well. In bittercress (*Cardamine hirsuta* L.), herbivory by a leaf-mining fly (*Scaptomyza nigrita* Latreille) led to a higher abundance of various microbiome taxa. However, the increased abundance reflected community-wide compositional shifts toward lower ecological diversity in the bittercress. Such shifts included plant-pathogenic members of the microbiome, such as *Pseudomonas syringae*, growing toward higher populations (103). A similar observation was made when the effect of pea aphid (*Acyrtosiphon pisum* Harris) was studied on the distribution of *P. syringae* among epiphytic communities, as the bacterium thrived better on aphid-infested leaves. The aphid feeding, attributed to the presence of honeydew, had a pronounced effect on *P. syringae* populations, which were initially small on the leaves (104). In beech and oak, the feeding of lepidopterous larvae caused changes in the bacterial communities of the leaves, which were associated with the increase of available nitrogen (105). However, in strawberry, the flower microbiome was primarily shaped by the plant genotype and not by *Vanessa cardui* L. caterpillars. Specifically, the plant volatiles, such as terpenoids and benzenoids, were mainly responsible for determining the structure of the bacterial and fungal communities of the strawberry flowers (106). This suggests that the plant microbiome is reasonably resilient to changes by herbivory or other plant-insect interactions.

Benefits and disadvantages of obtaining new microbiome members. In general, microbes can have a prominent effect on the host range of insects (107). For example, the red turpentine beetle (*D. valens* LeConte) acts as a vector for fungi that have enabled their invasion from Northern America to China (81, 108). In its native range, the beetle is only a minor threat to pine trees, but it has caused serious mortality in pines on the other continent (108). Taerum et al. (81) identified a large shift in the assemblage of the invasive red turpentine beetle fungal symbionts. They concluded that the fungal community shift had enabled the beetle to become invasive in China (81).

Adaptation to a new host can happen fast via horizontal gene transfer from microbes to insects (109), in which the insect microbiome can enable rapid adaptation to disturbances in the environment (110). Microbes may detoxify plant defense compounds and provide enzymes with which to digest the plant material (111). For example, the mountain pine beetle (*D. ponderosae* Hopkins) can consume terpenes with the help of gut microbes that belong to the genera *Pseudomonas* and *Rahnella*, which have the capacity to degrade terpenes (33). The gut microbes of wood-boring Cerambycidae, Curculionidae, and Siricidae, as well as some lepidopteran species, can produce cellulases for the food digestion of their hosts (112–115). Similarly, proteinase inhibitors that are produced by the plant host may be

overcome via the hyperproduction of proteases by the symbiotic bacteria (116). In Cerambycidae, the midgut microbes also synthesize essential nutrients for the host (117–119), and they additionally possess the capacity for nitrogen fixation (120). Specifically, in the sap-feeding Hemiptera, in the process of evolution, the insects have become dependent on the biosynthesis of amino acids and vitamins by the gut microbiome, which is obtained from the host plant (11).

Mutualists of herbivorous insects that are plant pathogens can also help in the invasion of the plant by circumventing plant defenses. Such relationship may have evolved from a symbiotic association with the host, plant, or insect that was followed by adaptation to the new host (92, 121). An insect-associated microbe can suppress the plant defense via various mechanisms of manipulating plant signaling (17, 122). The weakened plant defenses can also result from a synergy between multiple microorganisms (123). For example, many phytopathogenic bacteria and fungi manipulate cytokinin signaling to attack the plant. Microbes can either produce cytokinins or modify cytokinins that are produced by the plant (124). The plant defense system can further be manipulated via the exploitation of the existing antagonism between the SA and JA signaling networks (125). Microbes can also cause indirect changes in the plant, such as altered volatile production and visual cues (17, 23, 126).

The timing of the interactions between plants, insects, and microbial symbionts may be crucial for a successful microbiome exchange. For example, in Silver birch (*Betula pendula* L.), the phenological synchrony between the emergence of the overwintering herbivore gypsy moth (*Lymantria dispar* L.) and the budding of the host plant affected herbivore survival through the gut microbiome of the insect. There was clearly a lower diversity of bacteria in the guts of asynchronous larvae than in those of synchronous larvae, which lead to the lower susceptibility of asynchronous larvae toward the entomopathogenic bacterium *Bacillus thuringiensis* (127).

Besides defeating plant defenses and supporting exploitation by the insect herbivore, the transmitted microbes can alter the survival, fecundity, and immunity of the insect host and can therefore alter the fitness of the herbivore (126). The manipulation of the insect microbiome can shorten the life span, increase the latent period, block pathogen transmission, change reproduction, and increase insect susceptibilities to natural enemies (90). The newly acquired microbes can cause direct and indirect changes in insect behavior, such as altered host preferences or feeding (126). For example, in the lodgepole pine (*Pinus ponderosa* Douglas ex C. Lawson), the beneficial phloem endophyte *Bacillus pumilus* is antagonistic toward a fungal symbiont of the mountain pine beetle (*Dendroctonus ponderosae*) (128).

Besides the plant pathogens vectoring in insects, the consequences of microbiome exchange have rarely been studied in plants. The best example is the widely studied and used *B. thuringiensis*, which causes toxicity to the larvae of many insects through the formation of crystal proteins but acts as a beneficial endophyte inside plants (91). When the phloem-feeding grasshopper *S. titanus* transferred an entire endophytic community to new grapevine plants, the receiver plants were healthy. López-Fernández et al. (56) suggested that the endophytes had a mutualistic role in the grapevine, as their earlier studies had shown that the endophytes stimulate growth and protect the grapevines from pathogens (129).

CONCLUSIONS

Gupta and Nair (130) recently summed up the various roles of gut microbes in the survival and adaptation of insects to specific environments. However, no great generalizations can be made regarding the interactions between plant and herbivore microbiomes or on how the hosts affect each other's microbial communities and further fitness. The interactions between each species of plant, each species of insect, and their microbiomes is likely highly specific and dependent on the feeding mode and environmental conditions of the insect (Fig. 1; Table 1). As data from high-throughput sequencing studies accumulate, it becomes more evident that the microbial transfer between plants and insects is much more extensive than previously thought. Therefore, studies similar to the ones performed




FIG 1 Microbiome exchange and the interactions between microbial communities of plants and insect herbivores. The microbial interaction between each species of plant and insect is likely highly specific and dependent on the insect’s feeding mode and environmental conditions. Endophytic fungi may play an important role by producing secondary metabolites that are toxic to the insect herbivores. However, together with endophytic bacteria, they can also become part of the insect gut microbiome. Experiments revealing the microbial community structures will enable the identification of the drivers of diversity and the indicator strains for increasing or reducing the fitness of each host as well as the further manipulation of the microbiome exchange.

by Lòpez-Fernàndez et al. (56) on the bacterial communities in the sap-feeding grasshopper and grapevine, by Mogouong et al. (100) on the microbial communities in the emerald ash borer and ash trees, by Šigut et al. (102) on different caterpillar and tree species, and by Gonzalez-Escobedo et al. (101) on the bacterial communities in the bark beetle and pine trees, should become mainstream. Such studies would provide information on the involvement of microbial symbionts in the performance of an herbivore on a specific plant species and in a given environment, such as forest ecosystems. For example, the expansion of many forest pests, such as the invasive woodwasp (*Sirex noctilio* Fabricius) and spruce sawfly (*Gilpinia hercyniae* Hartig) has thus far been explained by the lack of parasitoids in the new environment (131, 132). However, the exchange of microbial communities with new host plants may have played an important role in their success in spreading to the new areas.

Studies on the transmission of microbial communities by insects could further revolutionize approaches to plant protection and even plant growth improvement. The analysis of the microbial community structures in each host, plant and insect will enable the identification of the drivers of diversity and the indicator strains for specific

TABLE 1 Microbiome exchange in plant-herbivore interactions^a

Ecosystem	Plant species	Insect species/Feeding mode	Microbial species	Reference(s)
Forest	Douglas fir (<i>Pseudotsuga menziesii</i>)	Needle midge (<i>Contarinia pseudotsugae</i>)/ Internal (gall-making), needle chewing	Endophytic fungus (<i>Rhodocone parkeri</i>)	66–68
Forest	White spruce	Spruce budworm (<i>Choristoneura fumiferana</i>)/External (spun needles), needle chewing	Endophytic fungus (<i>Phialocephala scopiformis</i>)	69–72
Forest	Scots pine (<i>Pinus sylvestris</i> L.)	Pine sawfly (<i>Neodiprion sertifer</i>)/External (needles), chewing	Endophytic fungi	74
Forest	Lodgepole pine (<i>Pinus ponderosa</i>)	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)/Internal (phloem), chewing	Phloem endophyte (<i>Bacillus pumilus</i>)	128
Agriculture, Forest	Many plant species	>700 insect species	Endophytic fungus (<i>Beauveria bassiana</i>)	93
Agriculture, Forest	Many plant species	Several insect species	Endophytic bacterium (<i>Bacillus thuringiensis</i>)	91
Agriculture	Grapevine (<i>Vitis vinifera</i> L.)	Leafhopper (<i>Scaphoideus titanus</i>)/ External, sucking (sap)	Endophytic bacteria	56
Forest	Ash (<i>Fraxinus</i> spp.)	Emerald ash borer (<i>Agrilus planipennis</i>)/ Internal (phloem, wood), chewing	Fungal and bacterial endophytes	100
Forest	Arizona pine (<i>Pinus arizonica</i>), Durango pine (<i>Pinus durangensis</i>)	Bark beetle (<i>Dendroctonus rhizophagus</i>)/ Internal (phloem), chewing	Endophytic bacteria	101
Agriculture	Strawberry (<i>Fragaria ananassa</i>)	Caterpillar (<i>Vanessa cardui</i>)/External (leaves), chewing	Epiphytic and endophytic bacteria and fungi	106
Forest	Conifers (several species)	Bark beetles (Coleoptera: Curculionidae: Scolytinae)/Internal (phloem), chewing	Insect symbiotic fungi (Ophiostomatales)	82
Agriculture	Bittercress (<i>Cardamine hirsute</i>)	Leaf-mining fly (<i>Scaptomyza nigrita</i>)/ Internal (leaf mining), chewing	Epiphytic and endophytic bacteria	103
Forest	Pinaceae	Red turpentine beetle (<i>Dendroctonus valens</i>)/Internal (phloem), chewing	Insect symbiotic fungi	81
Forest	Mountain pine (<i>Pinus contorta</i> , <i>Pinus contorta</i> - <i>Pinus banksiana</i>)	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)/Internal (phloem), chewing	Insect symbiotic bacteria (<i>Pseudomonas</i> and <i>Rahnella</i>)	33
Forest	Norway spruce (<i>Picea abies</i>)	European spruce bark beetle (<i>Ips typographus</i>)/Internal (phloem), chewing	Insect symbiotic fungus (<i>Ceratocystis polonica</i>)	87

^aLight grey, positive, dark grey, negative; white, neutral effect on host.

conditions (Fig. 1). The next step could be manipulations of the microbiome exchange. This could take place through the enrichment of species of bacteria or fungi that are beneficial for the plant in that they enhance plant growth, kill pathogens or pests (133), or eliminate plant-pathogenic strains through, for example, phage-based strategies (134, 135). The microbial biocontrol of herbivores would provide an environmentally friendly alternative to chemical pesticides that are, to a great extent, responsible for the global insect decline (17, 135–139). Therefore, research on the interactions between the microbial communities of plants and insects remains highly important for harnessing microbiome exchanges for forest management and crop production in the future.

ACKNOWLEDGMENTS

We thank EU COST Action FA1405 for providing the inspiration for the writing of this article. This work was supported by research grants 343565 and 308766 to A.M.P. and J.J.K. from the Academy of Finland. A.M.P., V.B., P.R.W., A.L.R. and A.M.M. contributed to the conception of the manuscript, and all authors contributed to the writing and review process as well as approved the final version for submission. We declare no conflicts of interest.

REFERENCES

- Hicke JA, Allen CD, Desai AR, Dietze MC, Hall RJ, Ted Hogg EH, Kashian DM, Moore D, Raffa KF, Sturrock RN, Vogelman J. 2012. Effects of biotic disturbances on forest carbon cycling in the United States and Canada. *Glob Change Biol* 18:7–34. <https://doi.org/10.1111/j.1365-2486.2011.02543.x>.

2. Dietze MC, Matthes JH. 2014. A general ecophysiological framework for modelling the impact of pests and pathogens on forest ecosystems. *Ecol Lett* 17:1418–1426. <https://doi.org/10.1111/ele.12345>.
3. Jactel H, Koricheva J, Castagneyrol B. 2019. Responses of forest insect pests to climate change: not so simple. *Curr Opin Insect Sci* 35:103–108. <https://doi.org/10.1016/j.cois.2019.07.010>.
4. Hammer TJ, Bowers MD. 2015. Gut microbes may facilitate insect herbivory of chemically defended plants. *Oecologia* 179:1–14. <https://doi.org/10.1007/s00442-015-3327-1>.
5. Müller DB, Vogel C, Bai Y, Vorholt JA. 2016. The plant microbiota: systems-level insights and perspectives. *Annu Rev Genet* 50:211–234. <https://doi.org/10.1146/annurev-genet-120215-034952>.
6. Hardoim PR, van Overbeek LS, Berg G, Pirttilä AM, Compant S, Campisano A, Döring M, Sessitsch A. 2015. The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol Mol Biol Rev* 79:293–320. <https://doi.org/10.1128/MMBR.00050-14>.
7. Leach JE, Triplett LR, Argueso CT, Trivedi P. 2017. Communication in the phytobiome. *Cell* 169:587–596. <https://doi.org/10.1016/j.cell.2017.04.025>.
8. Rodríguez RJ, White JF, Jr, Arnold AE, Redman RS. 2009. Fungal endophytes: diversity and functional roles. *New Phytologist* 182:314–330. <https://doi.org/10.1111/j.1469-8137.2009.02773.x>.
9. Müller MM, Hallaksela A-M. 1998. Diversity of Norway spruce needle endophytes in various mixed and pure Norway spruce stands. *Mycol Res* 102:1183–1189. <https://doi.org/10.1017/S0953756298006285>.
10. Müller MM, Valjakka R, Suokko A, Hantula J. 2001. Diversity of endophytic fungi of single Norway spruce needles and their role as pioneer decomposers. *Mol Ecol* 10:1801–1810. <https://doi.org/10.1046/j.1365-294X.2001.01304.x>.
11. Douglas AE. 2015. Multiorganismal insects: diversity and function of resident microorganisms. *Annu Rev Entomol* 60:17–34. <https://doi.org/10.1146/annurev-ento-010814-020822>.
12. Giron D, Dedeine F, Dubreuil G, Huguet E, Mouton L, Outreman Y, Vavre F, Simon J-C. 2017. Chapter seven - influence of microbial symbionts on plant-insect interactions, p 225–257. *In* Sauvion N, Thiéry D, Calatayud P-A (ed), *Advances in Botanical Research*. Academic Press.
13. Ren C, Webster P, Finkel SE, Tower J. 2007. Increased internal and external bacterial load during *Drosophila* aging without life-span trade-off. *Cell Metab* 6:144–152. <https://doi.org/10.1016/j.cmet.2007.06.006>.
14. Chen B, Teh B-S, Sun C, Hu S, Lu X, Boland W, Shao Y. 2016. Biodiversity and activity of the gut microbiota across the life history of the insect herbivore *Spodoptera littoralis*. *Sci Rep* 6:29505. <https://doi.org/10.1038/srep29505>.
15. Douglas AE. 2006. Phloem-sap feeding by animals: problems and solutions. *J Exp Bot* 57:747–754. <https://doi.org/10.1093/jxb/erj067>.
16. Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, van Wees SCM, Bakker PAHM. 2014. Induced systemic resistance by beneficial microbes. *Annu Rev Phytopathol* 52:347–375. <https://doi.org/10.1146/annurev-phyto-082712-102340>.
17. Coolen S, Magda Rd-M, Welte CU. 2022. The secret life of insect-associated microbes and how they shape insect-plant interactions. *FEMS Microbiol Ecol* 98:fiac083. <https://doi.org/10.1093/femsec/fiac083>.
18. Chagas FO, Pessotti R de C, Caraballo-Rodríguez AM, Pupo MT. 2018. Chemical signaling involved in plant-microbe interactions. *Chem Soc Rev* 47:1652–1704. <https://doi.org/10.1039/c7cs00343a>.
19. Chaudhary R, Atamian HS, Shen Z, Briggs SP, Kaloshian I. 2014. GroEL from the endosymbiont *Buchnera aphidicola* betrays the aphid by triggering plant defense. *Proc Natl Acad Sci U S A* 111:8919–8924. <https://doi.org/10.1073/pnas.1407687111>.
20. Pangesti N, Reichelt M, van de Mortel JE, Kapsomenou E, Gershenzon J, van Loon JJA, Dicke M, Pineda A. 2016. Jasmonic acid and ethylene signaling pathways regulate glucosinolate levels in plants during rhizobacteria-induced systemic resistance against a leaf-chewing herbivore. *J Chem Ecol* 42:1212–1225. <https://doi.org/10.1007/s10886-016-0787-7>.
21. Conrath U, Beckers GJM, Flors V, García-Agustín P, Jakab G, Mauch F, Newman M-A, Pieterse CMJ, Poinssot B, Pozo MJ, Pugin A, Schaffrath U, Ton J, Wendehenne D, Zimmerli L, Mauch-Mani B. 2006. Priming: getting ready for battle. *Mol Plant Microbe Interact* 19:1062–1071. <https://doi.org/10.1094/MPMI-19-1062>.
22. Wari D, Kabir MA, Mujiono K, Hojo Y, Shinya T, Tani A, Nakatani H, Galis I. 2019. Honeydew-associated microbes elicit defense responses against brown planthopper in rice. *J Exp Bot* 70:1683–1696. <https://doi.org/10.1093/jxb/erz041>.
23. Bruce TJA. 2015. Interplay between insects and plants: dynamic and complex interactions that have coevolved over millions of years but act in milliseconds. *J Exp Bot* 66:455–465. <https://doi.org/10.1093/jxb/eru391>.
24. Ali JG, Agrawal AA. 2012. Specialist versus generalist insect herbivores and plant defense. *Trends Plant Sci* 17:293–302. <https://doi.org/10.1016/j.tplants.2012.02.006>.
25. Novotny V, Miller SE, Baje L, Balagawi S, Basset Y, Cizek L, Craft KJ, Dem F, Drew RAJ, Hulcr J, Leps J, Lewis OT, Pokon R, Stewart AJA, Samuelson GA, Weiblen GD. 2010. Guild-specific patterns of species richness and host specialization in plant-herbivore food webs from a tropical forest. *J Anim Ecol* 79:1193–1203. <https://doi.org/10.1111/j.1365-2656.2010.01728.x>.
26. Jones OT, Coaker TH. 1978. A basis for host plant finding in phytophagous larvae. *Entomol Exp Appl* 24:472–484. <https://doi.org/10.1111/j.1570-7458.1978.tb02807.x>.
27. Soler R, Erb M, Kaplan I. 2013. Long distance root-shoot signalling in plant-insect community interactions. *Trends Plant Sci* 18:149–156. <https://doi.org/10.1016/j.tplants.2012.08.010>.
28. Jander G, Cui J, Nhan B, Pierce NE, Ausubel FM. 2001. The TASTY locus on chromosome 1 of *Arabidopsis* affects feeding of the insect herbivore *Trichoplusia ni*. *Plant Physiol* 126:890–898. <https://doi.org/10.1104/pp.126.2.890>.
29. Züst T, Agrawal AA. 2017. Trade-offs between plant growth and defense against insect herbivory: an emerging mechanistic synthesis. *Annu Rev Plant Biol* 68:513–534. <https://doi.org/10.1146/annurev-arplant-042916-040856>.
30. Gatehouse JA. 2002. Plant resistance towards insect herbivores: a dynamic interaction. *New Phytol* 156:145–169. <https://doi.org/10.1046/j.1469-8137.2002.00519.x>.
31. Howe GA, Jander G. 2008. Plant immunity to insect herbivores. *Annu Rev Plant Biol* 59:41–66. <https://doi.org/10.1146/annurev-arplant.59.032607.092825>.
32. Shavit R, Ofek-Lalzar M, Burdman S, Morin S. 2013. Inoculation of tomato plants with rhizobacteria enhances the performance of the phloem-feeding insect *Bemisia tabaci*. *Front Plant Sci* 4:306. <https://doi.org/10.3389/fpls.2013.00306>.
33. Adams AS, Aylward FO, Adams SM, Erbilgin N, Aukema BH, Currie CR, Suen G, Raffa KF. 2013. Mountain pine beetles colonizing historical and naive host trees are associated with a bacterial community highly enriched in genes contributing to terpene metabolism. *Appl Environ Microbiol* 79:3468–3475. <https://doi.org/10.1128/AEM.00068-13>.
34. Mason CJ, Rubert-Nason KF, Lindroth RL, Raffa KF. 2015. Aspen defense chemicals influence midgut bacterial community composition of gypsy moth. *J Chem Ecol* 41:75–84. <https://doi.org/10.1007/s10886-014-0530-1>.
35. Carvalhais LC, Dennis PG, Badri DV, Tyson GW, Vivanco JM, Schenk PM. 2013. Activation of the jasmonic acid plant defence pathway alters the composition of rhizosphere bacterial communities. *PLoS One* 8:e56457. <https://doi.org/10.1371/journal.pone.0056457>.
36. Kong HG, Kim BK, Song GC, Lee S, Ryu C-M. 2016. Aboveground whitefly infestation-mediated reshaping of the root microbiota. *Front Microbiol* 7:1314. <https://doi.org/10.3389/fmicb.2016.01314>.
37. de Souza EM, Granada CE, Sperotto RA. 2016. Plant pathogens affecting the establishment of plant-symbiont interaction. *Front Plant Sci* 7:15. <https://doi.org/10.3389/fpls.2016.00015>.
38. Hirsch AM, Lum MR, Downie JA. 2001. What makes the rhizobia-legume symbiosis so special? *Plant Physiol* 127:1484–1492. <https://doi.org/10.1104/pp.010866>.
39. Dean JM, Mescher MC, de Moraes CM. 2014. Plant dependence on rhizobia for nitrogen influences induced plant defenses and herbivore performance. *Int J Mol Sci* 15:1466–1480. <https://doi.org/10.3390/ijms15011466>.
40. Pölme S, Bahram M, Kóljalg U, Tedersoo L. 2014. Global biogeography of *Alnus*-associated *Frankia* actinobacteria. *New Phytol* 204:979–988. <https://doi.org/10.1111/nph.12962>.
41. Ballhorn DJ, Elias JD, Balkan MA, Fordyce RF, Kennedy PG. 2017. Colonization by nitrogen-fixing *Frankia* bacteria causes short-term increases in herbivore susceptibility in red alder (*Alnus rubra*) seedlings. *Oecologia* 184:497–506. <https://doi.org/10.1007/s00442-017-3888-2>.
42. Vincent M, Boubakri H, Gasser M, Hay A-E, Herrera-Belaroussi A. 2023. What contribution of plant immune responses in *Alnus glutinosa*-*Frankia* symbiotic interactions? *Symbiosis*:1–26.
43. Goswami D, Thakker JN, Dhandhukia PC. 2016. Portraying mechanics of plant growth promoting rhizobacteria (PGPR): a review. *Cogent Food & Agriculture* 2.
44. Zebelo S, Song Y, Kloepper JW, Fadamiro H. 2016. Rhizobacteria activates (+)- δ -cadinene synthase genes and induces systemic resistance in cotton against beet armyworm (*Spodoptera exigua*). *Plant Cell Environ* 39:935–943. <https://doi.org/10.1111/pce.12704>.

45. Bell K, Naranjo-Guevara N, dos Santos RC, Meadow R, Bento JMS. 2020. Predatory earwigs are attracted by herbivore-induced plant volatiles linked with plant growth-promoting rhizobacteria. *Insects* 11:271. <https://doi.org/10.3390/insects11050271>.
46. Friman J, Pineda A, Gershenzon J, Dicke M, Loon JJA. 2021. Differential effects of the rhizobacterium *Pseudomonas simiae* on above- and belowground chewing insect herbivores. *J Appl Entomol* 145:250–260. <https://doi.org/10.1111/jen.12842>.
47. Koricheva J, Gange AC, Jones T. 2009. Effects of mycorrhizal fungi on insect herbivores: a meta-analysis. *Ecology* 90:2088–2097. <https://doi.org/10.1890/08-1555.1>.
48. Brundrett MC. 2017. Global diversity and importance of mycorrhizal and nonmycorrhizal plants, p 533–556. *In* Tedersoo L (ed), *Biogeography of Mycorrhizal Symbiosis*. Springer International Publishing, Cham.
49. Jakobsen I, Smith SE, Smith FA. 2003. Function and diversity of arbuscular mycorrhizae in carbon and mineral nutrition, p 75–92. *In* van der Heijden MGA, Sanders IR (ed), *Mycorrhizal Ecology*. Springer Berlin Heidelberg, Berlin, Heidelberg.
50. Kempel A, Nater P, Fischer M, Kleunen M. 2013. Plant-microbe-herbivore interactions in invasive and non-invasive alien plant species. *Funct Ecol* 27:498–508. <https://doi.org/10.1111/1365-2435.12056>.
51. Bennett AE, Bever JD, Deane BM. 2009. Arbuscular mycorrhizal fungal species suppress inducible plant responses and alter defensive strategies following herbivory. *Oecologia* 160:771–779. <https://doi.org/10.1007/s00442-009-1338-5>.
52. Saikkonen K, Faeth SH, Helander M, Sullivan TJ. 1998. Fungal endophytes: a continuum of interactions with host plants. *Annu Rev Ecol Syst* 29:319–343. <https://doi.org/10.1146/annurev.ecolsys.29.1.319>.
53. Ruotsalainen AL, Kauppinen M, Wäli PR, Saikkonen K, Helander M, Tuomi J. 2022. Dark septate endophytes: mutualism from by-products? *Trends Plant Sci* 27:247–254. <https://doi.org/10.1016/j.tplants.2021.10.001>.
54. Koivusaari P, Pohjanen J, Wäli PR, Ahonen SHK, Saravesi K, Markkola AM, Haapala K, Suokas M, Koskimäki JJ, Tejesvi MV, Pirttilä AM. 2018. Different endophyte communities colonize buds of sprouts compared with mature trees of mountain birch recovered from moth herbivory. *Tree Physiol* 38:1437–1444. <https://doi.org/10.1093/treephys/tpy012>.
55. Arnold E, Lewis LC. 2005. Ecology and evolution of fungal endophytes, and their roles against insects, p 74–96. *In* Vega FE, Blackwell M (ed), *Insect-Fungal Associations: Ecology and Evolution*. Oxford University Press, New York.
56. López-Fernández S, Mazzoni V, Pedrazzoli F, Pertot I, Campisano A. 2017. A phloem-feeding insect transfers bacterial endophytic communities between grapevine plants. *Front Microbiol* 8:834. <https://doi.org/10.3389/fmicb.2017.00834>.
57. Bacon CW, Porter JK, Robbins JD, Luttrell ES. 1977. Epichloë typhina from toxic tall fescue grasses. *Appl Environ Microbiol* 34:576–581. <https://doi.org/10.1128/aem.34.5.576-581.1977>.
58. Faeth SH. 2002. Are endophytic fungi defensive plant mutualists? *Oikos* 98:25–36. <https://doi.org/10.1034/j.1600-0706.2002.980103.x>.
59. Clay K. 1996. Interactions among fungal endophytes, grasses and herbivores. *Popul Ecol* 38:191–201. <https://doi.org/10.1007/BF02515727>.
60. Saikkonen K, Saari S, Helander M. 2010. Defensive mutualism between plants and endophytic fungi? *Fungal Divers* 41:101–113. <https://doi.org/10.1007/s13225-010-0023-7>.
61. Jaber LR, Vidal S. 2010. Fungal endophyte negative effects on herbivory are enhanced on intact plants and maintained in a subsequent generation. *Ecol Entomol* 35:25–36. <https://doi.org/10.1111/j.1365-2311.2009.01152.x>.
62. Härrilä SA, Krauss J, Müller CB. 2008. Fungal endosymbionts of plants reduce lifespan of an aphid secondary parasitoid and influence host selection. *Proc Biol Sci* 275:2627–2632. <https://doi.org/10.1098/rspb.2008.0594>.
63. Härrilä SA, Krauss J, Müller CB. 2008. Trophic cascades initiated by fungal plant endosymbionts impair reproductive performance of parasitoids in the second generation. *Oecologia* 157:399–407. <https://doi.org/10.1007/s00442-008-1094-y>.
64. de Sassi C, Müller CB, Krauss J. 2006. Fungal plant endosymbionts alter life history and reproductive success of aphid predators. *Proc Biol Sci* 273:1301–1306. <https://doi.org/10.1098/rspb.2005.3442>.
65. Pirttilä AM, Wäli P. 2009. Conifer endophytes, p 235–246. *In* White J, Torres MS (ed), *Defensive Mutualism in Microbial Symbiosis*. CRC Press, Boca Raton, FL, USA.
66. Carroll GC, Carroll FE. 1978. Studies on the incidence of coniferous needle endophytes in the Pacific Northwest. *Can J Bot* 56:3034–3043. <https://doi.org/10.1139/b78-367>.
67. Sherwood-Pike M, Stone JK, Carroll GC. 1986. *Rhabdocline parkeri*, a ubiquitous foliar endophyte of Douglas-fir. *Can J Bot* 64:1849–1855. <https://doi.org/10.1139/b86-245>.
68. Miller J. 1986. Toxins of endophytic and epiphytic fungi of conifer needles, p 223–231. *In* Fokkema NJ, Van Huevel J (ed), *Microbiology of the phyllosphere*. Cambridge University.
69. Miller JD, Mackenzie S, Foto M, Adams GW, Findlay JA. 2002. Needles of white spruce inoculated with rugulosin-producing endophytes contain rugulosin reducing spruce budworm growth rate. *Mycol Res* 106:471–479. <https://doi.org/10.1017/S0953756202005671>.
70. Miller JD, Sumarah MW, Adams GW. 2008. Effect of a rugulosin-producing endophyte in *Picea glauca* on *Choristoneura fumiferana*. *J Chem Ecol* 34:362–368. <https://doi.org/10.1007/s10886-007-9412-0>.
71. Sumarah MW, Adams GW, Berghout J, Slack GJ, Wilson AM, Miller JD. 2008. Spread and persistence of a rugulosin-producing endophyte in *Picea glauca* seedlings. *Mycol Res* 112:731–736. <https://doi.org/10.1016/j.mycres.2008.01.007>.
72. Sumarah MW, Miller JD. 2009. Anti-insect secondary metabolites from fungal endophytes of conifer trees. *Nat Prod Commun* 4:1934578X0900401–1934578X0900401.
73. Sumarah MW, Puniani E, Sørensen D, Blackwell BA, Miller JD. 2010. Secondary metabolites from anti-insect extracts of endophytic fungi isolated from *Picea rubens*. *Phytochemistry* 71:760–765. <https://doi.org/10.1016/j.phytochem.2010.01.015>.
74. Saikkonen K, Helander M, Ranta H, Neuvonen S, Virtanen T, Suomela J, Vuorinen P. 1996. Endophyte-mediated interactions between woody plants and insect herbivores? *Entomol Exp Appl* 80:269–271. <https://doi.org/10.1111/j.1570-7458.1996.tb00932.x>.
75. Ahlholm J, Helander M, Elamo P, Saloniemi I, Neuvonen S, Hanhimäki S, Saikkonen K. 2002. Micro-fungi and invertebrate herbivores on birch trees: fungal mediated plant-herbivore interactions or responses to host quality? *Ecol Lett* 5:648–655. <https://doi.org/10.1046/j.1461-0248.2002.00368.x>.
76. Agrios G. 2004. *Plant Pathology: Fifth Edition* 1–922.
77. Six DL. 2003. Bark beetle-fungus symbioses, p 97–114. *In* Bourtzis K, Miller TA (ed), *Insect Symbiosis*. CRC Press, New York.
78. Mann RS, Pelz-Stelinski K, Hermann SL, Tiwari S, Stelinski LL. 2011. Sexual transmission of a plant pathogenic bacterium, *Candidatus Liberibacter asiaticus*, between conspecific insect vectors during mating. *PLoS One* 6:e29197. <https://doi.org/10.1371/journal.pone.0029197>.
79. Galambos N, Compant S, Wäckers F, Sessitsch A, Anfora G, Mazzoni V, Pertot I, Perazzoli M. 2021. Beneficial insects deliver plant growth-promoting bacterial endophytes between tomato plants. *Microorganisms* 9:1294. <https://doi.org/10.3390/microorganisms9061294>.
80. Pronk LUJ, Bakker PAHM, Keel C, Maurhofer M, Flury P. 2022. The secret life of plant-beneficial rhizosphere bacteria: insects as alternative hosts. *Environ Microbiol* 24:3273–3289. <https://doi.org/10.1111/1462-2920.15968>.
81. Taerum SJ, Duong TA, de Beer ZW, Gillette N, Sun J-H, Owen DR, Wingfield MJ. 2013. Large shift in symbiont assemblage in the invasive red turpentine beetle. *PLoS One* 8:e78126. <https://doi.org/10.1371/journal.pone.0078126>.
82. Paine TD, Raffa KF, Harrington TC. 1997. Interactions among scolytid bark beetles, their associated fungi, and live host conifers. *Annu Rev Entomol* 42:179–206. <https://doi.org/10.1146/annurev.ento.42.1.179>.
83. Kirisits T. 2004. Fungal associates of European bark beetles with special emphasis on the ophiostomatoid fungi, p 181–236. *In* *Bark and Wood Boring Insects in Living Trees in Europe: A Synthesis*.
84. Whitney HS. 1982. Relationships between bark beetles and symbiotic organisms, p 183–211. *In* Mitton JB, Sturgeon KB (ed), *Bark Beetles in North American Conifers*. University of Texas Press, Austin.
85. Six DL, Paine TD. 1998. Effects of mycorrhizal fungi and host tree species on progeny survival and emergence of *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Environ Entomol* 27:1393–1401. <https://doi.org/10.1093/ee/27.6.1393>.
86. Lieutier F, Yart A, Salle A. 2009. Stimulation of tree defenses by ophiostomatoid fungi can explain attack success of bark beetles on conifers. *Ann For Sci* 66:801–801. <https://doi.org/10.1051/forest/2009066>.
87. Hammerbacher A, Schmidt A, Wadke N, Wright LP, Schneider B, Bohlmann J, Brand WA, Fenning TM, Gershenzon J, Paetz C. 2013. A common fungal associate of the spruce bark beetle metabolizes the stilbene defenses of Norway spruce. *Plant Physiol* 162:1324–1336. <https://doi.org/10.1104/pp.113.218610>.
88. Kandasamy D, Gershenzon J, Andersson MN, Hammerbacher A. 2019. Volatile organic compounds influence the interaction of the Eurasian

- spruce bark beetle (*Ips typographus*) with its fungal symbionts. *ISME J* 13:1788–1800. <https://doi.org/10.1038/s41396-019-0390-3>.
89. Kandasamy D, Gershenzon J, Hammerbacher A. 2016. Volatile organic compounds emitted by fungal associates of conifer bark beetles and their potential in bark beetle control. *J Chem Ecol* 42:952–969. <https://doi.org/10.1007/s10886-016-0768-x>.
 90. Chucho J, Auricau-Bouverly N, Danet J-L, Thiéry D. 2017. Use the insiders: could insect facultative symbionts control vector-borne plant diseases? *J Pest Sci* 90:51–68. <https://doi.org/10.1007/s10340-016-0782-3>.
 91. Monnerat RG, Soares CM, Capdeville G, Jones G, Martins ES, Praça L, Cordeiro BA, Braz SV, dos Santos RC, Berry C. 2009. Translocation and insecticidal activity of *Bacillus thuringiensis* living inside of plants. *Microb Biotechnol* 2:512–520. <https://doi.org/10.1111/j.1751-7915.2009.00116.x>.
 92. Caspi-Fluger A, Zchori-Fein E. 2010. Do plants and insects share the same symbionts? *Isr J Plant Sci* 58:113–119. <https://doi.org/10.1560/IJPS.58.2.113>.
 93. Ortiz-Urquiza A. 2021. The split personality of *Beauveria bassiana*: understanding the molecular basis of fungal parasitism and mutualism. *mSystems* 6:e00766-21. <https://doi.org/10.1128/mSystems.00766-21>.
 94. Vega FE. 2018. The use of fungal entomopathogens as endophytes in biological control: a review. *Mycologia* 110:4–30. <https://doi.org/10.1080/00275514.2017.1418578>.
 95. Shikano I, Rosa C, Tan C-W, Felton GW. 2017. Tritrophic interactions: microbe-mediated plant effects on insect herbivores. *Annu Rev Phytopathol* 55:313–331. <https://doi.org/10.1146/annurev-phyto-080516-035319>.
 96. Strano CP, Malacrinò A, Campolo O, Palmeri V. 2018. Influence of host plant on *Thaumetopoea pityocampa* gut bacterial community. *Microb Ecol* 75:487–494. <https://doi.org/10.1007/s00248-017-1019-6>.
 97. Landry M, Comeau AM, Derome N, Cusson M, Levesque RC. 2015. Composition of the spruce budworm (*Choristoneura fumiferana*) midgut microbiota as affected by rearing conditions. *PLoS One* 10:e0144077. <https://doi.org/10.1371/journal.pone.0144077>.
 98. Broderick NA, Raffa KF, Goodman RM, Handelsman J. 2004. Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Appl Environ Microbiol* 70:293–300. <https://doi.org/10.1128/AEM.70.1.293-300.2004>.
 99. Chung SH, Scully ED, Peiffer M, Geib SM, Rosa C, Hoover K, Felton GW. 2017. Host plant species determines symbiotic bacterial community mediating suppression of plant defenses. *Sci Rep* 7:39690. <https://doi.org/10.1038/srep39690>.
 100. Mogouong J, Constant P, Legendre P, Guertin C. 2021. The phyllosphere microbiome of host trees contributes more than leaf phytochemicals to variation in the *Agrilus planipennis* Fairmaire gut microbiome structure. *Sci Rep* 11:15911. <https://doi.org/10.1038/s41598-021-95146-9>.
 101. Gonzalez-Escobedo R, Briones-Roblero CI, Pineda-Mendoza RM, Rivera-Orduña FN, Zúñiga G. 2018. Bacteriome from *Pinus arizonica* and *P. durangensis*: diversity, comparison of assemblages, and overlapping degree with the gut bacterial community of a bark beetle that kills pines. *Front Microbiol* 9:77. <https://doi.org/10.3389/fmicb.2018.00077>.
 102. Šigut M, Pyszko P, Šigutová H, Vlišnovská D, Kostovčík M, Kotásková N, Dornák O, Kolařík M, Drozd P. 2022. Fungi are more transient than bacteria in caterpillar gut microbiomes. *Sci Rep* 12:15552. <https://doi.org/10.1038/s41598-022-19855-5>.
 103. Humphrey PT, Whiteman NK. 2020. Insect herbivory reshapes a native leaf microbiome. *Nat Ecol Evol* 4:221–229. <https://doi.org/10.1038/s41599-019-1085-x>.
 104. Smee MR, Hendry TA. 2022. Context-dependent benefits of aphids for bacteria in the phyllosphere. *Am Nat* 199:380–392. <https://doi.org/10.1086/718264>.
 105. Müller T, Müller M, Behrendt U, Stadler B. 2003. Diversity of culturable phyllosphere bacteria on beech and oak: the effects of lepidopterous larvae. *Microbiol Res* 158:291–297. <https://doi.org/10.1078/0944-5013-00207>.
 106. Wei N, Whyte RL, Ashman T, Jamieson MA. 2022. Genotypic variation in floral volatiles influences floral microbiome more strongly than interactions with herbivores and mycorrhizae in strawberries. *Hortic Res* 9:uhab005. <https://doi.org/10.1093/hr/uhab005>.
 107. Feldhaar H. 2011. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecol Entomol* 36:533–543. <https://doi.org/10.1111/j.1365-2311.2011.01318.x>.
 108. Sun J, Lu M, Gillette NE, Wingfield MJ. 2013. Red turpentine beetle: innocuous native becomes invasive tree killer in China. *Annu Rev Entomol* 58:293–311. <https://doi.org/10.1146/annurev-ento-120811-153624>.
 109. Acuña R, Padilla BE, Flórez-Ramos CP, Rubio JD, Herrera JC, Benavides P, Lee S-J, Yeats TH, Egan AN, Doyle JJ, Rose JKC. 2012. Adaptive horizontal transfer of a bacterial gene to an invasive insect pest of coffee. *Proc Natl Acad Sci U S A* 109:4197–4202. <https://doi.org/10.1073/pnas.1121190109>.
 110. Ferrater JB, de Jong PW, Dicke M, Chen YH, Horgan FG. 2013. Symbiont-mediated adaptation by planthoppers and leafhoppers to resistant rice varieties. *Arthropod Plant Interact* 7:591–605. <https://doi.org/10.1007/s11829-013-9277-9>.
 111. Hansen AK, Moran NA. 2014. The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* 23:1473–1496. <https://doi.org/10.1111/mec.12421>.
 112. Adams AS, Jordan MS, Adams SM, Suen G, Goodwin LA, Davenport KW, Currie CR, Raffa KF. 2011. Cellulose-degrading bacteria associated with the invasive woodwasp *Sirex noctilio*. *ISME J* 5:1323–1331. <https://doi.org/10.1038/ismej.2011.14>.
 113. Delalibera I, Jr, Handelsman JO, Raffa KF. 2005. Contrasts in cellulolytic activities of gut microorganisms between the wood borer, *Saperda vestita* (Coleoptera: Cerambycidae), and the bark beetles, *Ips pini* and *Dendroctonus frontalis* (Coleoptera: Curculionidae). *Environ Entomol* 34:541–547. <https://doi.org/10.1603/0046-225X-34.3.541>.
 114. Vilanova C, Baixeras J, Latorre A, Porcar M. 2016. The generalist inside the specialist: gut bacterial communities of two insect species feeding on toxic plants are dominated by *Enterococcus* sp. *Front Microbiol* 7:1005. <https://doi.org/10.3389/fmicb.2016.01005>.
 115. Xia X, Gurr GM, Vasseur L, Zheng D, Zhong H, Qin B, Lin J, Wang Y, Song F, Li Y, Lin H, You M. 2017. Metagenomic sequencing of diamondback moth gut microbiome unveils key holobiont adaptations for herbivory. *Front Microbiol* 8:663. <https://doi.org/10.3389/fmicb.2017.00663>.
 116. Visóto LE, Oliveira MGA, Guedes RNC, Ribon AOB, Good-God PIV. 2009. Contribution of gut bacteria to digestion and development of the velvetbean caterpillar, *Anticarsia gemmatilis*. *J Insect Physiol* 55:185–191. <https://doi.org/10.1016/j.jinsphys.2008.10.017>.
 117. Scully ED, Geib SM, Carlson JE, Tien M, McKenna D, Hoover K. 2014. Functional genomics and microbiome profiling of the Asian longhorned beetle (*Anoplophora glabripennis*) reveal insights into the digestive physiology and nutritional ecology of wood feeding beetles. *BMC Genomics* 15:1–21. <https://doi.org/10.1186/1471-2164-15-1096>.
 118. Scully ED, Geib SM, Mason CJ, Carlson JE, Tien M, Chen H-Y, Harding S, Tsai C-J, Hoover K. 2018. Host-plant induced changes in microbial community structure and midgut gene expression in an invasive polyphage (*Anoplophora glabripennis*). *Sci Rep* 8:1–16. <https://doi.org/10.1038/s41598-018-27476-0>.
 119. Ayayee PA, Larsen T, Rosa C, Felton GW, Ferry JG, Hoover K. 2016. Essential amino acid supplementation by gut microbes of a wood-feeding cerambycid. *Environ Entomol* 45:66–73. <https://doi.org/10.1093/ee/nvv153>.
 120. Morales-Jiménez J, Vera-Ponce de León A, García-Domínguez A, Martínez-Romero E, Zúñiga G, Hernández-Rodríguez C. 2013. Nitrogen-fixing and uricolytic bacteria associated with the gut of *Dendroctonus rhizophagus* and *Dendroctonus valens* (Curculionidae: Scolytinae). *Microb Ecol* 66:200–210. <https://doi.org/10.1007/s00248-013-0206-3>.
 121. Gonella E, Tedeschi R, Crotti E, Alma A. 2019. Multiple guests in a single host: interactions across symbiotic and phytopathogenic bacteria in phloem-feeding vectors – a review. *Entomol Exp Appl* 167:171–185. <https://doi.org/10.1111/eea.12766>.
 122. Wang J, Chung SH, Peiffer M, Rosa C, Hoover K, Zeng R, Felton GW. 2016. Herbivore oral secreted bacteria trigger distinct defense responses in preferred and non-preferred host plants. *J Chem Ecol* 42:463–474. <https://doi.org/10.1007/s10886-016-0712-0>.
 123. Chung SH, Rosa C, Scully ED, Peiffer M, Tooker JF, Hoover K, Luthe DS, Felton GW. 2013. Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc Natl Acad Sci U S A* 110:15728–15733. <https://doi.org/10.1073/pnas.1308867110>.
 124. Giron D, Frago E, Glevarec G, Pieterse CMJ, Dicke M. 2013. Cytokinins as key regulators in plant–microbe–insect interactions: connecting plant growth and defence. *Funct Ecol* 27:599–609. <https://doi.org/10.1111/1365-2435.12042>.
 125. Derksen H, Rampitsch C, Daayf F. 2013. Signaling cross-talk in plant disease resistance. *Plant Science* 207:79–87. <https://doi.org/10.1016/j.plantsci.2013.03.004>.
 126. Mauck KE, de Moraes CM, Mescher MC. 2016. Effects of pathogens on sensory-mediated interactions between plants and insect vectors. *Curr Opin Plant Biol* 32:53–61. <https://doi.org/10.1016/j.pbi.2016.06.012>.
 127. Martemyanov VV, Belousova IA, Pavlushin S, Dubovskiy IM, Ershov NI, Alikina TY, Kabilov MR, Glupov V. 2016. Phenological asynchrony between host plant and gypsy moth reduces insect gut microbiota and susceptibility

- to *Bacillus thuringiensis*. *Ecol Evol* 6:7298–7310. <https://doi.org/10.1002/ece3.2460>.
128. Adams AS, Six DL, Adams SM, Holben WE. 2008. In vitro interactions between yeasts and bacteria and the fungal symbionts of the mountain pine beetle (*Dendroctonus ponderosae*). *Microb Ecol* 56:460–466. <https://doi.org/10.1007/s00248-008-9364-0>.
 129. Campisano A, Pancher M, Puopolo G, Puddu A, Lòpez-Fernàndez S, Biagini B, Yousaf S, Pertot I. 2015. Diversity in endophyte populations reveals functional and taxonomic diversity between wild and domesticated grapevines. *Am J Enol Vitic* 66:12–21. <https://doi.org/10.5344/ajev.2014.14046>.
 130. Gupta A, Nair S. 2020. Dynamics of insect-microbiome interaction influence host and microbial symbiont. *Front Microbiol* 11:1357. <https://doi.org/10.3389/fmicb.2020.01357>.
 131. Kenis M, Hurley BP, Hajek AE, Cock MJW. 2017. Classical biological control of insect pests of trees: facts and figures. *Biol Invasions* 19:3401–3417. <https://doi.org/10.1007/s10530-017-1414-4>.
 132. Kroll SA, Hajek AE, Erin ME, Long SJ. 2013. Parasitism of *Sirex noctilio* by non-sterilizing *Deladenus siricidicola* in northeastern North America. *Biological Control* 67:203–211. <https://doi.org/10.1016/j.biocontrol.2013.08.005>.
 133. Corinne V, Bastien C, Emmanuelle J, Heidy S. 2021. Trees and insects have microbiomes: consequences for forest health and management. *Curr Forestry Rep* 7:81–96. <https://doi.org/10.1007/s40725-021-00136-9>.
 134. Villalpando-Aguilar JL, Matos-Pech G, Lòpez-Rosas I, Castelán-Sánchez HG, Alatorre-Cobos F. 2022. Phage therapy for crops: concepts, experimental and bioinformatics approaches to direct its application. *Int J Mol Sci* 24:325. <https://doi.org/10.3390/ijms24010325>.
 135. Pirttilä AM, Mohammad Parast Tabas H, Baruah N, Koskimäki JJ. 2021. Biofertilizers and biocontrol agents for agriculture: how to identify and develop new potent microbial strains and traits. *Microorganisms* 9:817. <https://doi.org/10.3390/microorganisms9040817>.
 136. Seibold S, Gossner MM, Simons NK, Blüthgen N, Müller J, Ambarlı D, Ammer C, Bauhus J, Fischer M, Habel JC, Linsenmair KE, Naus T, Penone C, Prati D, Schall P, Schulze E-D, Vogt J, Wöllauer S, Weisser WW. 2019. Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature* 574:671–674. <https://doi.org/10.1038/s41586-019-1684-3>.
 137. White JF, Kingsley KL, Zhang Q, Verma R, Obi N, Dvinskikh S, Elmore MT, Verma SK, Gond SK, Kowalski KP. 2019. Review: endophytic microbes and their potential applications in crop management. *Pest Manag Sci* 75:2558–2565. <https://doi.org/10.1002/ps.5527>.
 138. Chaudhary P, Agri U, Chaudhary A, Kumar A, Kumar G. 2022. Endophytes and their potential in biotic stress management and crop production. *Front Microbiol* 13:933017. <https://doi.org/10.3389/fmicb.2022.933017>.
 139. Kamran M, Imran QM, Ahmed MB, Falak N, Khatoon A, Yun B-W. 2022. Endophyte-mediated stress tolerance in plants: a sustainable strategy to enhance resilience and assist crop improvement. *Cells* 11:3292. <https://doi.org/10.3390/cells11203292>.